

Home

2. Tetracycline

Tetracycline ⊲ⁿ

Generic name: tetracycline [*TET-tra-SYE-kleen*]

Brand names: Ala-Tet, Brodspec, Panmycin, Sumycin, Tetracap, ... show all 8 brands

Drug class: Tetracyclines

Medically reviewed by Melisa Puckey, BPharm. Last updated on Feb 29, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is tetracycline?

Tetracycline is an antibiotic that fights infection caused by bacteria.

Tetracycline is used to treat many different bacterial infections of the skin, intestines, respiratory tract, urinary tract, genitals, lymph nodes, and other body systems. It is often used in treating severe acne, or sexually transmitted diseases such as syphilis, gonorrhea, or chlamydia.

Tetracycline is also used to treat infections you can get from direct contact with infected animals or contaminated food.

In some cases, tetracycline is used when penicillin or another antibiotic cannot be used to treat serious infections such as Anthrax, Listeria, Clostridium, Actinomyces, and others.

Warnings

Children younger than 8 years old should not take this medicine.

Using tetracycline during pregnancy could harm the unborn baby or cause permanent tooth discoloration later in the baby's life.

Throw away any unused tetracycline when it expires or when it is no longer needed. Do not take this medicine after the expiration date on the label has passed. Expired tetracycline can cause a dangerous syndrome resulting in damage to the kidneys.

Tetracycline passes into breast milk and may affect bone and tooth development in a nursing baby. Do not take this medication without telling your doctor if you are breast-feeding a baby.

Avoid exposure to sunlight or artificial UV rays (sunlamps or tanning beds). Tetracycline can make your skin more sensitive to sunlight and sunburn may result. Use a sunscreen (minimum SPF 15) and wear protective clothing if you must be out in the sun.

Before taking this medicine

You should not use this medicine if you are allergic to tetracycline or similar medicines such as demeclocycline, doxycycline, minocycline, or tigecycline.

To make sure tetracycline is safe for you, tell your doctor if you have:

- · liver disease; or
- · kidney disease.

If you are using tetracycline to treat gonorrhea, your doctor may test you to make sure you do not also have syphilis, another sexually transmitted disease.

Taking this medicine during pregnancy may affect tooth and bone development in the unborn baby. Taking tetracycline during the last half of pregnancy can cause permanent tooth discoloration later in the baby's life. Tell your doctor if you are pregnant or if you become pregnant while using this medicine.

Tetracycline can make birth control pills less effective. Ask your doctor about using a non-hormonal birth control (condom, diaphragm with spermicide) to prevent pregnancy.

Tetracycline can pass into breast milk and may affect bone and tooth development in a nursing infant. **Do not breast-feed while you are taking tetracycline.**

Children younger than 8 years old should not take tetracycline. Tetracycline can cause permanent tooth discoloration and can also affect a child's growth.

1 Tetracycline pregnancy and breastfeeding warnings (more detail)

How should I take tetracycline?

Take tetracycline exactly as it was prescribed for you. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Take tetracycline on an empty stomach, at least 1 hour before or 2 hours after a meal.

Do not take this medicine with milk or other dairy products, unless your doctor has told you to. Dairy products can make it harder for your body to absorb the medicine.

Shake the oral suspension (**liquid**) well just before you measure a dose. Measure liquid medicine with the dosing syringe provided, or with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

If you need surgery, tell the surgeon ahead of time that you are using tetracycline.

If you use this medicine long-term, you may need frequent medical tests.

Use this medicine for the full prescribed length of time. Your symptoms may improve before the infection is completely cleared. Skipping doses may also increase your risk of further infection that is resistant to antibiotics. Tetracycline will not treat a viral infection such as the flu or a common cold.

Do not share this medicine with another person, even if they have the same symptoms you have.

Store this medication at room temperature away from moisture and heat.

Throw away any unused tetracycline after the expiration date on the label has passed. Taking expired tetracycline can cause damage to your kidneys.

Detailed Tetracycline dosage information

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What to avoid

For at least 2 hours before or 2 hours after taking tetracycline: avoid taking iron supplements, multivitamins, calcium supplements, antacids, or laxatives.

Antibiotic medicines can cause diarrhea, which may be a sign of a new infection. If you have diarrhea that is watery or bloody, call your doctor. Do not use anti-diarrhea medicine unless your doctor tells you to.

Avoid exposure to sunlight or tanning beds. Tetracycline can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Tetracycline side effects

Get emergency medical help if you have signs of an allergic reaction to tetracycline: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- · severe blistering, peeling, and red skin rash;
- · fever, chills, body aches, flu symptoms;
- · pale or yellowed skin, easy bruising or bleeding;
- · any signs of a new infection.

Common tetracycline side effects may include:

- nausea, vomiting, diarrhea, upset stomach, loss of appetite;
- white patches or sores inside your mouth or on your lips;
- swollen tongue, black or "hairy" tongue, trouble swallowing;
- sores or swelling in your rectal or genital area; or
- · vaginal itching or discharge.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

<u>Tetracycline side effects</u> (more detail)

Related/similar drugs

Hydroxychloroquine

Hydroxychloroquine is a quinoline drug used to treat or prevent malaria. It's also used to treat ...

Reviews & ratings

6.4 / 10

356 Reviews

View more

Ciprofloxacin

Ciprofloxacin is an antibiotic belong to a group of drugs called fluoroquinolones. Learn about side ...

Reviews & ratings

5.0 / 10

935 Reviews

View more

FEATURED

Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used alone or with other ...

Reviews & ratings

5.7 / 10

145 Reviews

View more

Metronidazole

Metronidazole is an antibiotic used to fight bacteria in your body. Learn about side effects ...

Reviews & ratings

6.0 / 10

View more

Azithromycin

Azithromycin is an antibiotic used to treat many different types of infections caused by bacteria ...

Reviews & ratings

6.8 / 10

1,306 Reviews

View more

Cephalexin

Cephalexin is a cephalosporin antibiotic used to treat bacterial infections like respiratory, skin ...

Reviews & ratings

6.4 / 10

509 Reviews

View more

Doxycycline

Doxycycline is a tetracycline antibiotic used to treat many different bacterial infections such as ...

Reviews & ratings

6.4 / 10

1,635 Reviews

View more

Pantoprazole

Pantoprazole is a proton pump inhibitor used to treat GERD, erosive esophagitis, and ...

Reviews & ratings

5.0 / 10

430 Reviews

\/:---

Amoxicillin

Amoxicillin is a penicillin antibiotic that fights bacteria. It is used to treat many types of ...

Reviews & ratings

6.4 / 10

366 Reviews

View more

Augmentin

Augmentin is a prescription antibiotic combining amoxicillin and clavulanate to treat bacterial ...

Reviews & ratings

5.4 / 10

447 Reviews

View more

What other drugs will affect tetracycline?

Tell your doctor about all your current medicines and any you start or stop using, especially:

- any other antibiotic;
- isotretinoin (Accutane);
- tretinoin (Renova, Retin-A, Vesanoid);
- a blood thinner (warfarin, Coumadin, Jantoven);
- an antacid or laxative medicine; or
- a vitamin or mineral supplement that contains iron, zinc, calcium, or magnesium.

This list is not complete. Other drugs may interact with tetracycline, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

1 Tetracycline drug interactions (more detail)

Does tetracycline interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

tetracycline + Enter a drug name Add

Popular FAQ

What causes black hairy tongue?

More FAQ

- What are the best antibiotics for pneumonia?
- How long do antibiotics take to work?

View more FAQ...

More about tetracycline

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (64)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Support group
- Drug class: tetracyclines
- Breastfeeding
- En español

Patient resources

• Tetracycline (Systemic) patient information

Other brands

Actisite

Professional resources

• Tetracycline monograph

- Tetracycline (FDA)
- Tetracycline Tablets (FDA)

Other brands

Achromycin V, Sumycin

Related treatment guides

- Bacterial Infection
- Bladder Infection
- Acne
- Bronchitis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use tetracycline only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

Antibiotics 101

List of Common Antibiotics & Types

Antibiotics and Alcohol

Probiotics and Antibiotics

Antibiotics for Strep Throat

Antibiotics for a Tooth Infection

Antibiotics for UTI's

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

∾ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

6.6 / 10

64 Reviews

Images

Tetracycline Hydrochloride 500 mg (5266 5266)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.