

Home

2. Topicort

Topicort ⊲»

Generic name: desoximetasone topical [des-OX-i-MET-a-sone]

Brand names: Topicort, Topicort LP

Drug class: Topical steroids



Medically reviewed by Drugs.com on Dec 17, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Topicort?

Topicort (for the skin) is a potent steroid used to treat inflammation and itching caused by plaque psoriasis or skin conditions that respond to steroid medication.

Topicort may also be used for purposes not listed in this medication guide.

Topicort side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Topicort may cause serious side effects. Call your doctor at once if you have:

- · worsening of your skin condition;
- redness, warmth, swelling, oozing, or severe irritation of any treated skin;
- blurred vision, tunnel vision, eye pain, or seeing halos around lights;
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor; or
- possible signs of absorbing Topicort through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea, diarrhea, tiredness, mood changes, menstrual changes, sexual changes.

Common side effects of Topicort may include:

- rash, itching, or irritation around your mouth;
- burning, irritation, itching, or dryness of treated skin;
- redness or crusting around your hair follicles;
- · increased hair growth;

- blisters, pimples, or crusting of treated skin;
- changes in color of treated skin;
- white or "pruned" appearance of the skin; or
- · stretch marks.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



Topicort side effects (more detail)

Related/similar drugs

Dupixent

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

Reviews & ratings

7.2 / 10

499 Reviews

View more

Stelara

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

Reviews & ratings

7.6 / 10

252 Reviews

View more

FEATURED

Skyrizi

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

Reviews & ratings

7.9 / 10

56 Reviews

View more

Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

Reviews & ratings

7.0 / 10

223 Reviews

View more

Humira

Humira is a tumor necrosis factor blocker that is used to treat many inflammatory conditions such ...

Reviews & ratings

6.4 / 10

698 Reviews

View more

Adalimumab

Adalimumab is used to treat plaque psoriasis, Crohn's disease, ulcerative colitis, rheumatoid ...

Reviews & ratings

6.4 / 10

737 Reviews

View more

Triamcinolone

Triamcinolone is used to treat allergies, skin conditions, ulcerative colitis, and arthritis. Learn ...

Reviews & ratings

6.7 / 10

226 Reviews

View more

Methotrexate

Methotrexate is used to treat certain

types of cancer of the breast, skin, head and neck, or lung ...

Reviews & ratings

6.5 / 10

335 Reviews

View more

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use desoximetasone if you are allergic to it.

Tell your doctor if you have ever had:

- any type of skin infection;
- a skin reaction to any steroid medicine;
- liver disease; or
- an adrenal gland disorder.

Steroid medicines can increase the glucose (sugar) levels in your blood or urine. Tell your doctor if you have diabetes.

It is not known whether Topicort will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk. If you apply desoximetasone to your chest, avoid areas that may come into contact with the baby's mouth.

Topicort is not approved for use by anyone younger than 18 years old. Some brands or forms of this medicine can be used by people younger than 18 years.

Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects.

How should I use Topicort?

Use exactly as directed on the label, or as prescribed by your doctor.

Do not take by mouth. Topical medicine is for use only on the skin. Rinse with water if this medicine gets in your eyes or mouth.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions.

Wash your hands before and after using desoximetasone, unless you are using this medicine to treat the skin on your hands.

Apply a thin layer of medicine to the affected skin and rub it in gently. **Do not apply this medicine over a large area of skin unless your doctor has told you to.**

Do not cover the treated skin area with a bandage or other covering unless your doctor tells you to. Covering treated areas can increase the amount of medicine absorbed through your skin and may cause harmful effects.

If you are treating the diaper area, do not use plastic pants or tight-fitting diapers.

This medicine is for short-term use only (up to 4 weeks for psoriasis). Follow your doctor's dosing instructions very carefully.

If you use desoximetasone to treat plaque psoriasis, you should stop using the medicine once your skin symptoms are controlled.

Call your doctor if your symptoms do not improve after 2 weeks of using this medicine.

You should not stop using desoximetasone suddenly. Follow your doctor's instructions about tapering your dose.

Store at room temperature away from moisture and heat. Keep from freezing.

Desoximetasone **spray** is flammable. Do not use near high heat or open flame. Do not smoke until the gel has completely dried on your skin.

Throw away any Topicort spray not used within 30 days after you received it.

Detailed Topicort dosage information

What happens if I miss a dose?

Apply the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** apply two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

High doses or long-term use of Topicort can lead to thinning skin, easy bruising, changes in body fat (especially in your face, neck, back, and waist), increased acne or facial hair, menstrual problems, impotence, or loss of interest in sex.

What should I avoid while using desoximetasone topical?

Avoid applying Topicort to your face, underarms, or groin area.

Do not use Topicort to treat any skin condition that has not been checked by your doctor.

Avoid using other topical steroid medications on the areas you treat with desoximetasone unless your doctor tells you to.

What other drugs will affect Topicort?

Medicine used on the skin is not likely to be affected by other drugs you use. But many drugs can interact with each other. Tell each of your healthcare providers about all medicines you use, including prescription and over-the-counter medicines, vitamins, and herbal products.

1 Topicort drug interactions (more detail)

Does Topicort interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



More about Topicort (desoximetasone topical)

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (6)
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Drug class: topical steroids
- Breastfeeding
- En español

Patient resources

- · Topicort advanced reading
- Topicort Spray

Other brands

Topicort LP

Professional resources

• Topicort prescribing information

- Topicort Spray (FDA)
- Desoximetasone (AHFS Monograph)

Related treatment guides

- Atopic Dermatitis
- Dermatitis
- Eczema
- Psoriasis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ം Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 1977

User Reviews & Ratings

6 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







×