

Home

2. Meclizine

Meclizine ⊲

Generic name: meclizine [MEK-li-zeen]

Brand names: Antivert, Bonine, Bonine Max, Dramamine Less Drowsy, Dramamine Nausea Long Lasting, ... show all 14 brands

Dosage forms: oral tablet (12.5 mg; 25 mg; 50 mg), oral tablet, chewable (25 mg; 50 mg)

Drug class: Anticholinergic antiemetics

Medically reviewed by Sanjai Sinha, MD. Last updated on Jan 28, 2025.

Uses Side effects Warnings Before taking Dosage Interactions

What is meclizine?

Meclizine is used in adults and children aged 12 years and older to treat or prevent nausea, vomiting and dizziness caused by motion sickness.

Meclizine is also used in adults to treat symptoms of vertigo (dizziness or spinning sensation) caused by disease that affects your inner ear.

Meclizine is most effective if taken before symptoms appear.

Meclizine side effects

Get emergency medical help if you have **signs of an allergic reaction to meclizine**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

Common meclizine side effects may include:

- · drowsiness;
- dry mouth;
- · headache;
- · vomiting; or
- · feeling tired.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Meclizine side effects (more detail)

Related/similar drugs

Zofran

Zofran (ondansetron) is used to prevent nausea and vomiting caused by surgery or cancer medicines ...

Reviews & ratings

8.0 / 10

270 Reviews

View more

Metoclopramide

Metoclopramide is used short-term to treat heartburn caused by gastroesophageal reflux. Learn about

Reviews & ratings

3.9 / 10

517 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Diphenhydramine

Diphenhydramine is an antihistamine used to treat sneezing, runny nose, itching, hives and other ...

Reviews & ratings

5.5 / 10

541 Reviews

View more

Olanzapine

Olanzapine (brand name Zyprexa) is an atypical antipsychotic that may be used to treat adults and ...

Reviews & ratings

5.9 / 10

1,091 Reviews

View more

Promethazine

Promethazine is a phenothiazine and antihistamine used to treat allergies, motion sickness, nausea ...

Reviews & ratings

7.2 / 10

388 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

Reviews & ratings

7.9 / 10

1,054 Reviews

View more

Clonazepam

Clonazepam is used to treat seizures

and panic disorder. Learn about side effects, interactions and ...

Reviews & ratings

8.2 / 10

2,176 Reviews

View more

Ativan

Ativan is a benzodiazepine used to treat anxiety disorders or anxiety associated with depression ...

Reviews & ratings

8.0 / 10

479 Reviews

View more

Warnings

Use meclizine only as directed. Tell your doctor if you use other medicines or have other medical conditions or allergies.

You should not take this medication if you are allergic to meclizine.

Before you take this medicine, tell your doctor if you have liver or kidney disease, asthma, glaucoma, an enlarged prostate, or urination problems.

Meclizine may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

Drinking alcohol can increase certain side effects of meclizine.

Cold or allergy medicine, sedatives, narcotic pain medicine, sleeping pills, muscle relaxers, and medicine for seizures, depression or anxiety can add to sleepiness caused by meclizine.

Before taking this medicine

You should not use meclizine if you are allergic to it.

Do not give meclizine to anyone younger than 12 years old without medical advice.

To make sure this medicine is safe for you, tell your doctor if you have ever had:

- glaucoma;
- breathing problems such as asthma, emphysema, or chronic bronchitis;
- an enlarged prostate or urination problems;

- · recently used alcohol, sedatives, or tranquilizers; or
- liver or kidney disease.

Tell your doctor if you are pregnant or breastfeeding.

1 Meclizine pregnancy and breastfeeding warnings (more detail)

How should I take meclizine?

Take meclizine exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets.

Swallow the tablet whole and do not crush, chew, or break it.

You must chew the *chewable tablet* before you swallow it.

To prevent motion sickness, take meclizine about 1 hour before you travel or anticipate having motion sickness. You may take meclizine once every 24 hours while you are traveling, to further prevent motion sickness.

To treat vertigo, you may need to take meclizine several times daily. Follow your doctor's instructions.

Store at room temperature away from moisture, heat, and light.

Meclizine patient tips (more detail)

Dosing information

Usual Adult Dose for Vertigo:

- -25 to 100 mg orally per day in divided doses
- -Dosing dependent upon clinical response.

Use: Management of vertigo associated with diseases affecting the vestibular system.

Usual Adult Dose for Motion Sickness:

-Initial Dose: 25 to 50 mg orally 1 hour before travel

-Maintenance Dose: Repeat dose every 24 hours if needed

Use: Management of nausea, vomiting, and dizziness associated with motion sickness.

Usual Pediatric Dose for Vertigo:

Age 12 years and older:

- -25 to 100 mg orally per day in divided doses
- -Dosing dependent upon clinical response.

Use: Management of vertigo associated with diseases affecting the vestibular system.

Usual Pediatric Dose for Motion Sickness:

Age 12 years and older:

- -Initial Dose: 25 to 50 mg orally 1 hour before travel
- -Maintenance Dose: Repeat dose every 24 hours if needed

Use: Management of nausea, vomiting, and dizziness associated with motion sickness.

Detailed Meclizine dosage information

What happens if I miss a dose?

Meclizine is used when needed. If you are on a dosing schedule, skip any missed dose. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking meclizine?

Avoid driving or hazardous activity until you know how meclizine will affect you. Your reactions could be impaired.

Avoid drinking alcohol. Drinking alcohol with this medicine can cause side effects.

What other drugs will affect meclizine?

Using meclizine with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which may increase side effects or make the medicines less effective.

Other drugs may interact with meclizine, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all other medicines you use.

• Meclizine drug interactions (more detail)

Does meclizine interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

meclizine
+
Enter a drug name
Add

More about meclizine

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (234)
- Drug images
- · Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: anticholinergic antiemetics
- Breastfeeding

Patient resources

• Meclizine Chewable Tablets patient information

Other brands

Antivert, Meclicot, Dramamine II, Meni-D, Travel Sickness

Professional resources

- Meclizine Hydrochloride monograph
- Meclizine (FDA)

Other brands

Antivert

Related treatment guides

- Vertigo
- Meniere's Disease
- Nausea/Vomiting
- Motion Sickness

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use meclizine only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

Availability

Rx and/or OTC

Pregnancy & Lactation

ন্প Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

7.4 / 10

234 Reviews

Images

Meclizine Hydrochloride 25 mg (44 403)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

Х