

[Home](#)

2. Stalevo 50

# Stalevo 50

**Generic name:** [carbidopa](#), [entacapone](#), and [levodopa](#) [ *KAR-bi-DOE-pa*, *en-TAK-a-pone*, *LEE-voe-DOE-pa* ]

**Brand names:** [Stalevo 100](#), [Stalevo 125](#), [Stalevo 150](#), [Stalevo 200](#), [Stalevo 50](#), [Stalevo 75](#)

**Drug class:** [Dopaminergic antiparkinsonism agents](#)



[Medically reviewed](#) by Drugs.com on Oct 4, 2024. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

## What is Stalevo 50 (Stalevo)?

Stalevo 50 (Stalevo) is a combination medicine used to treat Parkinson symptoms such as muscle stiffness, tremors, spasms, and poor muscle control.

Stalevo may also be used for purposes not listed in this medication guide.

## Stalevo side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

**Stalevo 50 may cause serious side effects. Call your doctor at once if you have:**

- worsening symptoms such as tremors, twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs;
- depressed mood, thoughts about hurting yourself;
- hallucinations, unusual behaviors;
- severe nausea and vomiting;
- severe or ongoing diarrhea, diarrhea that is watery;
- unexplained [muscle pain](#), tenderness, or weakness;
- a light-headed feeling, like you might pass out;
- **fluid build-up in or around the lungs**--pain when you breathe, feeling short of breath while lying down, wheezing, gasping for breath, cough with foamy mucus, cold, clammy skin, anxiety, rapid heartbeats; or
- **severe nervous system reaction**--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats.

Stalevo 50 may cause you to fall asleep during normal daytime activities such as working, talking, eating, or driving. You

may fall asleep suddenly, even after feeling alert. Tell your doctor if you have any problems with daytime sleepiness or [drowsiness](#).

You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs.

You may notice that your sweat, urine, or saliva appears dark in color, such as red, brown, or black. This is not a harmful side effect, but it may cause staining of your clothes or bed sheets.

**Common side effects of Stalevo 50 may include:**

- involuntary muscle movement;
- diarrhea;
- nausea, vomiting, stomach pain;
- discolored urine; or
- dry mouth.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Stalevo 50 side effects](#) (more detail)

## Related/similar drugs

### Sinemet

Sinemet is used to treat symptoms of Parkinson's disease, such as muscle stiffness, tremors ...

#### Reviews & ratings

7.7 / 10

[30 Reviews](#)

[View more](#)

#### FEATURED

### Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

#### Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

## Exelon

Exelon (rivastigmine) is used to treat dementia caused by Alzheimer's or Parkinson's disease ...

### Reviews & ratings

6.5 / 10

[15 Reviews](#)

[View more](#)

## Gocovri

Gocovri (amantadine) is used in the treatment of Parkinson's disease. Includes Gocovri side ...

### Reviews & ratings

[2 Reviews](#)

[View more](#)

## Amantadine

Easy-to-read patient leaflet for amantadine. Includes indications, proper use, special ...

### Reviews & ratings

6.9 / 10

[54 Reviews](#)

[View more](#)

## Warnings

You should not use this medicine if you have narrow-angle glaucoma.

**Do not use this medicine if you have used an MAO inhibitor in the past 14 days**, such as [isocarboxazid](#), [linezolid](#), [methylene blue](#) injection, [phenelzine](#), [rasagiline](#), [selegiline](#), or [tranylcypromine](#).

## Before taking this medicine

You should not take Stalevo 50 if you are allergic to [carbidopa](#), [entacapone](#), or levodopa, or if you have:

- narrow-angle glaucoma.

**Do not use this medicine if you have used an MAO inhibitor in the past 14 days.** A dangerous [drug interaction](#) could occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline,

tranylcypromine, and others.

Tell your doctor if you have ever had:

- glaucoma;
- [depression](#) or other mental illness;
- a [stomach ulcer](#); or
- [liver disease](#).

People with [Parkinson's disease](#) may have a higher risk of [skin cancer](#) (melanoma). Ask your doctor about skin symptoms to watch for.

Tell your doctor if you are pregnant or breastfeeding.

## How should I take Stalevo?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Swallow the tablet whole and do not crush, chew, or break it.

Take only a single tablet at each dosing time. **Do not take more than 1 tablet at a time or take a portion of a broken tablet.** Combining more than one tablet or tablet portions may cause an overdose.

Take your doses at regular intervals to keep a steady amount of the drug in your body at all times. **Ask your doctor about whether you should take Stalevo with or without food.**

**Tell your doctor if the effects of this medicine seem to wear off quickly in between doses.**

Call your doctor if you are sick with vomiting or diarrhea, or if you are sweating more than usual. You can easily become dehydrated from vomiting or diarrhea. Drink plenty of liquids if you are ill.

**Do not take any other medications for Parkinson's disease that your doctor has not prescribed.**

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Stalevo.

**Do not stop using Stalevo suddenly**, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine.

Store at room temperature away from moisture and heat. Keep the bottle tightly closed when not in use.

 [Detailed Stalevo 50 dosage information](#)

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include rapid heartbeats and feeling light-headed.

## What should I avoid while taking Stalevo?

Avoid driving or hazardous activity until you know how Stalevo 50 will affect you. Your reactions could be impaired. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

Avoid taking iron supplements without your doctor's advice. Also avoid increasing the amount of protein in your diet (protein sources include meat, eggs, and cheese). Taking iron or eating too much protein can make it harder for your body to digest and absorb Stalevo.

## What other drugs will affect Stalevo?

Using Stalevo with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

**Many drugs can affect Stalevo.** This includes prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Not all possible interactions are listed here.** Tell your doctor about all your current medicines and any medicine you start or stop using.

 [Stalevo 50 drug interactions](#) (more detail)

## Does Stalevo 50 interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Stalevo 50

+

Enter a drug name

Add

## More about Stalevo 50 (carbidopa / entacapone / levodopa)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)

- [Drug class: dopaminergic antiparkinsonism agents](#)
- [En español](#)

## Patient resources

- [Stalevo 50 advanced reading](#)

## Professional resources

- [Carbidopa, Levodopa and Entacapone Tablets prescribing information](#)

## Other formulations

- [Stalevo](#)

## Related treatment guides

- [Parkinson's Disease](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

### [Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

### DRUG STATUS

#### Availability

**Rx** Prescription only

#### [Pregnancy & Lactation](#)

 Risk data available

#### CSA Schedule\*

**N/A** Not a controlled drug



#### Approval History

 Drug history at FDA



## User Reviews & Ratings

[Review this drug](#)

## Images

# Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



## About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

## Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

## Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT

Copyright © 2000-2025 Drugs.com. All rights reserved.

