

[Home](#)

2. Dasetta 1/35

Dasetta 1/35 (birth control)

Generic name: [ethinyl estradiol and norethindrone \(birth control\)](#) [*ETH-in-il-ess-tra-DYE-ole-and-nor-ETH-in-drone*]

Brand names: [Alyacen 1/35](#), [Alyacen 7/7/7](#), [Aranelle](#), [Aurovela 1.5/30](#), [Aurovela 1/20](#), ... [show all 71 brands](#)

Drug class: [Contraceptives](#)



[Medically reviewed](#) by Drugs.com on Apr 4, 2025. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is Dasetta 1/35?

Dasetta 1/35 is a combination [birth control pill](#) containing female hormones that prevent ovulation (the release of an egg from an ovary). This medication also causes changes in your cervical mucus and uterine lining, making it harder for sperm to reach the uterus and harder for a fertilized egg to attach to the uterus.

Dasetta 1/35 is used as contraception to prevent pregnancy. Dasetta 1/35 is also used to treat moderate acne in women who are at least 15 years old and have started having menstrual periods, and who wish to use [birth control pills](#). There are many available brands of [ethinyl estradiol](#) and norethindrone. Not all brands are listed on this leaflet.

Dasetta 1/35 may also be used for purposes not listed in this medication guide.

Birth control pills side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have:

- **signs of a stroke**--sudden numbness or weakness, severe headache, slurred speech, problems with vision or balance;
- **signs of a blood clot in the lung**--chest pain, sudden cough or [shortness of breath](#), dizziness, coughing up blood;
- **signs of a blood clot deep in the body**--pain, swelling, or warmth in one leg;
- **heart attack symptoms**--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- **liver problems**--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- **increased blood pressure**--severe headache, blurred vision, pounding in your neck or ears;
- **depression**--mood changes, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself;

- swelling in your hands, ankles, or feet, or a breast lump; or
- a change in the pattern or severity of migraine headaches.

Common side effects of Dasetta 1/35 may include:

- nausea, vomiting;
- breast tenderness, or [breakthrough bleeding](#); or
- headache, or problems with contact lenses.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Dasetta 1/35 side effects](#) (more detail)

Related/similar drugs

Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

Reviews & ratings

6.1 / 10

[2,624 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Depo Provera

Depo-Provera is used as contraception to prevent pregnancy. It is also used to reduce pain cause by ...

Reviews & ratings

4.2 / 10

[1 967 Reviews](#)

[1,907 Reviews](#)

[View more](#)

Nexplanon

Nexplanon is a long-term, reversible birth control implant that lasts for three years. Learn about ...

Reviews & ratings

5.0 / 10

[4,864 Reviews](#)

[View more](#)

Yaz

Yaz (drospirenone and ethinyl estradiol) is used as a contraceptive to prevent pregnancy. Includes ...

Reviews & ratings

5.9 / 10

[997 Reviews](#)

[View more](#)

Slynd

Slynd (drospirenone) is a birth control pill used for the prevention of pregnancy. Includes Slynd ...

Reviews & ratings

5.8 / 10

[386 Reviews](#)

[View more](#)

Warnings

Do not use birth control pills if you are pregnant or if you have recently had a baby.

You should not use birth control pills if you have: uncontrolled [high blood pressure](#), heart disease, [coronary artery disease](#), circulation problems (especially with diabetes), undiagnosed vaginal bleeding, liver disease or liver cancer, severe migraine headaches, if you also take certain hepatitis C medication, if you will have major surgery, if you smoke and are over 35, or if you have ever had a heart attack, a stroke, a blood clot, jaundice caused by pregnancy or birth control pills, or cancer of the breast, uterus/cervix, or vagina.

Smoking, especially if you are older than 35, or taking birth control pills can increase your risk of blood clots, stroke, or heart attack.

Before taking this medicine

Taking birth control pills can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, [high cholesterol](#), or if you are overweight. Your risk of stroke or blood clot is highest during your first year of taking birth control pills. Your risk is also high when you restart birth control pills after not taking them for 4 weeks or longer.

Smoking can increase your risk of blood clots, stroke, or heart attack caused by birth control pills, especially if you are older than 35.

Do not use if you are pregnant. Stop using Dasetta 1/35 and tell your doctor right away if you become pregnant, or if you miss 2 menstrual periods in a row. If you have recently had a baby, wait at least 4 weeks before taking birth control pills.

You should not take birth control pills if you have:

- untreated or uncontrolled high blood pressure, or heart disease (chest pain, coronary artery disease, history of heart attack, stroke, or blood clot);
- an increased risk of having blood clots due to a heart problem or a hereditary blood disorder;
- circulation problems (especially if caused by diabetes);
- a history of hormone-related cancer, or cancer of the breast, uterus/cervix, or vagina;
- unusual vaginal bleeding that has not been checked by a doctor;
- liver disease or liver cancer;
- severe migraine headaches (with aura, numbness, weakness, or vision changes), especially if you are older than 35;
- a history of jaundice caused by pregnancy or birth control pills;
- if you smoke and are over 35 years old; or
- if you take any hepatitis C medication containing [ombitasvir/paritaprevir/ritonavir](#) with or without dasabuvir.

Tell your doctor if you have ever had:

- heart problems, high blood pressure, or if you are prone to having blood clots;
- high cholesterol or triglycerides, or if you are overweight;
- depression;
- a seizure or [migraine headache](#);
- diabetes, [gallbladder disease](#), [underactive thyroid](#);
- liver or kidney disease; or
- irregular menstrual cycles, or [fibrocystic breast disease](#), lumps, nodules, or an abnormal mammogram.

Ask a doctor if it is safe to breastfeed while using this medicine.

How should I take birth control pills?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

You may need to use back-up birth control, such as condoms with spermicide, when you first start using this medication. Follow your doctor's instructions.

Take one pill every day, no more than 24 hours apart. When the pills run out, start a new pack the following day. **You may get pregnant if you do not take one pill daily.**

Some birth control packs contain "reminder" pills to keep you on your regular cycle. Your period will usually begin while you are using these reminder pills.


Use a back-up birth control if you are sick with severe vomiting or diarrhea.

You may have breakthrough bleeding. Tell your doctor if it continues or is very heavy.

If you need major surgery or will be on long-term bed rest, you may need to stop using this medicine for a short time. Any doctor or surgeon who treats you should know that you are using Dasetta 1/35.

While taking birth control pills, you will need to visit your doctor regularly.

Store at room temperature away from moisture and heat.

 [Detailed Dasetta 1/35 dosage information](#)

What happens if I miss a dose?

Follow the directions on your prescription label and read all medication guides or instruction sheets. Missing a pill increases your risk of becoming pregnant.

If you miss 1 active pill, take 2 pills on the day you remember. Then take 1 pill per day for the rest of the pack.

If you miss 2 active pills in a row in Week 1 or 2, take 2 pills per day for 2 days in a row. Then take 1 pill per day for the rest of the pack. Use back-up birth control for at least 7 days following the missed pills.

If you miss 2 active pills in a row in Week 3, throw out the rest of the pack and start a new pack the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss 3 active pills in a row in Week 1, 2, or 3, throw out the rest of the pack and start a new pack on the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss 2 or more active pills, you may not have a period during the month. **If you miss a period for 2 months in a row, call your doctor because you might be pregnant.**

If you miss a reminder pill, throw it away and keep taking 1 reminder pill per day until the pack is empty.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose may cause nausea or vaginal bleeding.

What should I avoid while taking birth control pills?

Do not smoke while taking birth control pills, especially if you are older than 35 years of age.

Birth control pills will not protect you from [sexually transmitted diseases](#)--including HIV and AIDS. Using a condom is the only way to protect yourself from these diseases.

What other drugs will affect birth control pills?

Tell your doctor about all your other medicines, especially:

- rifampin;
- troglitazone;
- atorvastatin;
- [glecaprevir/pibrentasvir](#); or
- [anticonvulsants](#) ([antiepileptics](#)) such as phenobarbital, phenytoin, and carbamazepine.

This list is not complete. Other drugs may affect Dasetta 1/35, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

i [Dasetta 1/35 drug interactions](#) (more detail)

Does Dasetta 1/35 interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Dasetta 1/35

+

Enter a drug name

Add

More about Dasetta 1 / 35 (ethinyl estradiol / norethindrone)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(1\)](#)
- [Drug images](#)
- [Side effects](#)

- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: contraceptives](#)
- [En español](#)

Patient resources

Other brands

[Lo Loestrin Fe](#), [Junel Fe 1/20](#), [Blisovi Fe 1/20](#), [Loestrin 21 1/20](#), ... +76 more

Professional resources

- [Dasetta 1/35 prescribing information](#)

Other brands

[Lo Loestrin Fe](#), [Blisovi Fe 1/20](#), [Loestrin 21 1/20](#), [Hailey Fe 1/20](#), ... +80 more

Other formulations

- [Dasetta 7/7/7 birth control](#)

Related treatment guides

- [Birth Control](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.


Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 20.04.

DRUG STATUS

Availability
Rx Prescription only

Pregnancy & Lactation
 Risk data available

CSA Schedule*
N/A Not a controlled drug

Approval History



User Reviews & Ratings

[Review this drug](#)

Related News

[Certain Birth Control Pills, Devices Linked to Higher Heart Risks](#)

Images

[Dasetta 1/35 ethinyl estradiol 0.035 mg / norethindrone 1 mg \(D3\)](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.