

Home

- Parafon Forte DSC
- Consumer

Parafon Forte DSC d

Generic name: Chlorzoxazone [klor-ZOKS-a-zone]

Brand names: Lorzone, Parafon Forte DSC Drug class: Skeletal muscle relaxants

Medically reviewed by Drugs.com. Last updated on Mar 28, 2024.

Uses Before taking Warnings Dosage Side effects Overdose



The Parafon Forte DSC brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be generic equivalents available.

Uses of Parafon Forte DSC:

• It is used to relax muscles.

What do I need to tell my doctor BEFORE I take Parafon Forte DSC?

- If you have an allergy to chlorzoxazone or any other part of Parafon Forte DSC (chlorzoxazone).
- If you are allergic to Parafon Forte DSC (chlorzoxazone); any part of Parafon Forte DSC (chlorzoxazone); or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.

This medicine may interact with other drugs or health problems.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take Parafon Forte DSC (chlorzoxazone) with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take Parafon Forte DSC?

- Tell all of your health care providers that you take Parafon Forte DSC (chlorzoxazone). This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how Parafon Forte DSC (chlorzoxazone) affects you.
- Talk with your doctor before you drink alcohol or use other drugs and natural products that slow your actions.

- This medicine is used with rest, PT (physical therapy), pain drugs, and other therapies.
- Rarely, Parafon Forte DSC (chlorzoxazone) may change the color of the urine. This is normal and not a cause for concern.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

How is this medicine (Parafon Forte DSC) best taken?

Use Parafon Forte DSC (chlorzoxazone) as ordered by your doctor. Read all information given to you. Follow all instructions closely.

• Take with or without food. Take with food if it causes an upset stomach.

What do I do if I miss a dose?

- If you take Parafon Forte DSC (chlorzoxazone) on a regular basis, take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- Many times Parafon Forte DSC (chlorzoxazone) is taken on an as needed basis. Do not take more often than told by the doctor.
- Detailed Parafon Forte DSC dosage information

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever;
 wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Liver problems have rarely happened with Parafon Forte DSC (chlorzoxazone). Sometimes, this has been deadly. Call your doctor right away if you have signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Parafon Forte DSC side effects (more detail)

Related/similar drugs

Flexeril

Flexeril (cyclobenzaprine) is a muscle relaxant used to treat skeletal muscle conditions such as ...

Reviews & ratings

237 Reviews

View more

Robaxin

Robaxin (methocarbamol) is used to treat skeletal muscle conditions such as pain or injury ...

Reviews & ratings

6.3 / 10

80 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Valium

Valium is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.6 / 10

341 Reviews

View more

Zanaflex

Zanaflex (tizanidine) is a short-acting muscle relaxer used to treat spasticity by temporarily ...

Reviews & ratings

7.6 / 10

116 Reviews

View more

Soma

Soma (carisoprodol) is used treat injuries and other painful muscle conditions. Includes Soma side ...

Reviews & ratings

9.1 / 10

200 Reviews

View more

Diazepam

Diazepam is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.4 / 10

884 Reviews

View more

Tizanidine

Tizanidine is a short-acting muscle relaxant and is used to treat spasticity. Learn about side ...

Reviews & ratings

6.2 / 10

380 Reviews

View more

Baclofen

Baclofen is a muscle relaxant used to treat spasticity and muscle pain in conditions like multiple ...

Reviews & ratings

7.0 / 10

447 Reviews

View more

What are some other side effects of Parafon Forte DSC?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- · Feeling sleepy.
- · Dizziness.
- · Feeling tired or weak.
- Feeling nervous and excitable.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

If OVERDOSE is suspected:

If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

How do I store and/or throw out Parafon Forte DSC?

- Store at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so.
 Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Consumer Information Use and Disclaimer

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions
 about Parafon Forte DSC (chlorzoxazone), please talk with your doctor, nurse, pharmacist, or other health care
 provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

More about Parafon Forte DSC (chlorzoxazone)

- · Check interactions
- · Compare alternatives
- Reviews (3)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: skeletal muscle relaxants

Patient resources

• Parafon Forte DSC advanced reading

Other brands

Lorzone, Paraflex, Remular-S

Professional resources

- Parafon Forte DSC prescribing information
- Chlorzoxazone (AHFS Monograph)

Other brands

Lorzone

Related treatment guides

• Muscle Spasm


Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

DRUG STATUS

Availability




Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug



User Reviews & Ratings

3 Reviews

Images

Parafon Forte DSC 500 mg (MCNEIL PARAFON FORTE DSC)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





