

[Home](#)

2. Zenatane

# Zenatane

**Generic name:** [isotretinoin](#) (oral) [ *EYE-so-TRET-i-noyn* ]**Brand names:** [Absorica](#), [Absorica LD](#), [Accutane](#), [Amnesteem](#), [Claravis](#), ... [show all 9 brands](#)**Drug class:** [Miscellaneous uncategorized agents](#)[Medically reviewed](#) by Drugs.com on Jan 21, 2025. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

## What is Zenatane?

Zenatane is a form of vitamin A that is used to treat severe nodular acne that has not responded to other treatments, including antibiotics.

Zenatane is available only from a certified pharmacy under a special program called iPLEDGE.

Zenatane may also be used for purposes not listed in this medication guide.

## Zenatane side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) or **a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

**Zenatane may cause serious side effects. Stop using Zenatane and call your doctor at once if you have:**

- problems with your vision or hearing;
- muscle or joint pain, bone pain, back pain;
- increased thirst, increased urination;
- hallucinations, (see or hearing things that are not real);
- **symptoms of depression**--unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself;
- **signs of liver or pancreas problems**--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, [fast heart rate](#), dark urine, jaundice (yellowing of the skin or eyes);
- **severe stomach problems**--severe stomach or chest pain, pain when swallowing, heartburn, diarrhea, rectal bleeding, bloody or tarry stools; or
- **increased pressure inside the skull**--severe headaches, ringing in your ears, dizziness, nausea, vision problems,

pain behind your eyes.

**Common side effects of Zenatane may include:**

- dryness of your skin, lips, eyes, or nose (you may have nosebleeds);
- vision problems;
- headache, back pain, joint pain, muscle problems;
- skin reactions; or
- [cold symptoms](#) such as stuffy nose, sneezing, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Zenatane side effects](#) (more detail)

## **Related/similar drugs**

## Accutane

Accutane (isotretinoin) is a form of vitamin A and is used to treat severe nodular acne. Includes ...

### Reviews & ratings

7.9 / 10

[657 Reviews](#)

[View more](#)

### FEATURED

## Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

### Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

## Soolantra

Soolantra (ivermectin) is used to treat rosacea. Includes Soolantra side effects, interactions and ...

### Reviews & ratings

7.7 / 10

[299 Reviews](#)

[View more](#)

## Tazorac

Tazorac cream and gel is used to treat acne and plaque psoriasis. Includes Tazorac side effects ...

### Reviews & ratings

7.9 / 10

[38 Reviews](#)

[View more](#)

# Warnings

**Zenatane in just a single dose can cause severe birth defects or death of a baby. Never use Zenatane if you are pregnant or able to become pregnant.**

You must have a negative pregnancy test before taking Zenatane. You will also be required to use two forms of [birth control](#) to prevent pregnancy while taking this medicine. **Stop using this medicine and call your doctor at once if you think you might be pregnant.**

## Before taking this medicine

**Zenatane can cause miscarriage, premature birth, severe birth defects, or death of a baby if the mother takes this medicine at the time of conception or during pregnancy.** Even one dose of [isotretinoin](#) can cause major birth defects of the baby's ears, eyes, face, skull, heart, and brain. **Never use this medicine if you are pregnant or able to become pregnant.**

**For Women:** Unless you have had your uterus and ovaries removed (total [hysterectomy](#)) or have been in menopause for at least 12 months in a row, you are **considered to be able to get pregnant**. You must have 2 negative pregnancy tests before you start taking Zenatane, before each prescription is refilled, right after you take your last dose of isotretinoin, and again 30 days later. All pregnancy testing is required by the iPLEDGE program.

You must agree in writing to use two specific forms of birth control beginning 30 days before you start taking Zenatane and ending 30 days after your last dose. **Both a primary and a secondary form of birth control must be used together.**

**Primary** forms of birth control include:

- tubal ligation (tubes tied);
- vasectomy of the male sexual partner;
- an IUD (intrauterine device);
- estrogen-containing [birth control pills](#) (not mini-pills); and
- hormonal birth control patches, implants, injections, or [vaginal ring](#).

**Secondary** forms of birth control include:

- a male latex condom with or without spermicide;
- a diaphragm plus a spermicide;
- a cervical cap plus a spermicide; and
- a vaginal sponge containing a spermicide.

Stop using Zenatane and call your doctor at once if you have unprotected sex, if you quit using birth control, if your period is late, or if you think you might be pregnant. If you get pregnant while taking this medicine, call the iPLEDGE pregnancy registry at 1-866-495-0654.

**Not having sexual intercourse (abstinence) is the most effective method of preventing pregnancy.**

You should not use Zenatane if you are allergic to it.

Tell your doctor if you have ever had:

- depression or mental illness;
- asthma;
- [liver disease](#);
- diabetes;
- [heart disease](#) or [high cholesterol](#);
- [osteoporosis](#) or low bone mineral density;
- an eating disorder such as anorexia;
- a food or drug allergy; or
- an intestinal disorder such as [inflammatory bowel disease](#) or [ulcerative colitis](#).

Do not breastfeed.

Not approved for use by anyone younger than 12 years old.

## How should I take Zenatane?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Each prescription of isotretinoin must be filled within 7 days of the date it was written by your doctor. You will receive no more than a 30-day supply of isotretinoin at one time.

Always take Zenatane with a full glass of water. Do not chew or suck on the capsule. Swallow it whole.

Follow all directions about taking Zenatane with or without food.

Use Zenatane for the full prescribed length of time. Your acne may seem to get worse at first, but should then begin to improve.

You may need frequent blood tests.

**Never share this medicine with another person, even if they have the same symptoms you have.**

Store at room temperature away from moisture, heat, and light.

 [Detailed Zenatane dosage information](#)

## What happens if I miss a dose?

Skip the missed dose and use your next dose at the regular time. **Do not** use two doses at one time.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose symptoms may include headache, dizziness, vomiting, stomach pain, warmth or tingling in your face, swollen or cracked lips, and [loss of](#)

[balance](#) or coordination.

## What should I avoid while taking Zenatane?

Do not take a vitamin or mineral supplement that contains vitamin A, unless your doctor tells you to.

**Do not donate blood while taking Zenatane and for at least 30 days after you stop taking it.** Donated blood that is later given to a pregnant woman could lead to birth defects in her baby if the blood contains any level of isotretinoin.

While you are taking Zenatane and for at least 6 months after your last dose: Do not use wax hair removers or have dermabrasion or laser skin treatments. Scarring may result.

Zenatane could make you sunburn more easily. Avoid sunlight or tanning beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

Avoid driving or hazardous activity until you know how Zenatane will affect you. Zenatane may impair your vision, especially at night.

## What other drugs will affect Zenatane?

Tell your doctor about all your other medicines, especially:

- phenytoin;
- [St. John's wort](#) (may make birth control pills less effective);
- vitamin or mineral supplements;
- progestin-only birth control pills (mini-pills, may not work as well when taken with Zenatane);
- steroid medicine; or
- a [tetracycline](#) antibiotic, including [doxycycline](#) or minocycline.

This list is not complete. Other drugs may affect Zenatane, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

**i** [Zenatane drug interactions](#) (more detail)

## Does Zenatane interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Zenatane

+

Enter a drug name

Add

## Popular FAQ

Can Accutane cause permanent liver damage?



Is Absorica the same as Accutane?



How long does Absorica take to work?



## More FAQ

- [What are the most common skin conditions? \(with photos\)](#)
- [Isotretinoin - Does accutane help to remove scars from old pimples?](#)

## More about Zenatane (isotretinoin)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(12\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: miscellaneous uncategorized agents](#)
- [Breastfeeding](#)
- [En español](#)

## Patient resources

- [Zenatane advanced reading](#)

## Other brands

[Accutane](#), [Claravis](#), [Amnesteem](#), [Absorica](#), ... +3 more

## Professional resources

- [Zenatane prescribing information](#)
- [Isotretinoin \(AHFS Monograph\)](#)

## Other brands

[Accutane](#), [Claravis](#), [Amnesteem](#), [Absorica](#), [Myorisan](#)

## Related treatment guides

- [Acne](#)

## Further information

- Your doctor pharmacist can provide more information about Zenatane.

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

### [Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 14.02.

### DRUG STATUS

#### Availability

**Rx** Prescription only

#### [Pregnancy & Lactation](#)


 Risk data available

#### CSA Schedule\*

**N/A** Not a controlled drug



#### Approval History

 Drug history at FDA



## User Reviews & Ratings

7.9 / 10

[12 Reviews](#)

## Images


[Zenatane 40 mg \(R137\)](#)



[View larger images](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.







## About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

## Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

## Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.