

Home

Levocetirizine

Levocetirizine sa

Generic name: levocetirizine [*LEE-voe-se-TIR-a-zeen*]

Brand names: Xyzal, Levrix

Dosage forms: oral solution, oral tablet

Drug class: Antihistamines

Medically reviewed by <u>Carmen Pope</u>, <u>BPharm</u>. Last updated on Jun 26, 2024.

Uses Warnings Before taking Side effects Dosage Interactions FAQ

What is levocetirizine?

Levocetirizine is an antihistamine that may be used to reduce allergy symptoms such as red, itchy, or watery eyes; a runny nose; sneezing; rashes; or reactions to insect bites or stings.

Levocetirizine works by reducing the effects of a natural chemical in the body called histamine. Histamine can produce symptoms such as a runny nose or hives.

Levocetirizine is used to treat symptoms of year-round (also called perennial) allergies in adults and children who are at least 6 months old.

Levocetirizine is also used to treat itching and swelling caused by chronic (long-lasting) urticaria (hives) in adults and children who are at least 6 months old.

Levocetirizine was FDA approved in 2007.

Warnings

Follow all the directions on your levocetirizine label and package. Tell all your healthcare providers that you take levocetirizine.

Before taking this medicine

You should not use levocetirizine if you are allergic to levocetirizine or cetirizine (Zyrtec).

You should not take levocetirizine if you have end-stage kidney disease or if you are on dialysis. Any child younger than 12 years old with kidney disease should not take levocetirizine.

Tell your doctor if you have ever had:

kidney disease;

- liver disease;
- urination problems (caused by conditions such as enlarged prostate or spinal cord lesion); or
- gallbladder problems.

Levocetirizine is not expected to harm an unborn baby. Tell your doctor if you are pregnant or intending to become pregnant.

You should not breast-feed while taking levocetirizine.

Levocetirizine is not approved for use by anyone younger than 6 months old.

1 Levocetirizine pregnancy and breastfeeding warnings (more detail)

How should I take levocetirizine?

Take levocetirizine exactly as directed on the label and read all medication guides or instruction sheets about levocetirizine.

A child's dose of levocetirizine is based on the age of the child. Follow your doctor's dosing instructions very carefully.

Do not give a child more than the prescribed dose of levocetirizine. A child's body absorbs twice as much of the same dose of levocetirizine as an adult's body.

Taking more levocetirizine than instructed will not make it more effective, and may cause severe drowsiness.

Levocetirizine is usually taken in the evening, with or without food.

Measure liquid levocetirizine carefully. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

Call your doctor if your symptoms do not improve, if they get worse, or if you also have a fever.

Store levocetirizine at room temperature away from moisture and heat.

What happens if I miss a dose?

If you miss a dose of levocetirizine, take the missed dose as soon as you can. Skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking levocetirizine?

Avoid driving or hazardous activity until you know how levocetirizine will affect you. Your reactions could be impaired.

Drinking alcohol with levocetirizine can cause side effects such as drowsiness.

Levocetirizine side effects

Get emergency medical help if you have **signs of an allergic reaction to levocetirizine:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Levocetirizine may cause serious side effects. Stop using levocetirizine and call your doctor at once if you have:

- · worsening allergy or urticaria symptoms;
- · painful or difficult urination;
- little or no urination;
- · a light-headed feeling, like you might pass out;
- · fever; or
- signs of an ear infection--ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child.

Common side effects of levocetirizine may include:

- drowsiness, tiredness;
- · sinus pain;
- · ear infection;
- · cough;
- fever:
- · nosebleed;
- · vomiting, diarrhea, constipation;
- · dry mouth; or
- · weight gain.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i <u>Levocetirizine side effects</u> (more detail)

Related/similar drugs

Zyrtec

Zyrtec (cetirizine) is used to treat allergy symptoms such as sneezing, itching, watery eyes, or ...

Reviews & ratings

5.8 / 10

197 Reviews

View more

Benadryl

Benadryl is an antihistamine and is used to treat sneezing, runny nose, itching and other ...

Reviews & ratings

7.0 / 10

79 Reviews

View more

FEATURED

Xolair

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

Reviews & ratings

7.5 / 10

229 Reviews

View more

Epinephrine

Epinephrine (adrenaline) Injection is used to treat severe allergic reactions (anaphylaxis) to ...

Reviews & ratings

7.4 / 10

59 Reviews

View more

Diphenhydramine

Diphenhydramine is an antihistamine used to treat sneezing, runny nose, itching, hives and other ...

Reviews & ratings

5.5 / 10

541 Reviews

View more

Prednisone

Prednisone is used to treat allernic

disorders, ulcerative colitis, psoriasis and arthritis. Learn ...

Reviews & ratings

7.5 / 10

867 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Loratadine

Loratadine is a non-drowsy antihistamine used to relieve allergy symptoms like sneezing, runny ...

Reviews & ratings

4.7 / 10

139 Reviews

View more

Levocetirizine dosing information

Usual Adult Dose for Urticaria:

5 mg orally once a day in the evening

Comments: Some patients may be adequately controlled on 2.5 mg orally once a day in the evening.

Usual Adult Dose for Allergic Rhinitis:

Seasonal Allergic Rhinitis (OTC): 5 mg orally once a day in the evening; some patients may be adequately controlled on 2.5 mg once a day

Usual Pediatric Dose for Allergic Rhinitis:

Perennial Allergic Rhinitis:

6 months to 2 years: 1.25 mg orally once a day in the evening

Comments: The recommended dose in pediatric patients 6 months to 2 years should not be exceeded.

Seasonal Allergic Rhinitis (OTC):

6 to 11 years: 2.5 mg orally once a day in the evening

12 years or older: 5 mg orally once a day in the evening; some patients may be adequately controlled on 2.5 mg once a

day

Comments: Avoid use in patients less than 6 years.

Usual Pediatric Dose for Urticaria:

6 months to 5 years: 1.25 mg orally once a day in the evening

6 to 11 years: 2.5 mg orally once a day in the evening

12 years or older: 5 mg orally once a day in the evening; some patients may be adequately controlled on 2.5 mg once a

day

Comments: The recommended doses in pediatric patients 6 months to 12 years should not be exceeded.

Detailed Levocetirizine dosage information

What other drugs will affect levocetirizine?

Using levocetirizine with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- · ritonavir; or
- theophylline.

This list is not complete. Other drugs may affect levocetirizine, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Levocetirizine drug interactions (more detail)

Does levocetirizine interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



More about levocetirizine

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (343)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- Drug class: antihistamines
- Breastfeeding
- En español

Patient resources

• Levocetirizine Tablets patient information

Other brands

Xyzal

Professional resources

- Levocetirizine monograph
- Levocetirizine (FDA)
- Levocetirizine Oral Solution (FDA)

Other brands

Xyzal

Related treatment guides

- Urticaria
- Allergic Rhinitis
- Allergies
- Food Allergies

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

DRUG STATUS

Availability

Rx and/or OTC

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

5.6 / 10

343 Reviews

Images

Levocetirizine Dihydrochloride 5 mg (H 161)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









×

Copyright © 2000-2025 Drugs.com. All rights reserved.