

[Home](#)

2. Requip XL

Requip XL

Generic name: [ropinirole](#) (oral) [roe-PIN-i-role]**Drug class:** [Dopaminergic antiparkinsonism agents](#)[Medically reviewed](#) by Drugs.com on Aug 4, 2023. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

The Requip XL brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be [generic equivalents available](#).

What is Requip XL?

Requip XL has some of the same effects as a chemical called [dopamine](#), which occurs naturally in your body. Low levels of dopamine in the brain are associated with [Parkinson's disease](#).

Requip XL is used to treat symptoms of Parkinson's disease (stiffness, tremors, muscle spasms, and poor muscle control). This medicine is also used to treat [restless legs syndrome](#) (RLS).

Only **immediate-release Requip XL (Requip)** is approved to treat either Parkinson symptoms or RLS. **Extended-release this medicine (Requip XL)** is approved only to treat Parkinson symptoms.

Parkinson's and RLS are two separate disorders. Having one of these conditions will not cause you to have the other condition.

Requip XL may also be used for purposes not listed in this medication guide.

Requip XL side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Some people taking Requip XL have fallen asleep during normal daytime activities such as working, talking, eating, or driving. Tell your doctor if you have any problems with daytime sleepiness or [drowsiness](#).

You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking Requip XL. Talk with your doctor if this occurs.

Requip XL may cause serious side effects. Call your doctor at once if you have:

- extreme drowsiness, falling asleep suddenly (even after feeling alert);
- worsening or no improvement in your symptoms;
- a light-headed feeling, like you might pass out;
- unusual changes in mood or behavior;
- tremors, twitching uncontrollable muscle movements; or
- hallucinations (seeing or hearing things that are not real).

Side effects such as confusion or hallucinations may be more likely in older adults.

Common side effects of Requip XL may include:

- drowsiness, dizziness, weakness;
- [headache](#), confusion, hallucinations;
- increased blood pressure (severe headache, pounding in your neck or ears, nosebleed, irregular heartbeats);
- nausea, [vomiting](#), [upset stomach](#), [constipation](#);
- flu symptoms (fever, chills, body aches);
- sudden muscle movements;
- increased sweating; or
- swelling in your legs or feet.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Requip XL side effects](#) (more detail)

Related/similar drugs

Sinemet

Sinemet is used to treat symptoms of Parkinson's disease, such as muscle stiffness, tremors ...

Reviews & ratings

7.7 / 10

[30 Reviews](#)

[View more](#)

FEATURED

Omvox

Omvox is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

Reviews & ratings

[1 Reviews](#)

[View more](#)

Requip

Requip is used to treat symptoms of Parkinson's disease, such as stiffness, tremors and muscle ...

Reviews & ratings

5.9 / 10

[122 Reviews](#)

[View more](#)

Mirapex

Mirapex (pramipexole) treats the symptoms of Parkinson's disease and is also used to treat Restless ...

Reviews & ratings

7.2 / 10

[121 Reviews](#)

[View more](#)

Exelon

Exelon (rivastigmine) is used to treat dementia caused by Alzheimer's or Parkinson's disease ...

Reviews & ratings

6.5 / 10

[15 Reviews](#)

[View more](#)

Gocovri

Gocovri (amantadine) is used in the treatment of Parkinson's disease. Includes Gocovri side ...

Reviews & ratings

[2 Reviews](#)

[View more](#)

Inbrija

Inbrija is an inhaled medicine used to treat the symptoms of Parkinson's disease. Includes side ...

Reviews & ratings

[Add a review](#)

[View more](#)

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use Requip XL if you are allergic to it.

To make sure Requip XL is safe for you, tell your doctor if you have:

- high or [low blood pressure](#);
- kidney disease (or if you are on dialysis);
- [heart disease](#), heart rhythm problems;
- a sleep disorder such as [narcolepsy](#), or other conditions that may cause daytime sleepiness; or
- if you smoke.

People with Parkinson's disease may have a higher risk of [skin cancer \(melanoma\)](#). Talk to your doctor about this risk and what skin symptoms to watch for.

It is not known whether Requip XL will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether [ropinirole](#) passes into breast milk or if it could affect the nursing baby. Ropinirole may slow breast milk production. Tell your doctor if you are breast-feeding.

Requip XL is not approved for use by anyone younger than 18 years old.

How should I take Requip XL?

Follow all directions on your prescription label. Your doctor may occasionally change your dose. Do not take this medicine in larger or smaller amounts or for longer than recommended.

If you are taking immediate-release Requip XL (**Requip**) you should not take extended-release this medicine (**Requip XL**) at the same time.

The dose and timing of ropinirole in treating Parkinson's disease is different from the dose and timing in treating RLS. Follow the directions on your prescription label. Ask your pharmacist if you have any questions about the kind of ropinirole you receive at the pharmacy.

Requip XL can be taken with or without food. Take the medicine at the same time each day.

Do not crush, chew, or break an **extended-release tablet (Requip XL)**. Swallow it whole.

Call your doctor if you see part of the Requip XL tablet in your stool. This is a sign that your body may not have absorbed all of the medicine.

If you are taking this medicine for RLS, tell your doctor if your symptoms get worse, if they occur in the morning or earlier than usual in the evening, or if you feel restless symptoms in your hands or arms.

It may take up to several weeks before your symptoms improve. Keep using the medication as directed and tell your doctor if your symptoms do not improve.

Do not stop using Requip XL suddenly, or you could have unpleasant withdrawal symptoms. Follow your doctor's instructions about tapering your dose.

Store at room temperature away from moisture, heat, and light. Keep the bottle tightly closed when not in use.

 [Detailed Requip XL dosage information](#)

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose.

Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Requip XL?

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Get up slowly and steady yourself to prevent a fall. Dizziness may impair your thinking or reactions. Avoid driving or operating machinery until you know how Requip XL will affect you.

Drinking alcohol can increase certain side effects of ropinirole.

What other drugs will affect Requip XL?

Taking Requip XL with other drugs that make you sleepy can worsen this effect. Ask your doctor before taking a sleeping pill, narcotic medication, muscle relaxer, or medicine for anxiety, [depression](#), or [seizures](#).

Other drugs may interact with ropinirole, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

 [Requip XL drug interactions](#) (more detail)

Does Requip XL interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Requip XL

+

Enter a drug name

Add

More about Requip XL (ropinirole)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(7\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: dopaminergic antiparkinsonism agents](#)
- [Breastfeeding](#)

Patient resources

- [Requip XL advanced reading](#)

Other brands

[Requip](#)

Professional resources

- [Requip-XL prescribing information](#)
- [ROPINIRole \(AHFS Monograph\)](#)

Other brands

[Requip](#)

Related treatment guides

- [Parkinson's Disease](#)
- [Periodic Limb Movement Disorder](#)

Further information


Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 16.01.

DRUG STATUS

Availability	
	Discontinued
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	Drug history at FDA

User Reviews & Ratings

[7 Reviews](#)

Images

[Requip XL 8 mg \(GS 5CC\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)


[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.