

[Home](#)[2. Dosage](#)[3. Cyclizine](#)

Cyclizine Dosage

Applies to the following strengths: 25 mg; 50 mg

Usual Pediatric Dose for:

- [Motion Sickness](#)

Additional dosage information:

- [Renal Dose Adjustments](#)
- [Liver Dose Adjustments](#)
- [Precautions](#)
- [Dialysis](#)
- [Other Comments](#)

Usual Pediatric Dose for Motion Sickness

6 years and older: 25 mg orally every 6 to 8 hours

- Maximum dose: 75 mg/day

Comments:

- For the prevention of travel sickness, patients should take the first dose approximately 1 hour prior to departure or at the onset of symptoms.
- Patients should be cautioned to avoid driving or operating machinery while taking this drug.

Use: To prevent and treat nausea, vomiting, or dizziness associated with motion sickness

Renal Dose Adjustments

Data not available

Liver Dose Adjustments

Data not available

Precautions

CONTRAINDICATIONS:

- Use in children under 6 years of age
- Use in children with a breathing problem (e.g., chronic bronchitis) or glaucoma without consulting with a healthcare provider
- Use in children taking sedatives or tranquilizers without prior consultation with a healthcare provider

Safety and efficacy have not been established in patients younger than 6 years; this drug is not recommended for use in these patients.

Consult WARNINGS section for additional precautions.

Dialysis

Data not available

Other Comments

Administration advice:

- IV formulations should be injected slowly.
- Oral formulations should be given at least 1 hour prior to departure to prevent motion sickness.

Storage requirements:

- IV: Protect from light.

Patient advice:

- Inform patients that this drug may cause drowsiness, and they should avoid driving or operating machinery if these side effects occur.
- This drug may increase the side effects of alcohol; patients should avoid the concomitant use of alcohol.
- Advise patients to speak to their healthcare provider if they become pregnant, intend to become pregnant, or are breastfeeding.

More about cyclizine

- [Check interactions](#)
- [Compare alternatives](#)

- [Reviews \(55\)](#)
- [Side effects](#)
- [During pregnancy](#)
- [Drug class: anticholinergic antiemetics](#)
- [Breastfeeding](#)

Related treatment guides


- [Nausea/Vomiting](#)
- [Motion Sickness](#)


Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.


[Medical Disclaimer](#)

DRUG STATUS

Availability
 Discontinued

Pregnancy & Lactation
 Risk data available

CSA Schedule*
N/A Not a controlled drug

Approval History
 Drug history at FDA



User Reviews & Ratings

6.1 / 10

[55 Reviews](#)

Related Drugs

[Benadryl](#), [Phenergan](#), [scopolamine](#), [hydroxyzine](#), [Ativan](#), [diphenhydramine](#), [Dramamine](#), [promethazine](#), [Zofran](#), [olanzapine](#)

Images

[Marezine \(cyclizine\) 50 mg \(MAREZINE T4A\)](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.