

Home

2. Sucralfate

Sucralfate 4

Generic name: sucralfate (oral) [soo-KRAL-fate]

Brand name: Carafate

Dosage forms: oral suspension (1 g/10 mL), oral tablet (1 g)

Drug class: Miscellaneous GI agents

9

Medically reviewed by Drugs.com on Feb 24, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is sucralfate?

Sucralfate is used short-term (up to 8 weeks) to treat an active duodenal ulcer.

Sucralfate works mainly in the lining of the stomach and is not highly absorbed into the body. sucralfate adheres to ulcer sites and protects them from acids, enzymes, and bile salts.

Sucralfate can heal an active ulcer, but it will not prevent future ulcers from occurring.

Sucralfate may also be used for purposes not listed in this medication guide.

Sucralfate side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Common side effects of sucralfate may include:

- · constipation, diarrhea;
- nausea, vomiting, gas, indigestion;
- itching, rash;
- dizziness, drowsiness;
- sleep problems (insomnia);
- · headache; or
- back pain.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Related/similar drugs

Protonix

Protonix (pantoprazole) is used to treat erosive esophagitis and other conditions involving excess ...

Reviews & ratings

7.2 / 10

83 Reviews

View more

Nexium

Nexium is used to treat GERD and other conditions involving excessive stomach acid. Learn about ...

Reviews & ratings

6.2 / 10

211 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Prilosec

Prilosec (omeprazole) is used to treat symptoms of GERD and other conditions caused by excess ...

Reviews & ratings

6.9 / 10

53 Reviews

View more

Carafate

Carafate is an anti-ulcer medication used to treat active duodenal ulcers. Learn about side ...

Reviews & ratings

6.2 / 10

72 Reviews

View more

Lansoprazole

Lansoprazole is used to treat stomach and intestinal ulcers, erosive esophagitis and ...

Reviews & ratings

6.6 / 10

143 Reviews

View more

Warnings

The liquid form of sucralfate should never be injected through a needle into the body, or death may occur.

Before taking this medicine

You should not use sucralfate if you are allergic to it.

Tell your doctor if you have ever had:

- · diabetes;
- kidney disease (or if you are on dialysis); or
- trouble swallowing tablets.

Older adults may be more sensitive to the effects of sucralfate.

Tell your doctor if you are pregnant or breastfeeding.

Do not give this medicine to a child without medical advice.

How should I take sucralfate?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine

exactly as directed.

Take sucralfate on an empty stomach.

Shake the **oral suspension** (liquid) before you measure a dose. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

If you are diabetic, check your blood sugar regularly. Your doctor may adjust your dose based on your blood sugar levels.

The liquid from of this medicine should never be injected through a needle into the body, or death may occur. Sucralfate oral suspension is to be taken only by mouth.

It may take 2 to 8 weeks before you receive the full benefit of taking sucralfate. Use this medicine for the full prescribed length of time, even if your symptoms quickly improve.

Your doctor may want you to keep taking sucralfate at a lower dose once your active ulcer has healed. **Follow your doctor's dosing instructions very carefully.**

Store at room temperature away from moisture and heat. Do not allow the liquid medicine to freeze.

1 Sucralfate patient tips (more detail)

Sucralfate dosing information

Usual Adult Dose for Duodenal Ulcer:

1 g orally 4 times a day

-Duration of therapy: 4 to 8 weeks

Comments:

- -Each dose should be taken on an empty stomach.
- -For most patients, treatment should continue for 4 to 8 weeks; however, treatment may be stopped early if healing is demonstrated on x-ray/endoscopic tests.
- -Antacids may be given, but use should be separated by at least 30 minutes.

Use: Short-term treatment of active duodenal ulcers

Usual Adult Dose for Duodenal Ulcer Prophylaxis:

Oral Tablets: 1 g orally 2 times a day -Duration of therapy: Up to 1 year

Comments:

- -Each dose should be taken on an empty stomach.
- -Controlled studies did not extend beyond 12 months.

Use: Maintenance therapy for patients after healing of acute duodenal ulcers

Detailed Sucralfate dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking sucralfate?

Avoid taking any other medications within 2 hours before or after you take sucralfate. Sucralfate can make it harder for your body to absorb other medications you take by mouth.

Ask your doctor before using an antacid, and use only the type your doctor recommends. Some antacids can make it harder for sucralfate to work in your stomach. **Avoid taking an antacid within 30 minutes before or after taking sucralfate.**

What other drugs will affect sucralfate?

Other drugs may affect sucralfate, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

Sucralfate drug interactions (more detail)

Does sucralfate interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

sucralfate	
+	
Enter a drug name	Add

More about sucralfate

- Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (241)
- Drug images
- Latest FDA alerts (1)
- · Side effects

- Dosage information
- · Patient tips
- During pregnancy
- Support group
- Drug class: miscellaneous GI agents
- Breastfeeding
- En español

Patient resources

• Sucralfate Oral Suspension patient information

Other brands

Carafate

Professional resources

- Sucralfate monograph
- Sucralfate (FDA)
- Sucralfate Suspension (FDA)
- Sucralfate Tablet (FDA)

Other brands

Carafate

Related treatment guides

- GERD
- Duodenal Ulcer
- Hyperphosphatemia of Renal Failure
- Duodenal Ulcer Prophylaxis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

6.2 / 10

241 Reviews

Images

Sucralfate 1 g (TEVA 22 10)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×