

Home

2. Ivacaftor

Ivacaftor

Generic name: ivacaftor [EYE-va-KAF-tor]

Brand name: Kalydeco

Dosage forms: oral granule for reconstitution (13.4 mg; 25 mg; 5.8 mg; 50 mg; 75 mg), oral tablet (150 mg)

Drug class: CFTR potentiators



Medically reviewed by Drugs.com on Aug 21, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is ivacaftor?

Ivacaftor is used to treat cystic fibrosis in adults and children who are at least 1 month old and weigh at least 6 pounds (3 kilograms).

Ivacaftor **tablet** can be used in children aged 6 years and older. Ivacaftor **granules** can be used in children aged 1 month to under 6 years old.

Your doctor will make sure you have the specific gene mutation related to cystic fibrosis to be treated with ivacaftor.

Ivacaftor may also be used for purposes not listed in this medication guide.

Ivacaftor side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

lvacaftor may cause serious side effects. Call your doctor at once if you have:

- blurred vision, eye pain or redness, seeing halos around lights;
- low blood sugar--headache, hunger, weakness, sweating, confusion, irritability, dizziness, fast heart rate, or feeling
 jittery; or
- **liver problems**--loss of appetite, nausea, vomiting, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Common side effects of ivacaftor may include:

- muscle or joint pain;
- dizziness;
- · rash, acne;

- · headache;
- · mouth pain;
- · stomach pain, nausea, diarrhea; or
- cold symptoms such as stuffy nose, sneezing, runny nose, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



Ivacaftor side effects (more detail)

Related/similar drugs

Zithromax

Zithromax (azithromycin) treats infections caused by bacteria, such as respiratory infections, skin ...

Reviews & ratings

7.1 / 10

127 Reviews

View more

Creon

Creon (pancrelipase) contains digestive enzymes and is used to improve food digestion in people who

Reviews & ratings

6.3 / 10

59 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Zenpep

Zenpep (pancrealipase) is used to replace pancreatic enzymes when the body does not have enough of ...

Reviews & ratings

7.7 / 10

16 Reviews

View more

Vitamin e

Vitamin E is an antioxidant that occurs naturally in some foods and is used to treat or prevent ...

Reviews & ratings

7 Reviews

View more

Warnings

Use only as directed. Tell your doctor if you use other medicines or have other medical conditions or allergies.

Before taking this medicine

You should not be treated with ivacaftor if you are allergic to it, or if you have or have ever had:

• liver or kidney disease.

It is not known if ivacaftor will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

Ask a doctor if it is safe to breastfeed while using ivacaftor.

How should I take ivacaftor?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Take ivacaftor with food that contains fat, such as eggs, milk products such as whole milk, cheese, yogurt, breast milk or infant formula. Follow your doctor's instructions carefully.

Swallow the tablet whole and do not crush, chew, or break it.

Mix the ivacaftor **granules** with 1 teaspoon of soft food such as pureed fruits or vegetables, yogurt, applesauce, water, breast milk, infant formula, milk, or juice. Mix only 1 dose at a time and use within 1 hour after mixing. Feed the child a high-fat food after or before giving the granule mixture.

Take this medicine at evenly spaced intervals every 12 hours.

Doses are based on weight in children. Your child's dose may change if the child gains or loses weight.

You will need frequent blood tests to check your liver function.

Your child may need frequent vision exams.

Store at room temperature away from moisture and heat. Keep the medicine in the original packaging.

Detailed Ivacaftor dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, **with a fatty meal**, but skip the missed dose if you are more than 6 hours late for the dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking ivacaftor?

Avoid driving or hazardous activity until you know how ivacaftor will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Grapefruit may interact with ivacaftor and cause side effects. Avoid consuming grapefruit products.

What other drugs will affect ivacaftor?

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which may increase side effects or make the medicines less effective.

Tell your doctor about all your other medicines, especially:

- St. John's wort;
- seizure medicine such as carbamazepine, phenobarbital, or phenytoin;
- an antibiotic--clarithromycin, erythromycin, telithromycin, rifabutin, rifampin; or
- antifungal medicine--fluconazole, itraconazole, ketoconazole, posaconazole, voriconazole.

This list is not complete. Other drugs may affect ivacaftor, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Ivacaftor drug interactions (more detail)

Does ivacaftor interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

ivacaftor	
+	
Enter a drug name	Add

Popular FAQ

What is an orphan drug?

How does Kalydeco work?

More FAQ

- What is the difference between Symdeko and Orkambi?
- What is the difference between Symdeko and Kalydeco?

More about ivacaftor

- Check interactions
- Compare alternatives
- Reviews (1)
- Side effects
- Dosage information
- During pregnancy
- Drug class: CFTR potentiators
- Breastfeeding

Patient resources

• Ivacaftor advanced reading

Other brands

Kalydeco

Professional resources

· Ivacaftor monograph

Other brands

Kalydeco

Related treatment guides

Cystic Fibrosis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 6.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্প Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

Review this drug

Images

Kalydeco (ivacaftor) 150 mg (V 150)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









×

Copyright © 2000-2025 Drugs.com. All rights reserved.