

#### Home

2. Donepezil

# **Donepezil** ⊲<sub>0</sub>

**Pronunciation:** *don-eh-pee-zill* **Generic name:** donepezil

Brand names: Aricept, Aricept ODT

Dosage forms: Tablets (5 mg, 10 mg, 23 mg), ODT Tablets (5 mg, 10 mg)

Drug class: Cholinesterase inhibitors

Medically reviewed by Carmen Pope, BPharm. Last updated on Jul 4, 2024.

Uses Side effects Before taking Dosage What to avoid Interactions FAQ

## What is Donepezil?

Donepezil (brand name Aricept) may be used to treat dementia caused by Alzheimer's disease. **Donepezil can help** improve attention, memory, behavior, and ability to do daily activities in dementia patients.

Donepezil is not a cure for Alzheimer's disease; it may help improve symptoms, but the condition will progress over time, even in people who take this medicine. Research has determined that donepezil is effective in mild, moderate, and severe Alzheimer's disease.

Donepezil improves the function of nerve cells in the brain and works by preventing the breakdown of a chemical called acetylcholine (ah SEET il KOE leen). People with dementia usually have lower levels of this chemical, which is important for memory, thinking, and reasoning. Donepezil is from the drug class cholinesterase inhibitors.

Donepezil was FDA-approved on November 25, 1996, under the brand name Aricept. Generic donepezil became available in 2009.

## Donepezil side effects

The most common side effects of donepezil are:

- nausea
- diarrhea
- · not sleeping well
- vomiting
- · muscle cramps
- · feeling tired
- · not wanting to eat.

These side effects may get better after you take donepezil for a while. This is not a complete list of side effects. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA1088.

### Serious side effects and warnings

Get emergency medical help if you have signs of an allergic reaction which may include symptoms such as hives, difficulty breathing, and swelling of your face, lips, tongue, or throat.

Donepezil can cause serious side effects including:

**Slow heartbeat and fainting**. This happens more often in people with heart problems. Call your doctor right away if you feel faint or lightheaded while taking donepezil.

**More stomach acid**. This raises the chance of ulcers and bleeding, especially when taking higher dosages of donepezil. The risk is higher for people who have had stomach ulcers or also take aspirin or other NSAIDs.

Worsening of lung problems in people with asthma or other lung disease.

Donepezil may also exaggerate the response to certain muscle relaxants during surgery.

There is a risk of generalized convulsive seizures, gastrointestinal bleeding, and difficulty passing urine in people taking donepezil.

Donepezil can cause vomiting which may be severe. Your healthcare provider will monitor you for this.

Stop using this medicine and call your doctor at once if you have:

- severe or ongoing vomiting
- a light-headed feeling, like you might pass out
- slow heartbeats
- seizure (convulsions)
- · painful or difficult urination
- · new or worsening breathing problems or
- signs of stomach bleeding severe heartburn or stomach pain, bloody or tarry stools, coughing up blood or vomit
  that looks like coffee grounds.

This is not a complete list of side effects of donepezil, and others may occur. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Donepezil side effects (more detail)

## Related/similar drugs

#### **Namenda**

Namenda is used to treat dementia associated with Alzheimer's disease. Learn about side effects ...

### **Reviews & ratings**

5.8 / 10

33 Reviews

#### View more

### Leqembi

Leqembi (lecanemab) is used to treat Alzheimer's disease to help slow the progression of dementia s ...

### Reviews & ratings

1 Reviews

#### View more

#### FEATURED

### **Aricept**

Aricept is used to treat mild to severe dementia caused by Alzheimer's disease. Learn about side ...

### **Reviews & ratings**

5.7 / 10

47 Reviews

#### View more

### **Exelon**

Exelon (rivastigmine) is used to treat dementia caused by Alzheimer's or Parkinson's disease ...

### Reviews & ratings

6.5 / 10

15 Reviews

#### View more

### **Kisunla**

Kisunla (donanemab) is used to treat Alzheimer's disease to help slow down the decline in memory,  $t \dots$ 

### **Reviews & ratings**

Add a review

#### Vitamin e

Vitamin E is an antioxidant that occurs naturally in some foods and is used to treat or prevent ...

### **Reviews & ratings**

7 Reviews

View more

### **Memantine**

Memantine (brand name Namenda) is an oral NMDA blocker that may be used to treat moderate to severe ...

#### **Reviews & ratings**

6.3 / 10

108 Reviews

View more

## Before taking this medicine

You should not use this medicine if you are allergic to donepezil, Aricept, or any inactive ingredients in the medicine. Tell your doctor if you have any allergies to medicines.

To make sure this medicine is safe for you, tell your doctor if you have:

- a heart rhythm disorder such as "sick sinus syndrome" (slow heartbeats)
- an enlarged prostate
- · urination problems
- a seizure disorder such as epilepsy
- · a history of stomach ulcers
- · liver or kidney disease
- · trouble swallowing
- asthma, chronic obstructive pulmonary disease (COPD), or another breathing disorder
- are pregnant or intending to become pregnant
- · are breastfeeding.

If you need to have any type of surgery, tell the surgeon ahead of time that you are taking Donepezil.

### **Pregnancy**

It is not known whether donepezil will harm an unborn baby.

### **Breastfeeding**

It is not known whether donepezil passes into breast milk or if it could harm a nursing baby.

1 Donepezil pregnancy and breastfeeding warnings (more detail)

### How should I take donepezil?

Take donepezil exactly as it was prescribed for you. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

- Take donepezil one time each day.
- You may take donepezil with or without food.

Regular tablets: Do not crush, chew, or break the regular tablet. Swallow it whole.

How to take donepezil ODT (orally disintegrating tablet):

- Keep the tablet in its blister pack until you are ready to take it. Open the package and peel or cut the backing from the tablet blister. Do not push a tablet through the backing or you may damage the tablet.
- Use dry hands to remove the tablet and place it in your mouth.
- Do not swallow the tablet whole. Allow it to dissolve on your tongue.
- Swallow several times as the tablet dissolves. After the tablet dissolves completely, drink a glass of water.

If you need surgery or dental work, tell the surgeon ahead of time that you are using donepezil. You may need to stop using the medicine for a short time.

- Do should not stop using donepezil without your doctor's advice.
- Donepezil patient tips (more detail)

## **Donepezil dosing information**

### Mild to moderate Alzheimer's disease

Initial dose: donepezil 5 mg orally once a day, in the evening before going to bed.

Maintenance dose: donepezil 10 mg orally once a day after the patient has been on an initial dose of 5 mg once a day for 4 to 6 weeks.

### Moderate to severe Alzheimer's disease

Initial dose: donepezil 5 mg orally once a day, in the evening before going to bed.

Maintenance dose: donepezil 10 mg orally once a day after the patient has been on an initial dose of 5 mg once a day for 4 to 6 weeks.

• This dosage may be increased to a maximum of donepezil 23 mg once a day if needed and the person has been taking donepezil 10 mg once a day for at least 3 months.

**Comments**: A higher dose of 10 mg may or may not provide a statistically significantly greater clinical benefit than a 5 mg dose. Prescriber and patient's preferences should be considered.

Detailed Donepezil dosage information

### Donepezil dosage forms

Donepezil is available as tablets (5 mg, 10 mg, 23 mg) and orally disintegrating tablets (ODT) (5 mg, 10 mg).

## What happens if I miss a dose?

Take the missed donepezil dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up for the missed dose.

If you miss your doses for more than 7 days in a row, call your doctor before taking the medicine again.

### What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe nausea, vomiting, drooling, sweating, blurred vision, feeling light-headed, slow heartbeat, shallow breathing, muscle weakness, fainting, or seizure (convulsions).

### What to avoid

Donepezil can cause side effects that may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be awake and alert.

## What other drugs will affect Donepezil?

Tell your doctor about all medicines you use, and those you start or stop using during your treatment with Donepezil, especially:

- NSAIDs (nonsteroidal anti-inflammatory drugs) aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others
- Anticholinergics such as benztropine or scopolamine
- Succinylcholine, similar neuromuscular blocking agents, or cholinergic agonists (such as varenicline).

This list is not complete. Other drugs may interact with donepezil, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed in this medication guide.

<u>Donepezil drug interactions</u> (more detail)

### Does donepezil interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



## **Storage**

Store at room temperature between 59° to 86°F (15° to 30°C).

## Ingredients

Active ingredient: donepezil hydrochloride

**Inactive ingredients:** These vary depending on the manufacturer. Refer to the donepezil package insert with your medicine for more information.

## Who makes donepezil?

Various manufacturers make donepezil tablets. Examples include:

Aricept -- Eisai Inc.

Generics -- Camber Pharmaceuticals, Quallent Pharmaceuticals Health LLC, Alembic Pharmaceuticals Limited, and others.

## **Popular FAQ**

When is the best time to take Aricept (donepezil)?

How does Aricept work? 

✓

#### References

- 1. Donepezil hydrochloride tablets
- 2. Aricept Package Insert

## More about donepezil

- · Check interactions
- Compare alternatives

- Pricing & coupons
- Reviews (140)
- Drug images
- · Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: cholinesterase inhibitors

### **Patient resources**

- Donepezil (Oral) advanced reading
- Donepezil (Transdermal) (Advanced Reading)
- Donepezil 23 mg Tablets
- Donepezil Orally Disintegrating Tablets

### Other brands

Aricept, Adlarity

### **Professional resources**

- Donepezil monograph
- Donepezil (FDA)
- Donepezil ER Tablets (FDA)
- Donepezil Orally Disintegrating Tablet (FDA)

### Other brands

Aricept, Adlarity

### Related treatment guides

- Mild Cognitive Impairment
- Alzheimer's Disease
- Lewy Body Dementia

### **Further information**

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

#### **DRUG STATUS**

### **Availability**

Rx Prescription only

### **Pregnancy & Lactation**

ন্প Risk data available

#### **CSA Schedule\***

N/A Not a controlled drug

### **Approval History**

The Drug history at FDA


### **User Reviews & Ratings**

5.8 / 10

140 Reviews

### **Images**

Donepezil Hydrochloride 5 mg (5)



# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





### **About**

About Drugs.com

Advertising policy

Attribution & citations

#### Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

### **Support**

Help center

Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

×