



[Home](#)

2. [Natazia](#)

Natazia

Generic name: [dienogest and estradiol](#) [*dye-EN-oh-jest-and-ESS-tra-DYE-ole*]

Drug class: [Contraceptives](#)

[Medically reviewed](#) by Drugs.com. Last updated on Apr 23, 2024.

[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#)

What is Natazia?

Natazia contains a combination of dienogest and estradiol. Dienogest and estradiol are female hormones that prevent ovulation (the release of an egg from an ovary). This medicine also causes changes in your cervical mucus and uterine lining, making it harder for sperm to reach the uterus and harder for a fertilized egg to attach to the uterus.

Natazia is used to [prevent pregnancy](#).

Natazia is also used to treat [heavy menstrual bleeding](#) that is not caused by any medical condition of the uterus.

Warnings

Do not use Natazia if you are pregnant or if you have recently had a baby.

You should not take Natazia if you have any of the following conditions: uncontrolled high blood pressure, heart disease, a blood-clotting disorder, circulation problems, diabetic problems with your eyes or kidneys, unusual vaginal bleeding, liver disease or liver cancer, severe migraine headaches, if you smoke and are over 35, or if you have ever had breast or uterine cancer, a heart attack, a stroke, or a blood clot.

Taking Natazia can increase your risk of blood clots, stroke, or heart attack, especially if you have certain other conditions, or if you are overweight.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. You should not take Natazia if you smoke and are over 35 years old.

Missing a pill increases your risk of becoming pregnant. Carefully follow the "missed dose" instructions if you forget to take your medicine.

Some drugs can make Natazia less effective in preventing pregnancy, including antibiotics, hepatitis C medications, HIV/AIDS medications, seizure medications, or barbiturate sedatives. Tell your doctor about all other medications you use.

Before taking this medicine

Taking Natazia can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, high cholesterol, or if you are overweight. Your risk of stroke or blood clot is highest during your first year of taking Natazia. Your risk is also high when you restart this medicine after not taking it for 4 weeks or longer.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. Your risk increases the older you are and the more you smoke. You should not take Natazia if you smoke and are over 35 years old.

Do not use if you are pregnant. Stop taking Natazia and tell your doctor if you become pregnant, or if you miss two menstrual periods in a row. If you have recently had a baby, wait at least 4 weeks before taking this medicine.

You should not take Natazia if you have:

- untreated or uncontrolled high blood pressure;
- heart disease (coronary artery disease, uncontrolled heart valve disorder, history of heart attack, stroke, or blood clot);
- a blood-clotting disorder or circulation problems;
- problems with your eyes, kidneys or circulation caused by diabetes;
- a history of hormone-related cancer such as breast or uterine cancer;
- unusual vaginal bleeding that has not been checked by a doctor;
- liver disease or liver cancer;
- severe migraine headaches; or
- if you smoke and are over 35 years old.

To make sure Natazia is safe for you, tell your doctor if you have:

- high blood pressure, high cholesterol or triglycerides, or if you are overweight;
- a history of depression;
- diabetes, underactive thyroid, gallbladder disease;
- seizures or epilepsy; or
- a history of jaundice caused by pregnancy or birth control pills.

The hormones in Natazia can pass into breast milk and may harm a nursing baby. This medicine may also slow breast milk production. Do not use if you are breast feeding a baby.

 [Natazia pregnancy and breastfeeding warnings](#) (more detail)

How should I take Natazia?

Natazia contains a combination of dienogest and estradiol. contains 5 different colors of pills. Take 1 pill each day in the exact order directed on the blister pack. Follow the arrows shown on each row of pills in the pack. Use a back-up birth control, such as condoms or a spermicide, for the first 9 days. Do not take two different kinds of birth control pills at the same time.

If you are switching from another birth control pill, start taking Natazia on the first day of your period. If you were taking

progestin-only pills, start taking Natazia on the day you would have taken your next pill.

If you are switching from a birth control implant, intrauterine device (IUD), vaginal ring, or skin patch, start taking Natazia on the day the other birth control device is removed. If you are switching from a birth control injection, start taking Natazia on the day you would have received your next scheduled injection.

Tell your doctor if you have heavy breakthrough bleeding. You may also have very little or no bleeding during your periods.

If you need surgery or medical tests or if you will be on bed rest, you may need to stop using this medicine for a short time. Any doctor or surgeon who treats you should know that you are using Natazia.

While taking Natazia, you will need to visit your doctor regularly.

Store at room temperature away from moisture and heat.

 [Detailed Natazia dosage information](#)

What happens if I miss a dose?

Follow the patient instructions provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions. Missing a pill by more than 12 hours increases your risk of becoming pregnant. Vomiting or having diarrhea within 4 hours after taking a pill is the same as missing that dose.

If you miss one pill:

- During Days 1 through 17, take the missed pill as soon as you remember, then take your next pill at the usual time. Use back-up birth control for at least 9 days.
- During Days 18 through 24, throw out the rest of the pack and start a new one the same day. Take the Day 1 pill from the new pack and then take one pill per day in the order directed on the pack. Use back-up birth control for at least 9 days.
- During Days 25 through 28, take the missed pill as soon as you remember, then take your next pill at the usual time. You do not need back-up birth control if you miss one pill during Days 25 through 28.

If you miss two pills:

- During Days 1 through 16, skip the missed pills and start with the pill that corresponds to the day you remember you missed your doses. Then take one pill per day in the order directed on the pack. Use your back up birth control for at least 9 days
- During Days 17 through 24, throw out the rest of the pack and start a new one the same day. Take the Day 3 pill from the new pack and then take one pill per day in the order directed on the pack. Use your back up birth control for at least 9 days.
- During Days 25 through 28, throw out the rest of the pack. Start a new pack on the same day or on the day you would normally start a new pack. Take one pill per day in the order directed on the pack. No back up birth control is needed.

If you miss a period for two months in a row, call your doctor because you might be pregnant.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose symptoms may include nausea, vomiting, and vaginal bleeding.

What to avoid

Do not smoke while taking this medicine, especially if you are older than 35 years of age.

Natazia will not protect you from sexually transmitted diseases - including HIV and AIDS. Using a condom is the only way to protect yourself from these diseases.

Natazia side effects

Get emergency medical help if you have **signs of an allergic reaction to Natazia**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop taking Natazia and call your doctor at once if you have:

- signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance;
- signs of a blood clot - chest pain, sudden cough, wheezing, coughing up blood, swelling or warmth in one or both legs;
- heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- liver problems - severe stomach pain, fever, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- sudden and severe pelvic pain;
- a change in the pattern or severity of migraine headaches;
- swelling in your hands, ankles, or feet;
- a breast lump; or
- symptoms of depression - sleep problems, weakness, tired feeling, mood changes.

Common Natazia side effects may include:

- light vaginal bleeding or spotting;
- nausea (especially during the first few months of taking this medicine), vomiting;
- breast pain or tenderness;
- weight gain; or
- problems with contact lenses.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Natazia side effects](#) (more detail)

Related/similar drugs

Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

Reviews & ratings

6.1 / 10

[2,624 Reviews](#)

[View more](#)

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

[685 Reviews](#)

[View more](#)

Depo Provera

Depo-Provera is used as contraception to prevent pregnancy. It is also used to reduce pain cause by ...

Reviews & ratings

4.2 / 10

[1,967 Reviews](#)

[View more](#)

Medroxyprogesterone

Medroxyprogesterone (Provera) treats conditions such as absent or irregular menstrual periods, or ...

Reviews & ratings

4.5 / 10

[2,469 Reviews](#)

[View more](#)

Nexplanon

Nexplanon is a long-term, reversible birth control implant that lasts for three years. Learn about ...

Reviews & ratings

5.0 / 10

[4,864 Reviews](#)

[View more](#)

Yaz

Yaz (drospirenone and ethinyl estradiol) is used as a contraceptive to prevent pregnancy. Includes ...

Reviews & ratings

5.9 / 10

[997 Reviews](#)

[View more](#)

Slynd

Slynd (drospirenone) is a birth control pill used for the prevention of pregnancy. Includes Slynd ...

Reviews & ratings

5.8 / 10

[386 Reviews](#)

[View more](#)

What other drugs will affect Natazia?

Many drugs can interact with dienogest and estradiol and make them less effective, which may result in pregnancy. Estradiol can also affect blood levels of certain other drugs, making them less effective or increasing side effects. You may need to use a back-up birth control method while using certain other drugs. Tell your doctor about all your current medicines and any you start or stop using. Not all possible interactions are listed in this medication guide.

 [Natazia drug interactions](#) (more detail)

Does Natazia interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Natazia

+

Enter a drug name

Add

More about Natazia (dienogest / estradiol)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(67\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [FDA approval history](#)
- [Drug class: contraceptives](#)
- [En español](#)

Professional resources

- [Natazia prescribing information](#)

Related treatment guides

- [Birth Control](#)
- [Heavy Menstrual Bleeding](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Natazia only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)


Copyright 1996-2025 Cerner Multum, Inc. Version: 7.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

 Risk data available

CSA Schedule*

N/A Not a controlled drug



Approval History

10+ years FDA approved 2010



User Reviews & Ratings

6.5 / 10

[67 Reviews](#)

Images

[Natazia estradiol valerate 3 mg \(DD\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)





[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.