

#### Home

Aspirin/caffeine

# Aspirin and caffeine

Generic name: aspirin and caffeine [ AS-pir-in-and-KAF-een ]

Brand names: Alka-Seltzer Hangover Relief, Alka-Seltzer Morning Relief, Anacin, Anacin Extra Strength, Bayer Back and Body, ... show all

16 brands

Dosage forms: oral powder for reconstitution (1000 mg-150 mg; 1000 mg-65 mg; 650 mg-65 mg; 845 mg-65 mg), ... show all 3 dosage

forms

Drug class: Analgesic combinations



Medically reviewed by Drugs.com on Jan 13, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

# What is aspirin and caffeine?

Aspirin and caffeine is a combination medicine used in adults and children at least 12 years old to temporarily relieve minor aches and pains such as headaches, muscle pain, cold, minor arthritis pain, menstrual cramps, and fever or body aches. Aspirin and caffeine is sometimes used in adults to relieve symptoms associated with hangover.

Aspirin and caffeine should not be used as a substitute for sleep.

Aspirin and caffeine may also be used for purposes not listed in this medication guide.

# Aspirin and caffeine side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

Aspirin and caffeine may cause serious side effects. Stop using aspirin and caffeine and call your doctor at once if you have:

- hearing loss or a ringing in your ears;
- · redness or swelling;
- severe nausea, vomiting, or stomach pain;
- nervousness, irritability, rapid heartbeat;
- sleep problems (insomnia);
- · fever lasting longer than 3 days;
- · pain lasting longer than 10 days; or
- stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds.

Less serious side effects may be more likely, or you may have none at all.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Aspirin / caffeine side effects (more detail)

# Related/similar drugs

# **Oxycodone**

Oxycodone is an opioid analgesic used to treat moderate to severe pain; it has a high potential for ...

### **Reviews & ratings**

6.9 / 10

1,142 Reviews

View more

## **Naproxen**

Naproxen is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

### **Reviews & ratings**

6.8 / 10

683 Reviews

View more

## FEATURED

# **Ozempic**

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

### **Reviews & ratings**

6.6 / 10

1,504 Reviews

View more

### **Paracetamol**

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-thecounter painkiller and fever ...

### **Reviews & ratings**

6.1 / 10

13 Reviews

#### View more

# **Hydroxyzine**

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

#### **Reviews & ratings**

5.9 / 10

1,404 Reviews

#### View more

### **Tramadol**

Tramadol is an opioid medication that may be used to treat moderate to moderately severe chronic ...

#### **Reviews & ratings**

7.0 / 10

1,876 Reviews

## View more

# Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

# Reviews & ratings

6.0 / 10

618 Reviews

### View more

## **Meloxicam**

Meloxicam is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

### **Reviews & ratings**

6.3 / 10

600 Reviews

....

#### View more

## **Aspirin**

Aspirin is used to treat mild to moderate pain and to reduce fever or inflammation. Learn about ...

#### **Reviews & ratings**

7.8 / 10

51 Reviews

View more

# **Tylenol**

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

#### **Reviews & ratings**

5.8 / 10

30 Reviews

View more

# Warnings

Aspirin may cause stomach or intestinal bleeding, which can be fatal.

Aspirin can also cause a severe allergic reaction. Get emergency medical help if you have **signs of an allergic reaction**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

Limit the use of caffeine-containing medications, coffee, tea, cola, energy drinks or other sources of caffeine while taking aspirin and caffeine. Using too much caffeine may cause nervousness, irritability, sleeplessness, and rapid heartbeats.

Using this medicine in a child or teenager with flu symptoms or chickenpox can cause a serious or fatal condition called Reye's syndrome.

# Before taking this medicine

You should not use aspirin and caffeine if you are allergic to it, or if you have:

• an allergy to an NSAID (nonsteroidal anti-inflammatory drug) such as Advil, Motrin, Aleve, naproxen, ibuprofen, celecoxib, diclofenac, indomethacin, meloxicam, and others.

Do not give this medication to a child or teenager with a fever, flu symptoms, or chicken pox. Aspirin and caffeine can cause a condition called Reye's syndrome, which can be serious and sometimes fatal in children.

Ask a doctor or pharmacist if this medicine is safe to use if you have or have ever had:

- stomach ulcers or bleeding problems;
- heartburn;
- if you are age 60 or older;
- a habit of drinking more than 3 alcoholic beverages per day;
- · heart disease, high blood pressure;
- · asthma;
- · gout;
- · diabetes:
- if you are on a low-salt diet; or
- liver or kidney disease.

This medicine may contain phenylalanine and could be harmful if you have phenylketonuria (PKU).

Do not use aspirin and caffeine during pregnancy unless your doctor tells you to. Using an NSAID in the last 20 weeks can harm the unborn baby or cause pregnancy complications.

Ask a doctor before using this medicine if you are breastfeeding.

# How should I take aspirin and caffeine?

Use exactly as directed on the label, or as prescribed by your doctor.

Aspirin and caffeine is usually taken every 6 hours with a full glass of water.

Taking more than your recommended dose will not make aspirin and caffeine more effective, and may cause serious side effects.

Take with food if aspirin and caffeine upsets your stomach.

Limit the use of caffeine-containing medications, coffee, tea, cola, energy drinks or other sources of caffeine while taking this product.

Call your doctor if your symptoms do not improve, or if they get worse.

If you need surgery, tell the surgeon ahead of time that you are using aspirin and caffeine. You may need to stop using the medicine for a short time.

Store at room temperature away from moisture and heat.

Detailed Aspirin / caffeine dosage information

# What happens if I miss a dose?

Aspirin and caffeine is used when needed. If you are on a dosing schedule, skip any missed dose. **Do not** use two doses at one time.

# What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

# What should I avoid while taking aspirin and caffeine?

Drinking alcohol may increase your risk of stomach bleeding.

Ask a doctor or pharmacist before using other medicines for pain, fever, swelling, or cold/flu symptoms. They may contain ingredients similar to aspirin and caffeine (such as ibuprofen, ketoprofen, or naproxen).

# What other drugs will affect aspirin and caffeine?

Ask a doctor or pharmacist before using aspirin and caffeine with any other medicines, especially:

- gout medication;
- a diuretic or "water pill";
- arthritis medication;
- · diabetes medications;
- steroids:
- a blood thinner--warfarin, Coumadin, Jantoven; or
- NSAIDs (nonsteroidal anti-inflammatory drugs)--aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others.

This list is not complete. Other drugs may affect aspirin and caffeine, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Aspirin / caffeine drug interactions (more detail)

# Does aspirin/caffeine interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

aspirin/caffeine	
+	
Enter a drug name	Add

- Check interactions
- · Compare alternatives
- Reviews (310)
- Imprints, shape & color data
- · Side effects
- Dosage information
- During pregnancy
- Drug class: analgesic combinations

## **Patient resources**

#### Other brands

Anacin, BC Arthritis, Anacin Max Strength, Stanback, ... +7 more

# Related treatment guides

- Pain
- Fever
- Hangover

# **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 4.01.

## **DRUG STATUS**

### **Availability**

Over the counter

#### **Pregnancy & Lactation**

& Risk data available

# **CSA Schedule\***

N/A Not a controlled drug

# **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





#### **About**

About Drugs.com

Advertising policy

Attribution & citations

## Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

### **Support**

Help center

**Sitemap** 

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.