

Home

Amlodipine/valsartan

Amlodipine and valsartan

Generic name: amlodipine and valsartan [am-LOE-de-peen-val-SAR-tan]

Brand name: Exforge

Dosage form: oral tablet (10 mg-160 mg; 10 mg-320 mg; 5 mg-160 mg; 5 mg-320 mg)

Drug class: Angiotensin II inhibitors with calcium channel blockers



Medically reviewed by Drugs.com on Jun 5, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is amlodipine and valsartan?

Amlodipine and valsartan is a combination medicine used alone or together with other medicines to treat high blood pressure (hypertension) in adults.

Lowering blood pressure may lower your risk of a stroke or heart attack.

Amlodipine and valsartan may also be used for purposes not listed in this medication guide.

Amlodipine and valsartan side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, difficulty breathing, swelling of your face, lips, tongue, or throat.

In rare cases, when you first start taking amlodipine and valsartan, your chest pain may get worse or you could have a heart attack. Seek emergency medical attention or call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea, or sweating.

Amlodipine and valsartan may cause serious side effects. Call your doctor at once if you have:

- a light-headed feeling, like you might pass out;
- heart problems--swelling, rapid weight gain, feeling short of breath;
- · kidney problems--swelling, urinating less, feeling tired or short of breath; or
- high blood potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement.

Common side effects include:

- · swelling in your hands or feet;
- · spinning sensation, dizziness; or

· cold symptoms such as stuffy nose, sneezing, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Amlodipine / valsartan side effects (more detail)

Related/similar drugs

Vasotec

Vasotec (enalapril) is used to treat high blood pressure, heart failure and kidney problems caused ...

Reviews & ratings

4 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Lisinopril

Lisinopril is an ACE inhibitor. It is used to treat high blood pressure, congestive heart failure ...

Reviews & ratings

5.0 / 10

626 Reviews

View more

Atenolol

Atenolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

Reviews & ratings

- - - - -

View more

Valsartan

Valsartan is an angiotensin II receptor blocker (ARB) that may be used to treat high blood pressure ...

Reviews & ratings

5.7 / 10

152 Reviews

View more

Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

Reviews & ratings

5.2 / 10

185 Reviews

View more

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

6.4 / 10

56 Reviews

View more

Furosemide

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

Reviews & ratings

6.8 / 10

121 Reviews

View more

Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

Reviews & ratings

4.5 / 10

578 Reviews

View more

Warnings

Do not use if you are pregnant. Stop taking this medicine and tell your doctor right away if you become pregnant.

Tell your doctor about all your other medicines. Some drugs should not be used with amlodipine and valsartan.

Before taking this medicine

You should not use amlodipine and valsartan if you are allergic to it.

If you have diabetes, do not take amlodipine and valsartan together with any medication that contains aliskiren (a blood pressure medicine).

Tell your doctor if you have ever had:

- heart problems or low blood pressure;
- · a heart attack;
- a severe allergic reaction called angioedema to another blood pressure medicine;
- · vomiting or having a lot of diarrhea;
- · liver disease; or
- kidney disease (or if you are on dialysis).

You may also need to avoid taking amlodipine and valsartan with aliskiren if you have kidney disease.

Do not use if you are pregnant. Stop using the medicine and tell your doctor right away if you become pregnant. Amlodipine and valsartan can cause injury or death to the unborn baby if you take the medicine during your second or third trimester.

Do not breastfeed.

How should I take amlodipine and valsartan?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Amlodipine and valsartan is usually taken once per day.

You may take this medication with or without food.

Your blood pressure will need to be checked often and you may need frequent blood tests. Your treatment may also include diet, exercise, lifestyle changes, and other medications. Follow your doctor's instructions very carefully.

Your blood pressure may get too low during prolonged illness. Call your doctor if you are sick with vomiting or diarrhea.

Tell your doctor if you have a planned surgery.

Keep using amlodipine and valsartan even if you feel well. High blood pressure often has no symptoms.

Store at room temperature away from moisture and heat.

Amlodipine and valsartan dosing information

Usual Adult Dose for Hypertension:

Initial therapy: Amlodipine 5 mg-Valsartan 160 mg orally once a day

Add-on/Replacement therapy: Amlodipine 5 to 10 mg-Valsartan 160 to 320 mg orally once a day

Comments:

- -May increase dose after 1 to 2 weeks of therapy.
- -A patient who experiences dose-limiting adverse reactions on either component alone may be switched to this drug containing a lower dose of that component in combination with the other. If blood pressure remains uncontrolled after 3 to 4 weeks, may titrate up to a maximum of amlodipine 10 mg-valsartan 320 mg orally once a day.
- Detailed Amlodipine / valsartan dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include rapid heartbeats, low blood pressure, redness or warmth in your arms or legs, or fainting.

What should I avoid while taking amlodipine and valsartan?

Do not take potassium supplements or use salt substitutes, unless your doctor has told you to.

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Avoid driving or hazardous activity until you know how amlodipine and valsartan will affect you. Your reactions could be impaired.

Drinking alcohol with this medicine can cause side effects.

What other drugs will affect amlodipine and valsartan?

Sometimes it is not safe to use certain medicines at the same time. Some drugs can affect your blood levels of other drugs you use, which may increase side effects or make the medicines less effective.

Amlodipine and valsartan can harm your kidneys, especially if you also use certain medicines for infections, cancer, osteoporosis, organ transplant rejection, high blood pressure, or pain or arthritis (including Advil, Motrin, and Aleve).

Tell your doctor about all your current medicines. Many drugs can affect amlodipine and valsartan, especially:

- cyclosporine;
- sildenafil:
- rifampin;
- · ritonavir; or
- a heart or blood pressure medication, especially a diuretic ('water pill').

This list is not complete and many other drugs may affect amlodipine and valsartan. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Amlodipine / valsartan drug interactions (more detail)

Does amlodipine/valsartan interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

| amlodipine/valsartan | |
|----------------------|-----|
| + | |
| Enter a drug name | Add |

More about amlodipine / valsartan

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (60)
- Drug images
- Side effects

- Dosage information
- During pregnancy
- Drug class: angiotensin II inhibitors with calcium channel blockers
- En español

Patient resources

Other brands

Exforge

Professional resources

· Amlodipine and Valsartan prescribing information

Other brands

Exforge

Related treatment guides

• High Blood Pressure

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 13.01.

DRUG STATUS

Availability Rx Prescription only Pregnancy & Lactation Risk data available CSA Schedule* N/A Not a controlled drug Approval History Drug history at FDA

User Reviews & Ratings

60 Reviews

Images

Amlodipine Besylate and Valsartan 10 mg / 320 mg (MX44)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





