

Home

2. Micardis HCT

Micardis HCT ⊲

Generic name: hydrochlorothiazide and telmisartan [HYE-droe-KLOR-oh-THYE-a-zide-and-TEL-mi-SAR-tan]

Drug class: Angiotensin II inhibitors with thiazides



Medically reviewed by Drugs.com on Feb 5, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Micardis HCT?

Hydrochlorothiazide is a diuretic (water pill). Telmisartan is an angiotensin II receptor blocker (sometimes called an ARB blocker).

Micardis HCT is a combination medicine is used to treat high blood pressure (hypertension). Lowering blood pressure may lower your risk of a stroke or heart attack.

Micardis HCT may also be used for purposes not listed in this medication guide.

Micardis HCT side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

In rare cases, Micardis HCT can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine.

Also call your doctor at once if you have:

- · eye pain, vision problems;
- · a light-headed feeling, like you might pass out;
- jaundice (yellowing of the skin or eyes);
- swelling in your hands or feet, unusual weight gain;
- · easy bruising, unusual bleeding; or
- signs of an electrolyte imbalance--dry mouth, increased thirst, drowsiness, confusion, feeling restless, vomiting, muscle pain or weakness, lack of energy, fast heartbeats, little or no urine, or a seizure.

Common side effects of Micardis HCT may include:

- · dizziness, feeling tired;
- nausea, diarrhea;
- · back pain; or
- cold or flu symptoms.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



Related/similar drugs

Vasotec

Vasotec (enalapril) is used to treat high blood pressure, heart failure and kidney problems caused ...

Reviews & ratings

4 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Lisinopril

Lisinopril is an ACE inhibitor. It is used to treat high blood pressure, congestive heart failure ...

Reviews & ratings

5.0 / 10

626 Reviews

View more

Atenolol

Atenolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

Reviews & ratings

7.2 / 10

276 Reviews

View more

Valsartan

Valsartan is an angiotensin II receptor blocker (ARB) that may be used to treat high blood pressure ...

Reviews & ratings

5.7 / 10

152 Reviews

View more

Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

Reviews & ratings

5.2 / 10

185 Reviews

View more

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

6.4 / 10

56 Reviews

View more

Furosemide

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

Reviews & ratings

6.8 / 10

121 Reviews

View more

Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

Reviews & ratings

4.5 / 10

578 Reviews

View more

Warnings

Do not use if you are pregnant. Stop using the medicine and tell your doctor right away if you become pregnant.

You should not use this medicine if you are unable to urinate.

If you have diabetes, do not use Micardis HCT together with any medication that contains aliskiren (a blood pressure medicine).

Before taking this medicine

You should not use Micardis HCT if you are allergic to hydrochlorothiazide or telmisartan, if you are unable to urinate.

If you have diabetes, do not use Micardis HCT together with any medication that contains aliskiren (a blood pressure medicine).

You may also need to avoid taking Micardis HCT with aliskiren if you have kidney disease.

Tell your doctor if you have ever had:

- · kidney disease;
- liver disease;
- glaucoma;
- low or high levels of magnesium or potassium in your blood;
- asthma or allergies;
- lupus;
- · diabetes; or

· an allergy to penicillin or sulfa drugs.

Do not use if you are pregnant. Stop using the medicine and tell your doctor right away if you become pregnant. Micardis HCT can cause injury or death to the unborn baby if you take the medicine during your second or third trimester.

You should not breastfeed while using this medicine.

How should I take Micardis HCT?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You may take this medicine with or without food.

Your blood pressure will need to be checked often.

Call your doctor if you are sick with vomiting or diarrhea, or if you are sweating more than usual. You can easily become dehydrated while taking Micardis HCT. This can lead to very low blood pressure, a serious electrolyte imbalance, or kidney failure.

If you need surgery or medical tests, tell the surgeon ahead of time that you are using this medicine.

It may take up to 4 weeks for this medication to control your blood pressure. **Keep using this medicine as directed, even if you feel well.** High blood pressure often has no symptoms. You may need to use blood pressure medicine for the rest of your life.

Store at room temperature away from moisture and heat. Keep each tablet in its blister pack until you are ready to take it. Gently push the tablet through the foil to remove it.

Detailed Micardis HCT dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include fast or slow heart rate, feeling light-headed, or fainting.

What should I avoid while taking Micardis HCT?

Hydrochlorothiazide may increase your risk of skin cancer. Avoid sunlight or tanning beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors. Your doctor may want you to have skin examinations on a regular basis.

Do not use potassium supplements or salt substitutes, unless your doctor has told you to.

Drinking alcohol can further lower your blood pressure and may cause side effects.

Avoid becoming overheated or dehydrated during exercise, in hot weather, or by not drinking enough fluids.

What other drugs will affect Micardis HCT?

If you also take cholestyramine or colestipol, avoid taking these medications within 4 hours before or 4 hours after you take Micardis HCT.

Tell your doctor about all your other medicines, especially:

- any other heart or blood pressure medications;
- · digoxin;
- · lithium;
- a diuretic or "water pill";
- · insulin or oral diabetes medicine;
- · opioid pain medicine;
- · a sleeping pill;
- steroid medicine or adrenocorticotrophic hormone (ACTH);
- a barbiturate such as butabarbital, phenobarbital, or secobarbital; or
- NSAIDs (nonsteroidal anti-inflammatory drugs)--aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others.

This list is not complete. Other drugs may affect Micardis HCT, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Micardis HCT drug interactions (more detail)

Does Micardis HCT interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Micardis HCT
+
Enter a drug name
Add

More about Micardis HCT (hydrochlorothiazide / telmisartan)

- Check interactions
- · Compare alternatives
- Pricing & coupons

- Reviews (16)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Drug class: angiotensin II inhibitors with thiazides
- En español

Patient resources

· Micardis HCT advanced reading

Professional resources

• Micardis HCT prescribing information

Other formulations

• Micardis

Related treatment guides

• High Blood Pressure

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.01.

DRUG STATUS

Availability


Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug



User Reviews & Ratings

6.9 / 10

16 Reviews

Images

Micardis HCT 12.5 mg / 80 mg (LOGO H8)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

V