

Home

Q-Tapp DM

Q-Tapp DM

Generic name: Brompheniramine, Pseudoephedrine, and Dextromethorphan [brome-fen-IR-a-meen, soo-doe-e-FED-rin, & deks-troe-meth-OR-fan]

Brand names: Bromfed DM, Brotapp-DM, Neo DM, Q-Tapp DM, TGQ 30PSE/3BRM/15DM

Drug class: Upper respiratory combinations

Medically reviewed by Drugs.com. Last updated on Apr 16, 2024.

Uses Before taking Warnings Dosage Side effects Overdose



The Q-Tapp DM brand name has been **discontinued** in the U.S. If generic versions of this product have been approved by the FDA, there may be generic equivalents available.

Uses of Q-Tapp DM:

- · It is used to treat nose stuffiness.
- It is used to ease allergy signs.
- It is used to relieve coughing.

What do I need to tell my doctor BEFORE I take Q-Tapp DM?

- If you have an allergy to brompheniramine, pseudoephedrine, dextromethorphan, or any other part of Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan).
- If you are allergic to Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan); any part of Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan); or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have taken certain drugs for depression or Parkinson's disease in the last 14 days. This includes isocarboxazid, phenelzine, tranylcypromine, selegiline, or rasagiline. Very high blood pressure may happen.
- If you are taking any of these drugs: Linezolid or methylene blue.
- If you have a cough with a lot of mucus.
- If you have a long-term cough caused by smoking or being around smoke, or lung problems like asthma or emphysema.
- If you are breast-feeding. Do not breast-feed while you take Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan).

This is not a list of all drugs or health problems that interact with Q-Tapp DM (brompheniramine, pseudoephedrine, and

dextromethorphan).

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take Q-Tapp DM?

- Tell all of your health care providers that you take Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan). This includes your doctors, nurses, pharmacists, and dentists.
- Do not take more than what your doctor told you to take. Taking more than you are told may raise your chance of very bad side effects.
- Do not take Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) for longer than you were told by your doctor.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) affects you.
- Avoid drinking alcohol while taking Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan).
- Talk with your doctor before you use other drugs and natural products that slow your actions.
- If you are 65 or older, use Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) with care. You
 could have more side effects.
- Use with care in children. Talk with the doctor.
- Do not use to make a child sleepy. Talk with the doctor.
- This medicine may not be for use in children of all ages. Talk with the doctor before giving Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) to a child.
- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) while you are pregnant.

How is this medicine (Q-Tapp DM) best taken?

Use Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with or without food. Take with food if it causes an upset stomach.
- Measure liquid doses carefully. Use the measuring device that comes with Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan). If there is none, ask the pharmacist for a device to measure Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan).

What do I do if I miss a dose?

- If you take Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) on a regular basis, take a
 missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.

- Do not take 2 doses at the same time or extra doses.
- Many times Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan) is taken on an as needed basis. Do not take more often than told by the doctor.
- Detailed Q-Tapp DM dosage information

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever;
 wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Q-Tapp DM side effects (more detail)

Related/similar drugs

Mucinex D

Mucinex D (guaifenesin and pseudoephedrine) is used to treat nasal and sinus congestion, and to ...

Reviews & ratings

6.1 / 10

224 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

What are some other side effects of Q-Tapp DM?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your

doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- · Dizziness.
- · Feeling nervous and excitable.
- Trouble sleeping.
- · Feeling sleepy.
- Dry mouth.
- Dry nose.
- Thickening of mucus in nose or throat.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

If OVERDOSE is suspected:

If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

How do I store and/or throw out Q-Tapp DM?

- Store at room temperature. Do not freeze.
- · Keep lid tightly closed.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so.
 Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Consumer Information Use and Disclaimer

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about Q-Tapp DM (brompheniramine, pseudoephedrine, and dextromethorphan), please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

More about Q-Tapp DM (brompheniramine / dextromethorphan / pseudoephedrine)

- · Check interactions
- · Compare alternatives
- Reviews (2)
- · Side effects
- Dosage information
- Drug class: upper respiratory combinations

Patient resources

Other brands

Resperal-DM, Allanhist PDX Drops, Anaplex DM, Andehist DM NR Syrup, ... +2 more

Professional resources

• Brompheniramine, Pseudoephedrine, Dextromethorpan Syrup prescribing information

Other brands

Bromfed DM

Related treatment guides

• Cough and Nasal Congestion

Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

DRUG STATUS

Availability

Discontinued

CSA Schedule*

N/A Not a controlled drug

WADA Class

റ്റ് Anti-Doping Classification

User Reviews & Ratings

Review this drug

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





