

Home

2. Amrix

Amrix ⊲»

Generic name: cyclobenzaprine [sye-kloe-BEN-za-preen]

Brand names: Amrix, Fexmid

Drug class: Skeletal muscle relaxants



Medically reviewed by Drugs.com on Aug 4, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is Amrix?

Amrix is a muscle relaxant. It works by blocking nerve impulses (or pain sensations) that are sent to your brain.

Amrix is used together with rest and physical therapy to relieve muscle spasms caused by painful conditions such as an injury.

Amrix may also be used for purposes not listed in this medication guide.

Amrix side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Amrix may cause serious side effects. Stop using Amrix and call your doctor at once if you have:

- · fast or irregular heartbeats;
- chest pain or pressure, pain spreading to your jaw or shoulder; or
- sudden numbness or weakness (especially on one side of the body), slurred speech, balance problems.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Serious side effects may be more likely in older adults.

Common side effects of Amrix may include:

- · drowsiness, tiredness;
- · headache, dizziness;
- dry mouth; or

• upset stomach, nausea, constipation.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Amrix side effects (more detail)

Related/similar drugs

Ibuprofen

Ibuprofen (Advil, Motrin) treats minor aches and pains caused by the common cold, headaches ...

Reviews & ratings

7.3 / 10

240 Reviews

View more

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-thecounter painkiller and fever ...

Reviews & ratings

6.1 / 10

13 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

- - / / -

View more

Flexeril

Flexeril (cyclobenzaprine) is a muscle relaxant used to treat skeletal muscle conditions such as ...

Reviews & ratings

6.1 / 10

237 Reviews

View more

Robaxin

Robaxin (methocarbamol) is used to treat skeletal muscle conditions such as pain or injury ...

Reviews & ratings

6.3 / 10

80 Reviews

View more

Advil

Advil (ibuprofen) reduces fever, pain and inflammation caused by headaches, toothaches, arthritis ...

Reviews & ratings

6.9 / 10

41 Reviews

View more

Valium

Valium is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.6 / 10

341 Reviews

View more

Diazepam

Diazepam is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.4 / 10

884 Reviews

View more

Zanaflex

Zanaflex (tizanidine) is a short-acting muscle relaxer used to treat spasticity by temporarily ...

Reviews & ratings

7.6 / 10

116 Reviews

View more

Warnings

You should not use Amrix if you have a thyroid disorder, heart block, congestive heart failure, a heart rhythm disorder, or you have recently had a heart attack.

Do not use Amrix if you have taken an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, or tranylcypromine.

Before taking this medicine

You should not use Amrix if you are allergic to it, or if you have:

- a thyroid disorder;
- heart block, heart rhythm disorder, congestive heart failure; or
- if you have recently had a heart attack.

Amrix is not approved for use by anyone younger than 15 years old.

Do not use Amrix if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Some medicines can interact with Amrix and cause a serious condition called **serotonin syndrome**. Be sure your doctor knows if you also take stimulant medicine, opioid medicine, herbal products, or medicine for depression, mental illness, Parkinson's disease, migraine headaches, serious infections, or prevention of nausea and vomiting. **Ask your doctor before making any changes in how or when you take your medications**.

Tell your doctor if you have ever had:

- · liver disease:
- · glaucoma;
- · enlarged prostate; or
- problems with urination.

It is not known whether Amrix will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It may not be safe to breast-feed while using this medicine. Ask your doctor about any risk.

Older adults may be more sensitive to the effects of this medicine.

How should I take Amrix?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Amrix is usually taken once daily for only 2 or 3 weeks. Follow your doctor's dosing instructions very carefully.

Swallow the capsule whole and do not crush, chew, break, or open it.

Take the medicine at the same time each day.

Call your doctor if your symptoms do not improve after 3 weeks, or if they get worse.

Store at room temperature away from moisture, heat, and light.

Detailed Amrix dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. An overdose of Amrix can be fatal.

Overdose symptoms may include severe drowsiness, vomiting, fast heartbeats, tremors, agitation, or hallucinations.

What should I avoid while taking cyclobenzaprine?

Avoid driving or hazardous activity until you know how Amrix will affect you. Your reactions could be impaired.

Avoid drinking alcohol. Dangerous side effects could occur.

What other drugs will affect Amrix?

Using Amrix with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- bupropion (Zyban, for smoking cessation);
- meperidine;
- · tramadol;
- verapamil;
- cold or allergy medicine that contains an antihistamine (Benadryl and others);
- medicine to treat Parkinson's disease;
- medicine to treat excess stomach acid, stomach ulcer, motion sickness, or irritable bowel syndrome;
- medicine to treat overactive bladder; or
- bronchodilator asthma medication.

This list is not complete. Other drugs may affect Amrix, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Amrix drug interactions (more detail)

Does Amrix interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

Does cyclobenzaprine make you sleepy?

How long does it take for cyclobenzaprine to work?

How much cyclobenzaprine can I take?

∨

How long does it take for Flexeril to work?

More FAQ

- How long does Flexeril stay in your system?
- Can I take cyclobenzaprine and Tylenol together?

More about Amrix (cyclobenzaprine)

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (48)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- · Generic availability
- FDA approval history
- Drug class: skeletal muscle relaxants
- Breastfeeding
- En español

Patient resources

· Amrix advanced reading

Other brands

Flexeril, Fexmid

Professional resources

- Amrix prescribing information
- Cyclobenzaprine (AHFS Monograph)

Other brands

Flexeril, Fexmid

Related treatment guides

• Muscle Spasm

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 7.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্য Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

7.0 / 10

48 Reviews

Images

Amrix 15 mg (ECR 15 ECR 15)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

×