

Home

Follicle stimulating hormone

Follicle stimulating hormone

Generic name: follicle stimulating hormone [FOL-ik-al-STIM-ue-lay-ting-HOR-mone]

Brand names: Follistim AQ Cartridge, Gonal-F, Gonal-f RFF, Follistim, Follistim AQ, ... show all 10 brands

Dosage forms: subcutaneous powder for injection (1050 intl units; alpha 450 intl units; alpha 75 intl units), ... show all 2 dosage forms

Drug class: Gonadotropins



Medically reviewed by Drugs.com on Feb 4, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is follicle stimulating hormone?

Follicle stimulating hormone is used to treat infertility **in women** who cannot ovulate. follicle stimulating hormone is not effective in women with primary ovarian failure (when the ovaries are unable to produce an egg).

Follicle stimulating hormone is also used **in men** to stimulate sperm production. This medicine is not effective in men with primary testicular failure (when the testicles are unable to produce sperm).

Follicle stimulating hormone is often used together with another medication called human chorionic gonadotropin (hCG).

Follicle stimulating hormone may also be used for purposes not listed in this medication guide.

Follicle stimulating hormone side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Some women using this medicine develop ovarian hyperstimulation syndrome (OHSS), a potentially life-threatening condition. Stop using this medicine and call your doctor right away if you have symptoms of OHSS during or shortly after using follicle stimulating hormone:

- severe pain in your lower stomach, nausea, vomiting, diarrhea;
- · bloating, rapid weight gain;
- little or no urination; or
- · trouble breathing.

Also call your doctor at once if you have:

- · severe pelvic pain on one side;
- · a cough with bloody mucus;

- fever, sore throat, swollen glands;
- a lump or swelling in your testicles;
- signs of acute respiratory distress syndrome--fever with shortness of breath or rapid breathing; or
- signs of a blood clot--sudden numbness or weakness, problems with vision or speech, swelling or redness in an arm or leg.

Common side effects of follicle stimulating hormone may include:

- redness, swelling, or bruising where an injection was given;
- · an ovarian cyst;
- headache, tiredness;
- breast pain or swelling;
- · stuffy nose, sneezing, sinus pain;
- · acne; or
- nausea.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

• Follicle stimulating hormone side effects (more detail)

Related/similar drugs

Clomid

Clomid (clomiphene) is used to stimulate ovulation when the ovaries can produce a follicle but ...

Reviews & ratings

7.8 / 10

224 Reviews

View more

AndroGel

AndroGel is used to treat conditions in men that result from a lack of natural testosterone. Learn ...

Reviews & ratings

7.5 / 10

112 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Xyosted

Xyosted is used to treat men who have low or no testosterone due to certain medical conditions ...

Reviews & ratings

4 Reviews

View more

HCG

HCG is used to cause ovulation and to treat infertility in women. Learn about side effects ...

Reviews & ratings

7.2 / 10

13 Reviews

View more

Warnings

You should not use follicle stimulating hormone if you have an untreated or uncontrolled endocrine disorder (thyroid, pituitary gland, or adrenal gland), heavy or abnormal vaginal bleeding that has not been checked by a doctor, an ovarian cyst, or cancer of the breast, ovary, uterus, testicle, hypothalamus, or pituitary gland.

Do not use this medicine if you are pregnant. You may need to have a negative pregnancy test before starting this treatment.

Before taking this medicine

Your doctor will perform blood tests to make sure you do not have conditions that would prevent you from safely using follicle stimulating hormone.

You should not use follicle stimulating hormone if you are allergic to it, or if you have:

- an untreated or uncontrolled disorder of the thyroid, pituitary gland, or adrenal glands;
- · heavy or abnormal vaginal bleeding that has not been checked by a doctor;
- an ovarian cyst;
- · cancer of the breast, ovary, uterus, testicle, hypothalamus, or pituitary gland; or
- if you are allergic to benzyl alcohol, follitropin, sucrose, neomycin, sodium phosphate, or streptomycin.

Do not use this medicine if you are already pregnant. You may need to have a negative pregnancy test before starting this treatment. Tell your doctor right away if you become pregnant.

Tell your doctor if you have ever had:

- · unusual vaginal bleeding;
- · polycystic ovary disease;
- twisting (torsion) of your ovary;
- · a stroke or blood clot;
- · asthma; or
- stomach surgery.

Fertility treatment may increase your chance of having multiple births (twins, triplets, etc). These are high-risk pregnancies both for the mother and the babies. Talk to your doctor if you have concerns about this risk.

You should not breastfeed while using follicle stimulating hormone.

How should I use follicle stimulating hormone?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You must remain under the care of a doctor while using follicle stimulating hormone.

Follicle stimulating hormone is injected under the skin. A healthcare provider may teach you how to properly use the medication by yourself.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you don't understand all instructions.

You may need to mix follicle stimulating hormone with a liquid (diluent) before using it. When using injections by yourself, be sure you understand how to properly mix and store the medicine.

Prepare an injection only when you are ready to give it. **Do not use if the medicine looks cloudy or has particles in it.** Call your pharmacist for new medicine.

To get pregnant, you may need to have sex daily for several days in a row. The timing of sex and medication dosing is important for follicle stimulating hormone to work.

You will need frequent medical tests.

If you use this medicine to get pregnant, you may also need occasional ultrasound exams. **Based on your blood work,** you may need to stop having sex for a short time. Follow your doctor's instructions very carefully.

Storing this medicine unopened (not in use):

- Refrigerate (do not freeze) and use until expiration date; or
- Store at room temperature protected from light, and use within 3 months or until expiration date.

Storing this medicine opened (in use):

- Store in a refrigerator (do not freeze), or at room temperature protected from light.
- Use within 28 days.

Take the medicine out of the refrigerator and allow it to reach room temperature before injecting your dose. Do not heat the medicine.

Use a needle and syringe only once and then place them in a puncture-proof "sharps" container. Follow state or local laws about how to dispose of this container. Keep it out of the reach of children and pets.

1 Detailed Follicle stimulating hormone dosage information

What happens if I miss a dose?

Call your doctor for instructions if you miss a dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while using follicle stimulating hormone?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

What other drugs will affect follicle stimulating hormone?

Other drugs may affect follicle stimulating hormone, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all your current medicines and any medicine you start or stop using.

• Follicle stimulating hormone drug interactions (more detail)

Does follicle stimulating hormone interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

follicle stimulating hormone

Follicle stimulating hormone Biosimilars

Biosimilar and interchangeable products are biological products that are highly similar to and have no clinically meaningful differences from the reference product.

Reference products

These are biological products that have already been approved by the FDA, against which biosimilar products are compared. There are 6 for follicle stimulating hormone.

Follistim (follitropin	heta) - Organo	n IISA Inc	a subsidiary	of Marck & Co. li	nc
FOIIISLIM (TOIIILEODIN	i beta) - Ordano	n USA Inc	a subsidiary	oi werck & Co II	IIC.

romstilli (toliiti opiii beta) - Organori OSA ilic., a subsidiary of werck & Co., ilic.					
	Formulation type	Strength			
	Single-Dose Vial	150 IU Discontinued, Voluntarily Revoked			
	Single-Dose Vial	75 IU Discontinued, Voluntarily Revoked			
	Follistim				
	Follistim AQ (follitropin beta) - Organo	on USA Inc., a subsidiary of Merck & Co., Inc.	~		
Follistim AQ Cartridge (follitropin beta) - Organon USA LLC, a subsidiary of Organon & Co.			~		
Gonal-f (follitropin alfa) - EMD Serono, Inc.			~		
	Gonal-f REF (follitropin alfa) - EMD Serono Inc				
Gonal-f RFF (follitropin alfa) - EMD Serono, Inc.					
	Gonal-f RFF RediJect (follitropin alfa)	- EMD Serono, Inc.	~		
	zana				

More about follicle stimulating hormone

- · Check interactions
- · Compare alternatives
- Reviews (3)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: gonadotropins
- En español

Patient resources

- · Follitropin alfa advanced reading
- Follitropin beta (Advanced Reading)
- Follitropin Alfa
- Follitropin Beta

Other brands

Gonal-f, Gonal-f RFF Redi-ject, Follistim AQ, Gonal-f RFF, Follistim AQ Cartridge

Professional resources

Other brands

Gonal-f, Follistim AQ, Gonal-f RFF

Related treatment guides

- · Hypogonadism, Male
- Follicle Stimulation
- Ovulation Induction

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 6.01.

DRUG STATUS

Availability Rx Prescription only Pregnancy & Lactation Risk data available CSA Schedule* N/A Not a controlled drug Approval History Drug history at FDA

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

Х