

[Home](#)

2. Fioricet with Codeine

Fioricet with Codeine

Generic name: [acetaminophen](#), [butalbital](#), [caffeine](#), and [codeine](#) [*a-SEET-a-MIN-o-fen*, *bue-TAL-bi-tal*, *KAF-eeen*, and *KOE-deen*]

Drug class: [Narcotic analgesic combinations](#)



[Medically reviewed](#) by Drugs.com on Jan 20, 2025. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is this Fioricet with Codeine?

Codeine is an opioid pain medication. **Acetaminophen** is a less potent pain reliever. **Butalbital** is a barbiturate. **Caffeine** is a stimulant.

Fioricet with Codeine is a combination medicine used to treat tension headaches.

Fioricet with Codeine may also be used for purposes not listed in this medication guide.

Fioricet with Codeine side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains [acetaminophen](#).

Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up.

Fioricet with Codeine may cause serious side effects. Call your doctor at once if you have:

- noisy breathing, sighing, shallow breathing;
- a [slow heart rate](#) or weak pulse;
- extreme drowsiness, feeling like you might pass out;
- chest pain, fast or pounding heart rate, feeling short of breath;
- confusion, unusual thoughts or behavior;
- **high levels of serotonin in the body**--agitation, hallucinations, fever, sweating, shivering, [fast heart rate](#), muscle

stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea;

- **low cortisol levels**-- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness; or
- **liver problems**--nausea, upper stomach pain, itching, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Serious breathing problems may be more likely in older adults and those who are debilitated or have wasting syndrome or chronic breathing disorders.

Common side effects include:

- drowsiness, dizziness, feeling "drunk";
- headache, tiredness; or
- nausea, vomiting, stomach pain, [constipation](#).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Fioricet with Codeine side effects](#) (more detail)

Related/similar drugs

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-the-counter painkiller and fever ...

Reviews & ratings

6.1 / 10

[13 Reviews](#)

[View more](#)

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

5.8 / 10

[30 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight

management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Celebrex

Celebrex is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation. Learn about ...

Reviews & ratings

7.0 / 10

[226 Reviews](#)

[View more](#)

Botox

Botox is used to treat chronic migraines, excessive sweating, bladder conditions, eye muscle ...

Reviews & ratings

5.7 / 10

[473 Reviews](#)

[View more](#)

Celecoxib

Celecoxib is a nonsteroidal anti-inflammatory drug (NSAID) used to treat pain or inflammation ...

Reviews & ratings

7.0 / 10

[280 Reviews](#)

[View more](#)

Diclofenac

Diclofenac is a nonsteroidal anti-inflammatory drug used to treat pain and inflammation associated ...

Reviews & ratings

7.3 / 10

7.3 / 10

[695 Reviews](#)

[View more](#)

Propranolol

Propranolol is a beta-blocker that is used to treat tremors, chest pain, high blood pressure, heart ...

Reviews & ratings

8.2 / 10

[1,376 Reviews](#)

[View more](#)

Naproxen

Naproxen is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

[683 Reviews](#)

[View more](#)

Amitriptyline

Amitriptyline is a tricyclic antidepressant used to treat depression and, off-label, conditions ...

Reviews & ratings

7.5 / 10

[1,803 Reviews](#)

[View more](#)

Warnings

MISUSE OF OPIOID MEDICINE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medication in a place where others cannot get to it.

Do not give this medicine to anyone younger than 12 years old, or anyone under 18 who recently had surgery to remove the tonsils or adenoids.

Taking opioid medicine during pregnancy may cause life-threatening withdrawal symptoms in the newborn.

Fatal side effects can occur if you use opioid medicine with alcohol, or with other drugs that cause drowsiness or slow your breathing.

Before taking this medicine

You should not use this medicine if you are allergic to acetaminophen, butalbital, caffeine, or codeine, or if you have:

- severe asthma or breathing problems;
- a blockage in your stomach or intestines;
- porphyria; or
- if you have taken an MAO inhibitor in the past 14 days, such as [isocarboxazid](#), linezolid, [methylene blue](#) injection, phenelzine, rasagiline, selegiline, or [tranylcypromine](#).

This medicine is **not approved** for use by anyone younger than 12 years old.

Do not give this medicine to anyone younger than 18 years old who recently had surgery to remove the tonsils or adenoids.

Tell your doctor if you have ever had:

- breathing problems, [sleep apnea](#) (breathing stops during sleep);
- a [head injury](#), or seizures;
- liver or kidney disease;
- urination problems;
- problems with your gallbladder, thyroid, or adrenal gland; or
- alcoholism or [drug addiction](#).

If you use opioid medicine while you are pregnant, your baby could become dependent on the drug. This can cause life-threatening withdrawal symptoms in the baby after it is born. Babies born dependent on opioids may need medical treatment for several weeks.

Do not breastfeed. Codeine can pass into breast milk and may cause drowsiness, breathing problems, or death in a nursing baby.

How should I take Fioricet with Codeine?

Follow the directions on your prescription label and read all medication guides. **Never use this medicine in larger amounts, or for longer than prescribed.** Tell your doctor if you feel an increased urge to take more of this medicine.

Never share opioid medicine with another person, especially someone with a history of drug abuse or addiction. MISUSE CAN CAUSE ADDICTION, OVERDOSE, OR DEATH. Keep the medication in a place where others cannot get to it. Selling or giving away opioid medicine is against the law.

Do not stop using this medicine suddenly after long-term use, or you could have unpleasant withdrawal symptoms.

Ask your doctor how to safely stop using the medicine.

Store at room temperature away from moisture and heat. Keep track of your medicine. Keep track of your medicine. You should be aware if anyone is using it improperly or without a prescription.

Do not keep leftover opioid medication. **Just one dose can cause death in someone using this medicine accidentally or improperly.** Ask your pharmacist where to locate a drug take-back disposal program. If there is no take-back program, mix the leftover medicine with cat litter or coffee grounds in a sealed plastic bag throw the bag in the trash.

 [Detailed Fioricet with Codeine dosage information](#)

What happens if I miss a dose?

Since Fioricet with Codeine is used for pain, you are not likely to miss a dose. Skip any missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An opioid overdose can be fatal, especially in a child or other person using the medicine without a prescription.** Overdose symptoms may include severe drowsiness, pinpoint pupils, slow breathing, or no breathing.

Your doctor may recommend you get naloxone (a medicine to reverse an [opioid overdose](#)) and keep it with you at all times. A person caring for you can give the naloxone if you stop breathing or don't wake up. Your caregiver must still get emergency medical help and may need to perform CPR (cardiopulmonary resuscitation) on you while waiting for help to arrive.

Anyone can buy naloxone from a pharmacy or local health department. Make sure any person caring for you knows where you keep naloxone and how to use it.

What should I avoid while taking Fioricet with Codeine?

Do not drink alcohol. Dangerous side effects or death could occur.

Avoid driving or hazardous activity until you know how this medicine will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Ask a doctor or pharmacist before using any other medicine that may contain acetaminophen (sometimes abbreviated as APAP). **Taking too much acetaminophen can lead to a fatal overdose.**

What other drugs will affect Fioricet with Codeine?

You may have breathing problems or withdrawal symptoms if you start or stop taking certain other medicines. Tell your doctor if you also use an antibiotic, antifungal medication, heart or blood pressure medication, seizure medication, or medicine to treat HIV or [hepatitis C](#).

Opioid medication can interact with many other drugs and cause dangerous side effects or death. Be sure your doctor knows if you also use:

- cold or allergy medicines, bronchodilator asthma/COPD medication, or a diuretic ("water pill");
- medicines for [motion sickness](#), [irritable bowel syndrome](#), or [overactive bladder](#);
- **other opioids**--opioid pain medicine or prescription cough medicine;
- **a sedative like Valium**--diazepam, [alprazolam](#), lorazepam, Xanax, Klonopin, Versed, and others;
- **drugs that make you sleepy or slow your breathing**--a sleeping pill, muscle relaxer, medicine to treat mood disorders or mental illness;
- **drugs that affect serotonin levels in your body**--a stimulant, or medicine for depression, [Parkinson's disease](#), migraine headaches, serious infections, or nausea and vomiting.

This list is not complete. Other drugs may affect Fioricet with Codeine, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Not all possible interactions are listed here.**

 [Fioricet with Codeine drug interactions](#) (more detail)

Does Fioricet with Codeine interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Fioricet with Codeine

+

Add

More about Fioricet with Codeine (acetaminophen / butalbital / caffeine / codeine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(20\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: narcotic analgesic combinations](#)
- [En español](#)

Patient resources

Other brands

[Phrenilin with Caffeine and Codeine](#)

Professional resources

- [Fioricet with Codeine prescribing information](#)

Other formulations

- [Fioricet](#)

Related treatment guides

- [Headache](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer


Copyright 1996-2025 Cerner Multum, Inc. Version: 19.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

 Risk data available

CSA Schedule*

3 Moderate abuse potential



User Reviews & Ratings

9.2 / 10

[20 Reviews](#)

Images

[Fioricet with Codeine 325 mg / 50 mg / 40 mg / 30 mg \(FIORICET CODEINE LOGO PROFILES\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.