

[Home](#)2. [Uloric](#)

# Uloric

**Generic name:** [febuxostat](#) [ *fe-BUX-oh-stat* ]**Drug class:** [Antihyperuricemic agents](#)Medically reviewed by [Philip Thornton, DipPharm](#). Last updated on Mar 21, 2025.[Uses](#) [Warnings](#) [Before taking](#) [Dosage](#) [Side effects](#) [Interactions](#)

## What is Uloric?

Uloric reduces the production of uric acid in your body. A build-up of uric acid can cause [gout symptoms](#).

Uloric is used to keep uric acid levels from getting too high in people with [gout](#).

Uloric is for use only in people who cannot take another medicine called allopurinol, or when allopurinol has stopped working.

## Warnings

**Taking Uloric increase your risk of serious or fatal heart problems.** Seek medical attention if you have chest pain, trouble breathing, fast heartbeats, sudden severe headache, numbness or weakness on one side of your body, or problems with vision or speech.

You should not use Uloric if you also use [azathioprine](#) or [mercaptopurine](#).

Before taking Uloric, tell your doctor if you have kidney disease, liver disease, diabetes, heart disease, cancer, Lesch-Nyhan syndrome, a history of heart attack or stroke, or if you have ever received an organ transplant.

You may have an increase in gout symptom flares when you first start taking Uloric. For best results, keep taking the medication as directed. Your doctor may prescribe other gout medications for you to use during the first 6 months of treatment.

To be sure Uloric is helping your condition, your blood may need to be tested often. Visit your doctor regularly.

## Before taking this medicine

You should not use Uloric if you are allergic to febuxostat, or if you also use:

- azathioprine; or
- mercaptopurine.

**Taking febuxostat may increase your risk of serious or fatal heart problems.** Talk with your doctor about the risks and benefits of this medicine.

To make sure Uloric is safe for you, tell your doctor if you have ever had:

- heart problems;
- a [heart attack](#) or stroke;
- liver or kidney disease
- [cancer](#);
- Lesch-Nyhan syndrome; or
- an [organ transplant](#) (kidney, liver, lung, heart).

It is not known whether febuxostat will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It may not be safe to breast-feed while using Uloric. Ask your doctor about any risk.

 [Uloric pregnancy and breastfeeding warnings](#) (more detail)

## How should I take Uloric?

Take Uloric exactly as prescribed by your doctor. Follow all directions on your prescription label. Your doctor may occasionally change your dose. Do not use this medicine in larger or smaller amounts or for longer than recommended.

You may take Uloric with or without food. Uloric may be taken with an antacid if needed.

You may have an increase in gout symptoms (flares) when you first start taking Uloric. Your doctor may prescribe other medications for you to use during the first 6 months of treatment.

**You should not stop using febuxostat suddenly.** Stopping suddenly may make your condition worse.

For best results, keep using your medications as directed. Call your doctor if your symptoms do not improve after 6 months, or if they get worse.

You will need frequent blood tests to check your liver function.

Store at room temperature away from moisture, heat, and light.

Read all patient information, medication guides, and instruction sheets provided to you. Ask your doctor or pharmacist if you have any questions.

## Dosing information

### Usual Adult Dose for Gout:

Initial dose: 40 mg orally once a day

-If serum uric acid level is greater than 6 mg/dL after 2 weeks, increase the dose to 80 mg orally once a day

Maintenance dose: 40 to 80 mg orally once a day

Comments:

-Use of this drug should occur after an acute gout flare subsides.

-Gout flares may occur due to increased urate mobilization from tissue deposits, especially during initiation; gout flare prophylaxis with a non-steroidal anti-inflammatory or [colchicine](#) is recommended and may be beneficial for up to 6 months.

-Uric acid levels may be checked as soon as 2 weeks after beginning this drug.

Use: Chronic management of symptomatic hyperuricemia in patients with gout

 [Detailed Uloric dosage information](#)

## What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose.

Do not take extra medicine to make up the missed dose.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

## What to avoid

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

## Uloric side effects

Get emergency medical help if you have **signs of an allergic reaction to Uloric** ([hives](#), difficult breathing, swelling in your face or throat) **or a severe skin reaction** ([fever](#), [sore throat](#), burning in your eyes, skin [pain](#), red or purple [skin rash](#) that spreads and causes blistering and peeling).

**Seek medical treatment if you have a serious drug reaction that can affect many parts of your body.** Symptoms may include: skin [rash](#), fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using Uloric.

Call your doctor at once if you have:

- a light-headed feeling, like you might pass out;
- **heart attack symptoms** - chest pain or pressure, pain spreading to your jaw or shoulder, fast heartbeats, shortness of breath;
- **liver problems** - stomach pain (upper right side), unusual tiredness, dark urine, [jaundice](#) (yellowing of the skin or eyes); or
- **signs of a stroke** - sudden numbness or weakness (especially on one side of the body), sudden severe [headache](#), slurred speech, problems with vision or balance.

Common Uloric side effects may include:

- gout flares, [joint pain](#);

- nausea;
- mild rash; or
- abnormal liver function tests.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Uloric side effects](#) (more detail)

## Related/similar drugs

### Krystexxa

Krystexxa (pegloticase) is an enzyme that metabolizes uric acid and is used to treat chronic gout ...

#### Reviews & ratings

8.6 / 10

[16 Reviews](#)

[View more](#)

#### FEATURED

### Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

#### Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

### Colchicine

Colchicine is a plant-based alkaloid that may be used to treat and prevent gout and some other ...

#### Reviews & ratings

7.3 / 10

[102 Reviews](#)

[View more](#)

### Triamcinolone

Triamcinolone is used to treat

Iriamcinoione is used to treat allergies, skin conditions, ulcerative colitis, and arthritis. Learn ...

#### Reviews & ratings

6.7 / 10

[226 Reviews](#)

[View more](#)

### Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

#### Reviews & ratings

5.9 / 10

[162 Reviews](#)

[View more](#)

### Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

#### Reviews & ratings

7.1 / 10

[484 Reviews](#)

[View more](#)

## What other drugs will affect Uloric?

Tell your doctor about all your current medicines and any you start or stop using, especially:

- [theophylline](#).

This list is not complete. Other drugs may interact with febuxostat, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#). Not all possible interactions are listed in this medication guide.

 [Uloric drug interactions](#) (more detail)

## Does Uloric interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Uloric

+

Enter a drug name

Add

## More about Uloric (febuxostat)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(46\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [FDA approval history](#)
- [Drug class: antihyperuricemic agents](#)
- [Breastfeeding](#)
- [En español](#)

## Professional resources

- [Uloric prescribing information](#)
- [Febuxostat \(AHFS Monograph\)](#)

## Related treatment guides

- [Gout](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Uloric only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 6.01.

## DRUG STATUS

### Availability

**Rx** Prescription only

### Pregnancy & Lactation

 Risk data available

### CSA Schedule\*

**N/A** Not a controlled drug



### Approval History

**10+ years** FDA approved 2009



## User Reviews & Ratings

7.7 / 10

[46 Reviews](#)

## Images

[Uloric 40 mg \(TAP 40\)](#)



[View larger images](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



### About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

### Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

### Support

[Help center](#)

[Sitemap](#)

[Contact us](#)





[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Copyright © 2000-2025 Drugs.com. All rights reserved.

