

Home

2. Mysoline

Mysoline ⊲ⁿ

Generic name: primidone [*PRIM-i-done*] **Drug class:** Barbiturate anticonvulsants



Medically reviewed by Drugs.com on Sep 7, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Mysoline?

Mysoline is an anticonvulsant that is used alone or with other medications to control seizures in adults and children.

Mysoline may also be used for purposes not listed in this medication guide.

Mysoline side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself.

Mysoline may cause serious side effects. Call your doctor at once if you have:

- extreme drowsiness, falling asleep suddenly, even after feeling alert; or
- signs of infection--fever, swollen glands, shortness of breath, chills, sore throat, body aches, unusual tiredness, loss of appetite, bruising or bleeding.

Common side effects of Mysoline may include:

- dizziness, drowsiness, vertigo (spinning sensation);
- problems with walking or moving;
- nausea, vomiting, loss of appetite;
- · feeling tired or irritable;
- · vision problems; or
- impotence, sexual problems.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Mysoline side effects (more detail)

Related/similar drugs

Depakote

Depakote is used to treat various types of seizure disorders. Learn about side effects ...

Reviews & ratings

6.3 / 10

329 Reviews

View more

Lyrica

Lyrica is used to control seizures, treat nerve pain and fibromyalgia. Learn about side effects ...

Reviews & ratings

6.4 / 10

925 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Keppra

Keppra is an anti-epileptic drug used to treat partial onset seizures in people with epilepsy ...

Reviews & ratings

View more

Diazepam

Diazepam is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.4 / 10

884 Reviews

View more

Lamictal

Lamictal is an anti-epileptic medication used treat seizures in adults and children over 2 years ...

Reviews & ratings

8.0 / 10

792 Reviews

View more

Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

Reviews & ratings

7.9 / 10

1,054 Reviews

View more

Pregabalin

Pregabalin may be used to treat certain types of pain and used in combination with other ...

Reviews & ratings

6.8 / 10

1,668 Reviews

Warnings

Some people have thoughts about suicide while taking seizure medicine. Tell your doctor right away if you have any sudden changes in mood or behavior, or thoughts about suicide.

Seizures may increase if you stop using Mysoline suddenly. Ask your doctor before stopping the medicine.

Before taking this medicine

You should not use Mysoline if you have:

- an allergy to phenobarbital; or
- porphyria (a genetic disorder that causes symptoms affecting the skin or nervous system).

Tell your doctor if you have ever had:

- · depression;
- a mood disorder; or
- · suicidal thoughts or actions.

Some people have thoughts about suicide while taking seizure medicine. Stay alert to changes in your mood or symptoms. Your family or caregivers should also watch for sudden changes in your behavior.

Do not start or stop seizure medication during pregnancy without your doctor's advice. Having a seizure during pregnancy could harm both mother and baby. Tell your doctor if you become pregnant.

If you take Mysoline during pregnancy, be sure to tell the doctor who delivers your baby. Both you and the baby may need medication to prevent bleeding during delivery and just after birth.

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of primidone on the baby.

Tell your doctor if you are breastfeeding. If you are breastfeeding, tell your doctor if you notice drowsiness in the nursing baby.

How should I take Mysoline?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You may not fully benefit from this medicine for up to several weeks. Your doctor will determine how long to treat you with this medicine.

You should not stop using Mysoline suddenly. Seizures may increase if you stop using this medicine. Ask your doctor before stopping the medicine.

You may need blood tests, and your next dose may be changed based on the results. Your doctor will need to check your progress on a regular basis.

You may be given other medications such as folic acid to help prevent serious side effects. Keep taking these medicines for as long as your doctor has prescribed.

Doses may be based on weight for children under 8 years old. Your child's dose may change if the child gains or loses weight.

Store tightly closed at room temperature, away from moisture and heat.

Detailed Mysoline dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Mysoline?

Avoid driving or hazardous activity until you know how Mysoline will affect you. Your reactions could be impaired.

Drinking alcohol with this medicine can cause side effects.

What other drugs will affect Mysoline?

Using Mysoline with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Other drugs may affect Mysoline, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell your doctor about all other medicines you use.

Mysoline drug interactions (more detail)

Does Mysoline interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Mysoline
+
Enter a drug name
Add

More about Mysoline (primidone)

- · Check interactions
- · Compare alternatives
- · Pricing & coupons
- Reviews (11)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Drug class: barbiturate anticonvulsants
- Breastfeeding
- En español

Patient resources

· Mysoline advanced reading

Professional resources

- Mysoline prescribing information
- Primidone (AHFS Monograph)

Related treatment guides

- Seizures
- Epilepsy

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 6.01.

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 1954

User Reviews & Ratings

8.7 / 10

11 Reviews

Images

Mysoline 250 mg (Logo MYSOLINE 250)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

 \times