



[Home](#)

2. Fluticasone

Fluticasone inhalation

Generic name: fluticasone inhalation [*floo-TIK-a-zone*]

Brand names: [Arnuity Ellipta](#), [Flovent Diskus](#), [Flovent HFA](#), [Flovent](#), [Flovent Rotadisk](#), ... [show all 9 brands](#)

Dosage forms: inhalation aerosol (110 mcg/inh; 220 mcg/inh; 44 mcg/inh), inhalation powder (100 mcg/inh; 113 mcg/inh; 232 mcg/inh; 250 mcg/inh; 50 mcg/inh; 55 mcg/inh; furoate 100 mcg/inh; furoate 200 mcg/inh; furoate 50 mcg/inh)

Drug class: [Inhaled corticosteroids](#)



[Medically reviewed](#) by Drugs.com on Mar 21, 2025. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

What is fluticasone inhalation?

[Fluticasone](#) inhalation is a steroid that is used to prevent asthma attacks in adults and children. Fluticasone inhalation is sometimes used together with steroid medicine taken by mouth.

Flovent brands of fluticasone inhalation may be used in children as young as 4 years old. **Arnuity Ellipta** should not be used in children younger than 5 years old, and **ArmonAir** should not be used in anyone younger than 12 years old.

Fluticasone may also be used for purposes not listed in this medication guide.

Fluticasone inhalation side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Fluticasone inhalation may cause serious side effects. Call your doctor at once if you have:

- weakness, tired feeling, nausea, vomiting, feeling like you might pass out;
- [wheezing](#), choking, or other breathing problems after using fluticasone inhalation;
- blurred vision, tunnel vision, eye pain, or seeing halos around lights;
- worsening of your asthma symptoms;
- **blood vessel inflammation**--fever, cough, stomach pain, [weight loss](#), skin rash, severe tingling, numbness, chest pain; or
- **liver problems**--upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).

Fluticasone can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine.

Common side effects of fluticasone inhalation may include:

- [cold symptoms](#) such as stuffy nose, sneezing, [sore throat](#), sinus pain;
- low fever, cough, wheezing, chest tightness;
- hoarseness or deepened voice;
- white patches or sores inside your mouth or on your lips;
- headache; or
- nausea, vomiting, [upset stomach](#).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Fluticasone side effects](#) (more detail)

Related/similar drugs

Xolair

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

Reviews & ratings

7.5 / 10

[229 Reviews](#)

[View more](#)

Eohilia

Eohilia (budesonide) suspension is used to treat eosinophilic esophagitis (EoE), to reduce symptoms ...

Reviews & ratings

[Add a review](#)

[View more](#)

FEATURED

Dupixent

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

Reviews & ratings

7.2 / 10

[499 Reviews](#)

[View more](#)

Symbicort

Symbicort (budesonide and formoterol) is used to prevent bronchospasm in people with asthma or ...

Reviews & ratings

6.6 / 10

[226 Reviews](#)

[View more](#)

Ventolin HFA

Ventolin HFA (albuterol) is used to treat or prevent breathing problems in patients who have asthma ...

Reviews & ratings

7.5 / 10

[18 Reviews](#)

[View more](#)

Breo Ellipta

Breo Ellipta (fluticasone and vilanterol) is used to prevent airflow obstruction or bronchospasm in ...

Reviews & ratings

5.8 / 10

[228 Reviews](#)

[View more](#)

Spiriva

Spiriva (tiotropium) is used to prevent bronchospasm in people with bronchitis, emphysema, or COPD ...

Reviews & ratings

5.9 / 10

[122 Reviews](#)

[View more](#)

[View more](#)

Xopenex

Xopenex is a short-acting bronchodilator used to treat or prevent bronchospasm in asthma and other ...

Reviews & ratings

9.0 / 10

[15 Reviews](#)

[View more](#)

Warnings

Fluticasone inhalation is not a rescue medicine for asthma attacks. Seek medical attention if your breathing problems get worse quickly, or if you think your asthma medications are not working as well.

You should not use the **inhalation powder** if you are allergic to milk proteins.

Before taking this medicine

You should not use fluticasone inhalation if you are allergic to fluticasone. Do not use the inhalation powder (**ArmonAir RespiClick**, **Arnuity Ellipta** or **Flovent Diskus**) if you are allergic to milk proteins.

Do not use fluticasone inhalation to treat an asthma attack that has already begun.

Tell your doctor if you have ever had:

- any type of [infection](#) (bacterial, viral, or fungal);
- [glaucoma](#) or cataracts;
- an infection caused by parasites (such as giardia, malaria, [leishmaniasis](#), hookworm, pinworm, [toxoplasmosis](#), and many others);
- [tuberculosis](#);
- herpes infection of the eyes;
- [liver disease](#);
- low bone mineral density; or
- a weak immune system.

Long-term use of steroids may lead to bone loss (osteoporosis), especially if you smoke, if you do not exercise, if you do not get enough vitamin D or calcium in your diet, or if you have a family history of [osteoporosis](#). Talk with your doctor about your risk.

Tell your doctor if you are pregnant. It is not known whether fluticasone will harm an unborn baby. However, having untreated or uncontrolled asthma during pregnancy may cause complications such as low birth weight, premature birth,

or eclampsia (dangerously [high blood pressure](#) that can lead to medical problems in both mother and baby). The benefit of treating asthma may outweigh any risks to the baby.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk.

Do not give this medicine to a child without medical advice.

How should I use fluticasone inhalation?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed. **Do not use the medicine more often than prescribed.**

Fluticasone inhalation is not a rescue medicine for asthma attacks. Use only fast-acting inhalation medicine for an attack. Seek medical attention if your breathing problems get worse quickly, or if you think your asthma medications are not working as well.

Flovent is a liquid form of fluticasone used with an inhaler device. This device creates a spray that you inhale through your mouth and into your lungs.

Flovent Diskus, ArmonAir, and Arnuity Ellipta are powder forms of fluticasone that come with a special inhaler device preloaded with blister packs containing measured doses of fluticasone. The device opens and loads a blister of fluticasone each time you use the inhaler.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions.

Do not allow a young child to use this medicine without help from an adult.

To prevent a yeast infection, rinse your mouth with water (do not swallow) after using this medicine. Pay close attention to your dental hygiene. Fluticasone can cause cavities or tooth discoloration.

Your dose needs may change if you have surgery, are ill, are under stress, or have recently had an asthma attack. **Do not change your medication dose or stop using fluticasone without your doctor's advice.**

You should not stop using fluticasone inhalation suddenly or your asthma could get worse.

If you stop taking any steroid medicine, you may need to stop gradually. Follow your doctor's instructions.

In case of emergency, wear or carry medical identification to let others know you may need an oral steroid in an emergency.

Your doctor will need to check your progress on a regular basis. Your vision may also need to be checked at regular intervals.

Call your doctor if your symptoms do not improve after 2 weeks of treatment, or if symptoms get worse. If you use a peak flow meter at home, call your doctor if your numbers are lower than normal.

Store at room temperature away from moisture, light, and extreme heat or cold.

Store **ArmonAir** in the foil pouch and use the medicine right away after opening. Discard the empty pouch in a place children and pets cannot reach. Store in a dry place.

Store the powder forms in the unopened foil pouch or tray until ready to use. Throw away any unused medicine when the inhalations counter on your inhaler device shows "0." **Get your prescription refilled before you run out of medicine completely.**

Store the **Flovent HFA** canister with the mouthpiece down. Keep the canister away from open flame or high heat, such as in a car on a hot day. The canister may explode if it gets too hot. Do not puncture or burn an empty inhaler canister.

 [Detailed Fluticasone dosage information](#)

What happens if I miss a dose?

Skip the missed dose and use your next dose at the regular time. **Do not** use two doses at one time.

If you use Arnuity Ellipta, do not use more than 1 inhalation per day.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

An overdose of fluticasone is not expected to produce life threatening symptoms. **Long term use of an inhaled steroid** can lead to glaucoma, cataracts, thinning skin, changes in body fat (especially in your face, neck, back, and waist), increased acne or facial hair, menstrual problems, [impotence](#), or loss of interest in sex.

What should I avoid while using fluticasone inhalation?

Avoid being near people who are sick or have infections. Call your doctor for preventive treatment if you are exposed to [chickenpox](#) or measles. These conditions can be serious or even fatal in people who are using steroid medicine.

What other drugs will affect fluticasone inhalation?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Tell your doctor about all your other medicines, especially:

- an [antibiotic](#);
- antifungal medicine;
- antiviral medicine to treat HIV or AIDS; or
- steroid medicine.

This list is not complete. Other drugs may affect fluticasone, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

 [Fluticasone drug interactions](#) (more detail)

Does fluticasone interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

fluticasone

+

Enter a drug name

Add

Popular FAQ

- What are the risks of using Flonase long-term?

▼
- Can asthma inhalers cause a sore throat?

▼
- How often can you use Flonase nasal spray?

▼
- How long does it take for Flonase to work?

▼
- What's the difference between Xhance and Flonase Allergy Relief?

▼

More FAQ

- [Can I use fluticasone nasal spray to relieve stuffy nose due to a cold?](#)

[View more FAQ...](#)

More about fluticasone

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(74\)](#)
- [Side effects](#)
- [Dosage information](#)
- [Patient tips](#)
- [During pregnancy](#)
- [Support group](#)
- [Drug class: inhaled corticosteroids](#)
- [En español](#)

Patient resources

- [Fluticasone HFA Aerosol Inhaler patient information](#)
- [Fluticasone Inhalation Powder](#)

Other brands

[Flovent](#), [Flovent HFA](#), [Arnuity Ellipta](#), [Flovent Diskus](#), ... +3 more

Professional resources

- [Fluticasone Propionate \(Systemic, Oral Inhalation\) monograph](#)
- [Fluticasone HFA Aerosol \(FDA\)](#)

Other brands

[Flovent](#), [Arnuity Ellipta](#), [Flovent Diskus](#), [ArmonAir Digihaler](#), [ArmonAir RespiClick](#)

Related treatment guides

- [Eosinophilic Esophagitis](#)
- [Asthma](#)
- [Bronchopulmonary Dysplasia](#)
- [Asthma, Maintenance](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 17.02.

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	Drug history at FDA
WADA Class	
	Anti-Doping Classification

User Reviews & Ratings

6.3 / 10

[74 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.