

Home

2. Lovastatin

Lovastatin do

Generic name: lovastatin [*LOE-va-sta-tin*] **Brand names:** Altoprev, Mevacor, Altocor

Dosage forms: oral tablet (10 mg; 20 mg; 40 mg), oral tablet, extended release (20 mg; 40 mg; 60 mg)

Drug class: Statins



Medically reviewed by Drugs.com on Sep 26, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is lovastatin?

Lovastatin is used together with diet to lower blood levels of "bad" cholesterol (low-density lipoprotein, or LDL), to increase levels of "good" cholesterol (high-density lipoprotein, or HDL), and to lower triglycerides (a type of fat in the blood).

Lovastatin is used to lower the risk of stroke, heart attack, and other heart complications in adults with diabetes, coronary heart disease, or other risk factors.

Lovastatin may also be used for purposes not listed in this medication guide.

Lovastatin side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Lovastatin can cause the breakdown of muscle tissue, which can lead to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine.

Also call your doctor at once if you have:

- muscle weakness in your hips, shoulders, neck, and back;
- trouble lifting your arms, trouble climbing or standing;
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath; or
- **liver problems**--loss of appetite, stomach pain (upper right side), tiredness, dark urine, jaundice (yellowing of the skin or eyes).

Common side effects of lovastatin may include:

· infections;

- · headache; or
- · accidental injury.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Lovastatin side effects (more detail)

Related/similar drugs

Lipitor

Lipitor is used to treat high cholesterol. Learn about side effects, interactions and indications.

Reviews & ratings

4.3 / 10

102 Reviews

View more

Crestor

Crestor (rosuvastatin) is used to treat high cholesterol and high triglycerides in the blood ...

Reviews & ratings

5.6 / 10

138 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Pravastatin

Pravastatin (Pravachol) is a statin medicine used to lower cholesterol and triglycerides in the ...

Reviews & ratings

4.5 / 10

78 Reviews

View more

Fenofibrate

Fenofibrate is used to treat high cholesterol and high triglyceride levels. Learn about side ...

Reviews & ratings

5.7 / 10

69 Reviews

View more

Simvastatin

Simvastatin (Zocor) is a cholesterollowering medication that blocks the production of cholesterol ...

Reviews & ratings

4.8 / 10

131 Reviews

View more

Ezetimibe

Ezetimibe is used alone or with other medication to treat high cholesterol. Learn about side ...

Reviews & ratings

3.9 / 10

172 Reviews

View more

Rosuvastatin

Rosuvastatin is a prescription medication used to treat high cholesterol and prevent heart disease

• •

Reviews & ratings

5.0 / 10

258 Reviews

View more

Atorvastatin

Atorvastatin is a cholesterol-lowering medication that blocks the production of cholesterol. Learn ...

Reviews & ratings

4.3 / 10

298 Reviews

View more

Zetia

Zetia (ezetimibe) is used to treat high cholesterol. Includes Zetia side effects, interactions and ...

Reviews & ratings

4.3 / 10

75 Reviews

View more

Warnings

You should not take lovastatin if you have active liver disease, or if you are pregnant or breastfeeding.

Tell your doctor about all your current medicines and any you start or stop using. **Many drugs can interact, and some drugs should not be used together.**

Stop taking lovastatin and tell your doctor at once if you become pregnant.

Before taking this medicine

You should not use lovastatin if you are allergic to it, or if:

- you are pregnant or breastfeeding; or
- you have active liver disease.

Many drugs can interact and cause dangerous effects. Some drugs should not be used together with lovastatin.

Your doctor may change your treatment plan if you also use:

- · nefazodone;
- an antibiotic--clarithromycin, erythromycin, telithromycin;
- antifungal medicine--itraconazole, ketoconazole, posaconazole, voriconazole; or
- antiviral medicine to treat HIV or hepatitis C--boceprevir, cobicistat, dasabuvir, elvitegravir, indinavir, lopinavir/ritonavir, nelfinavir, ombitasvir, paritaprevir saquinavir, telaprevir, tipranavir.

Tell your doctor if you have ever had:

- liver disease;
- · muscle pain or weakness;
- · kidney disease;
- a thyroid disorder; or
- · if you drink large amounts of alcohol.

Lovastatin can cause the breakdown of muscle tissue, which can lead to kidney failure. This happens more often in older adults or people who have kidney disease or poorly controlled hypothyroidism (underactive thyroid).

Do not use if you are pregnant. This medicine can harm an unborn baby. Use effective birth control to prevent pregnancy. Stop taking this medicine and tell your doctor at once if you become pregnant.

Do not breastfeed while using this medicine.

Lovastatin is not approved for use by anyone younger than 18 years old.

How should I take lovastatin?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Swallow the tablet whole and do not crush, chew, or break it.

It may take up to 4 weeks before your cholesterol levels improve, and you may need frequent blood tests. Even if you have no symptoms, tests can help your doctor determine if lovastatin is effective.

You may need to stop using lovastatin for a short time if you have surgery or a medical emergency. **Do not stop taking this medicine unless your doctor tells you to.**

Lovastatin is only part of a complete treatment program that may also include diet, exercise, and weight control. Follow your doctor's instructions very closely.

Store at room temperature away from moisture, heat, and light.

Detailed Lovastatin dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking lovastatin?

Avoid eating foods high in fat or cholesterol, or lovastatin will not be as effective.

Avoid drinking alcohol. It can raise triglyceride levels and may increase your risk of liver damage.

Grapefruit may interact with lovastatin and lead to unwanted side effects. Avoid the use of grapefruit products.

What other drugs will affect lovastatin?

Some drugs can increase your risk of serious muscle problems if you take them together with lovastatin. Your doctor may need to change your treatment plan if you use any of the following drugs:

- cyclosporine;
- · danazol;
- · gemfibrozil, fenofibric acid, fenofibrate; or
- medicines that contain niacin (Advicor, Niaspan, Niacor, Simcor, Slo-Niacin, and others).

This list is not complete and many other drugs may affect lovastatin. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Lovastatin drug interactions (more detail)

Does lovastatin interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

lov	astatin	
+		
Ent	ter a drug name	Add

Frequently asked questions

- · What are the side effects of statins?
- Do statins help treat COVID-19?

More about lovastatin

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (18)
- Drug images
- Latest FDA alerts (3)
- Side effects
- Dosage information
- Patient tips
- During pregnancy
- · Drug class: statins
- Breastfeeding
- En español

Patient resources

- · Lovastatin advanced reading
- Lovastatin Extended-Release Tablets
- Lovastatin Tablets

Other brands

Mevacor, Altoprev, Altocor

Professional resources

- · Lovastatin monograph
- Lovastatin (FDA)

Other brands

Altoprev

Related treatment guides

- High Cholesterol
- High Cholesterol, Familial Heterozygous
- Hyperlipoproteinemia
- Hyperlipoproteinemia Type IIa, Elevated LDL
- Hyperlipoproteinemia Type IIb, Elevated LDL VLDL

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 16.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ঝ্প Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

The Drug history at FDA

User Reviews & Ratings

6.2 / 10

18 Reviews

Related News

Two-Drug Combo of a Statin Plus Ezetimibe, Lowers Cholesterol Better Than Statins Alone

Images

Lovastatin 20 mg (TEVA 576)





Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

×