

Home

2. Tambocor

Tambocor ⊲

Generic name: flecainide [FLEK-a-nide] **Drug class:** Group I antiarrhythmics



Medically reviewed by Drugs.com on Aug 4, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Tambocor?

Tambocor is a Class IC anti-arrhythmic that is used in certain situations to prevent serious heart rhythm disorders.

Tambocor may also be used for purposes not listed in this medication guide.

Tambocor side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Tambocor may cause serious side effects. Call your doctor at once if you have:

- · fast or pounding heartbeats;
- fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out);
- slow heart rate, weak pulse, slow breathing (breathing may stop);
- · feeling short of breath;
- swelling, rapid weight gain;
- pale skin, easy bruising or bleeding, unusual weakness;
- · fever, flu-like symptoms; or
- jaundice (yellowing of the skin or eyes).

Common side effects of Tambocor may include:

- · dizziness;
- · vision problems;
- · trouble breathing;
- · headache;

- · nausea; or
- · feeling weak or tired.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Tambocor side effects (more detail)

Related/similar drugs

Xarelto

Xarelto (rivaroxaban) is a factor Xa inhibitor used to reduce the risk of blood clots and stroke in ...

Reviews & ratings

5.4 / 10

224 Reviews

View more

Cardizem

Cardizem is used to treat hypertension (high blood pressure), angina (chest pain), and heart rhythm ...

Reviews & ratings

7.3 / 10

12 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Phenylephrine

Oral and nasal phenylephrine are used as nasal decongestants to temporarily

unblock a nose or ...

Reviews & ratings

6.2 / 10

23 Reviews

View more

Sotalol

Sotalol is a beta-blocker with antiarrhythmic properties that affects the heart and circulation ...

Reviews & ratings

6.5 / 10

56 Reviews

View more

Bisoprolol

Bisoprolol belongs to a group of drugs called beta-blockers and is used to treat hypertension (high ...

Reviews & ratings

6.5 / 10

115 Reviews

View more

Digoxin

Digoxin is used to treat congestive heart failure and to slow the heart rate in patients with ...

Reviews & ratings

5.8 / 10

17 Reviews

View more

Warnings

You should not use Tambocor if you have a serious heart condition such as bundle branch block or AV block (without a pacemaker), or if your heart cannot pump blood properly.

You may receive your first dose in a hospital or clinic setting to quickly treat any serious side effects.

Before taking this medicine

You should not use Tambocor if you are allergic to it, or if:

- you have a serious heart condition such as bundle branch block or AV block (unless you have a pacemaker); or
- your heart cannot pump blood properly.

Tell your doctor if you have ever had:

- a heart attack;
- chronic atrial fibrillation, or "AFib";
- · congestive heart failure;
- a heart condition called "sick sinus syndrome";
- an electrolyte imbalance (such as low levels of potassium or magnesium in your blood); or
- liver disease.

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It may not be safe to breast-feed while using this medicine. Ask your doctor about any risk.

How should I take Tambocor?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

You may receive your first few doses in a hospital or clinic setting to quickly treat any serious side effects.

Your heart rate will be monitored using an electrocardiograph or ECG (sometimes called an EKG). This will help your doctor determine how long to treat you with Tambocor.

You may also need frequent blood tests to check your liver or kidney function.

Store at room temperature away from moisture, heat, and light.

Detailed Tambocor dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include nausea, vomiting, slow heart rate, fainting, or seizure (convulsions).

What should I avoid while taking Tambocor?

Follow your doctor's instructions about any restrictions on food, beverages, or activity.

What other drugs will affect Tambocor?

Tell your doctor about all your other medicines, especially:

- · digoxin;
- a diuretic or "water pill";
- a beta-blocker (atenolol, metoprolol, propranolol, sotalol, and others);
- other heart medications such as amiodarone, diltiazem, disopyramide, nifedipine, quinidine, or verapamil; or
- · seizure medication.

This list is not complete. Other drugs may affect Tambocor, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

1 Tambocor drug interactions (more detail)

Does Tambocor interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Tambocor
+
Enter a drug name
Add

More about Tambocor (flecainide)

- · Check interactions
- Compare alternatives
- Reviews (21)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: group I antiarrhythmics
- Breastfeeding

Patient resources

• Tambocor advanced reading

Professional resources

- Tambocor prescribing information
- Flecainide Acetate (AHFS Monograph)

Related treatment guides

- Atrial Fibrillation
- Atrial Flutter
- Paroxysmal Supraventricular Tachycardia
- Ventricular Tachycardia
- Supraventricular Tachycardia
- Wolff-Parkinson-White Syndrome

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 6.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ড Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

10+ years FDA approved 1985

User Reviews & Ratings

7.3 / 10

Images

Tambocor 100 mg (3M TR 100)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







