

Home

- 2. Cyclessa
- Consumer

Cyclessa

Generic name: Ethinyl Estradiol and Desogestrel [ETH-in-il-es-tra-DYE-ole-& des-oh-JES-trel]

Brand names: Apri, Azurette, Bekyree, Caziant, Cyclessa, ... show all 22 brands

Drug class: Contraceptives

Medically reviewed by Drugs.com. Last updated on Feb 26, 2024.

Uses Before taking Warnings Dosage Side effects Overdose

Warning

• Smoking cigarettes while using Cyclessa (ethinyl estradiol and desogestrel) raises the chance of very bad heart and blood-related side effects. This chance is raised with age (mainly in women older than 35 years of age). It is also raised with the number of cigarettes smoked. It is strongly advised not to smoke. Do not use Cyclessa (ethinyl estradiol and desogestrel) if you smoke and are older than 35 years of age.

Uses of Cyclessa:

- It is used to prevent pregnancy.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take Cyclessa?

- If you are allergic to Cyclessa (ethinyl estradiol and desogestrel); any part of Cyclessa (ethinyl estradiol and desogestrel); or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have had any of these health problems: Blood clots, blood clotting problem, breast cancer or other cancer
 where hormones make it grow, diseased blood vessels in the brain or heart, disease of a heart valve with problems,
 heart disease, chest pain caused by angina, heart attack, stroke, high blood pressure, liver disease, liver tumor, very
 bad headache or migraine, or diabetes that affects blood flow.
- If you have had any of these health problems: Endometrial cancer, cancer of the cervix or vagina, or vaginal bleeding where the cause is not known.
- If you are taking ombitasvir, paritaprevir, and ritonavir (with or without dasabuvir).
- If you turned yellow during pregnancy or with estrogen-based or hormone contraceptive use.
- If you are pregnant or may be pregnant. Do not take Cyclessa (ethinyl estradiol and desogestrel) if you are pregnant.

This is not a list of all drugs or health problems that interact with Cyclessa (ethinyl estradiol and desogestrel).

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take Cyclessa (ethinyl estradiol and desogestrel) with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take Cyclessa?

- Tell all of your health care providers that you take Cyclessa (ethinyl estradiol and desogestrel). This includes your
 doctors, nurses, pharmacists, and dentists. This medicine may need to be stopped before certain types of surgery
 as your doctor has told you. If Cyclessa (ethinyl estradiol and desogestrel) is stopped, your doctor will tell you when
 to start taking Cyclessa (ethinyl estradiol and desogestrel) again after your surgery or procedure.
- This medicine may raise the chance of blood clots, a stroke, or a heart attack. Talk with the doctor.
- Talk with your doctor if you will need to be still for long periods of time like long trips, bedrest after surgery, or illness. Not moving for long periods may raise your chance of blood clots.
- If you have high blood sugar (diabetes), talk with your doctor. This medicine may raise blood sugar.
- Check your blood sugar as you have been told by your doctor.
- Tell your doctor if you have signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- High blood pressure has happened with drugs like this one. Have your blood pressure checked as you have been told by your doctor.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- Be sure to have regular breast exams and gynecology check-ups. You will also need to do breast self-exams as you
 have been told.
- If you drink grapefruit juice or eat grapefruit often, talk with your doctor.
- This medicine may affect certain lab tests. Tell all of your health care providers and lab workers that you take Cyclessa (ethinyl estradiol and desogestrel).
- This medicine may cause high cholesterol and triglyceride levels. Talk with the doctor.
- This medicine may cause dark patches of skin on your face. Avoid sun, sunlamps, and tanning beds. Use sunscreen and wear clothing and eyewear that protects you from the sun.
- Certain drugs, herbal products, or health problems could cause Cyclessa (ethinyl estradiol and desogestrel) to not work as well. Be sure your doctor knows about all of your drugs and health problems.
- This medicine does not stop the spread of diseases like HIV or hepatitis that are passed through having sex. Do not have any kind of sex without using a latex or polyurethane condom. If you have questions, talk with your doctor.
- If you are allergic to tartrazine, talk with your doctor. Some products have tartrazine.
- Do not use in children who have not had their first menstrual period.
- If you have any signs of pregnancy or if you have a positive pregnancy test, call your doctor right away.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

Use Cyclessa (ethinyl estradiol and desogestrel) as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take Cyclessa (ethinyl estradiol and desogestrel) at the same time of day.
- Take with or without food. Take with food if it causes an upset stomach.
- After starting Cyclessa (ethinyl estradiol and desogestrel), you may need to use a non-hormone type of birth control
 like condoms to prevent pregnancy for some time. Follow what your doctor has told you to do about using a non-hormone type of birth control.
- If you also take colesevelam, take it at least 4 hours before or after you take Cyclessa (ethinyl estradiol and desogestrel).
- Do not skip doses, even if you do not have sex very often.
- If you throw up or have diarrhea, Cyclessa (ethinyl estradiol and desogestrel) may not work as well to prevent pregnancy. Use an extra form of birth control, like condoms, until you check with your doctor.
- If you miss 2 periods in a row, take a pregnancy test before starting a new cycle.

What do I do if I miss a dose?

- If a dose is missed, check the package insert or call the doctor to find out what to do. If using Cyclessa (ethinyl
 estradiol and desogestrel) to prevent pregnancy, another form of birth control may need to be used for some time to
 prevent pregnancy.
- Detailed Cyclessa dosage information

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of gallbladder problems like pain in the upper right belly area, right shoulder area, or between the shoulder blades; yellow skin or eyes; fever with chills; bloating; or very upset stomach or throwing up.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Depression or other mood changes.
- · Trouble sleeping.
- · Feeling very tired or weak.
- Eyesight changes or loss, bulging eyes, or change in how contact lenses feel.

- Not able to pass urine or change in how much urine is passed.
- A lump in the breast, breast pain or soreness, or nipple discharge.
- · Vaginal itching or discharge.
- Spotting or vaginal bleeding that is very bad or does not go away.
- This medicine may cause you to swell or keep fluid in your body. Tell your doctor if you have swelling, weight gain, or trouble breathing.
- Call your doctor right away if you have signs of a blood clot like chest pain or pressure; coughing up blood;
 shortness of breath; swelling, warmth, numbness, change of color, or pain in a leg or arm; or trouble speaking or swallowing.
- Cyclessa side effects (more detail)

Related/similar drugs

Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

Reviews & ratings

6.1 / 10

2,624 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Depo Provera

Depo-Provera is used as contraception to prevent pregnancy. It is also used to reduce pain cause by

Reviews & ratings

4.2 / 10

1,967 Reviews

View more

Nexplanon

Nexplanon is a long-term, reversible birth control implant that lasts for three years. Learn about ...

Reviews & ratings

5.0 / 10

4,864 Reviews

View more

Yaz

Yaz (drospirenone and ethinyl estradiol) is used as a contraceptive to prevent pregnancy. Includes ...

Reviews & ratings

5.9 / 10

997 Reviews

View more

Slynd

Slynd (drospirenone) is a birth control pill used for the prevention of pregnancy. Includes Slynd ...

Reviews & ratings

5.8 / 10

386 Reviews

View more

What are some other side effects of Cyclessa?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- · Feeling more or less hungry.
- · Dizziness or headache.
- · Weight gain or loss.

- · Upset stomach or throwing up.
- · Stomach pain or cramps.
- Bloating.
- · Period (menstrual) changes. These include spotting or bleeding between cycles.
- Enlarged breasts.
- · Tender breasts.
- Lowered interest in sex.
- Hair loss.
- · Feeling nervous and excitable.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

If OVERDOSE is suspected:

If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

How do I store and/or throw out Cyclessa?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so.
 Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Consumer Information Use and Disclaimer

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions
 about Cyclessa (ethinyl estradiol and desogestrel), please talk with your doctor, nurse, pharmacist, or other health
 care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

More about Cyclessa (desogestrel / ethinyl estradiol)

- Check interactions
- Compare alternatives
- Reviews (6)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Drug class: contraceptives

Patient resources

· Cyclessa advanced reading

Other brands

Apri, Enskyce, Kariva, Isibloom, ... +17 more

Professional resources

• Cyclessa prescribing information

Other brands

Apri, Enskyce, Kariva, Isibloom, ... +18 more

Related treatment guides

• Birth Control

Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ড Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History ☐ Drug history at FDA

User Reviews & Ratings

6 Reviews

Images

Cyclessa inert (KH 2 ORGANON)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









