

Home

Fleet Bisacodyl

Fleet Bisacodyl Enema

Generic name: Bisacodyl Enema [bis-AK-oh-dil]

Drug class: Laxatives

Medically reviewed by Drugs.com. Last updated on Dec 28, 2023.

Uses Before taking Warnings Dosage Side effects Overdose FAQ

Uses of Fleet Bisacodyl Enema:

- It is used to treat constipation.
- It is used as a laxative to clean out the colon before an exam.

What do I need to tell my doctor BEFORE I take Fleet Bisacodyl Enema?

- If you have an allergy to bisacodyl or any other part of Fleet Bisacodyl Enema (bisacodyl enema).
- If you are allergic to Fleet Bisacodyl Enema (bisacodyl enema); any part of Fleet Bisacodyl Enema (bisacodyl enema); or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: Bowel block, belly pain, upset stomach, rectal bleeding, throwing up, or change in bowel habits lasting longer than 2 weeks.

This is not a list of all drugs or health problems that interact with Fleet Bisacodyl Enema (bisacodyl enema).

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take Fleet Bisacodyl Enema (bisacodyl enema) with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take Fleet Bisacodyl Enema?

- Tell all of your health care providers that you take Fleet Bisacodyl Enema (bisacodyl enema). This includes your doctors, nurses, pharmacists, and dentists.
- Do not use Fleet Bisacodyl Enema (bisacodyl enema) for more than 1 week unless told to do so by your doctor.
- Do not use other laxatives or stool softeners unless told to do so by the doctor.
- If you have rectal bleeding or you do not have a bowel movement after using Fleet Bisacodyl Enema (bisacodyl enema), talk with your doctor.
- Do not give to a child younger than 12 years of age.

- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using Fleet Bisacodyl Enema (bisacodyl enema) while you are pregnant.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

How is this medicine (Fleet Bisacodyl Enema) best taken?

Use Fleet Bisacodyl Enema (bisacodyl enema) as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Wash your hands before and after use.
- Use enema rectally.
- Shake well before use.
- Put enema tip into the rectum with gentle pressure. Do not force.
- Squeeze the bottle until almost all the liquid is gone.

What do I do if I miss a dose?

- If you take Fleet Bisacodyl Enema (bisacodyl enema) on a regular basis, take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- Many times Fleet Bisacodyl Enema (bisacodyl enema) is taken on an as needed basis. Do not take more often than told by the doctor.
- Detailed Fleet Bisacodyl dosage information

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever;
 wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Fleet Bisacodyl side effects (more detail)

Related/similar drugs

MiraLAX

MiraLAX (polyethylene glycol) is used as a laxative to treat occasional constipation or irregular ...

_ . . .

Reviews & ratings

7.2 / 10

262 Reviews

View more

Colace

Colace makes bowel movements softer and is used to treat or prevent constipation. Includes Colace ...

Reviews & ratings

6.3 / 10

18 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Linzess

Linzess (linaclotide) is used to treat chronic idiopathic constipation, irritable bowel syndrome ...

Reviews & ratings

6.7 / 10

884 Reviews

View more

Dulcolax

Dulcolax (bisacodyl) is a laxative used to treat constipation. Includes Dulcolax side effects ...

Reviews & ratings

5.3 / 10

1,158 Reviews

Sutab

Sutab (sodium sulfate, magnesium sulfate, and potassium chloride) is used for colonoscopy ...

Reviews & ratings

6.3 / 10

126 Reviews

View more

What are some other side effects of Fleet Bisacodyl Enema?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- · Belly pain.
- · Stomach cramps.
- · Burning.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

If OVERDOSE is suspected:

If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

How do I store and/or throw out Fleet Bisacodyl Enema?

- Store at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so.
 Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Consumer Information Use and Disclaimer

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions
 about Fleet Bisacodyl Enema (bisacodyl enema), please talk with your doctor, nurse, pharmacist, or other health
 care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Frequently asked questions

- Which drugs cause opioid-induced constipation?
- Dulcolax How long does it take for bisacodyl to clear your system?

More about Fleet Bisacodyl (bisacodyl)

- Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (3)
- Side effects
- · Dosage information
- During pregnancy
- Drug class: laxatives
- Breastfeeding
- En español

Patient resources

Other brands

Dulcolax, Correctol

Professional resources

- · Bisacodyl monograph
- Bisacodyl (FDA)

Other brands

Dulcolax

Related treatment guides

Constipation

Further information

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

DRUG STATUS

Availability

OTC Over the counter

Pregnancy & Lactation

প্ Risk data available

CSA Schedule*

N/A Not a controlled drug

User Reviews & Ratings

3 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.