

Home

Azor

Azor ⊲»

Generic name: amlodipine and olmesartan [am-LOE-de-peen-ol-me-SAR-tan]

Drug class: Angiotensin II inhibitors with calcium channel blockers

Medically reviewed by Sanjai Sinha, MD. Last updated on Apr 15, 2024.

Uses Warnings Before taking Dosage Side effects Interactions

What is Azor?

Azor contains a combination of amlodipine and olmesartan.

Amlodipine is a calcium channel blocker. It works by relaxing the muscles of your heart and blood vessels.

Olmesartan is an angiotensin II receptor antagonist (sometimes called an ARB). Olmesartan keeps blood vessels from narrowing, which lowers blood pressure and improves blood flow.

Azor is used to treat high blood pressure (hypertension). Lowering blood pressure may lower your risk of a stroke or heart attack.

Warnings

Do not use Azor if you are pregnant. If you become pregnant, stop taking this medicine and tell your doctor right away.

If you have diabetes, do not use Azor together with any medication that contains aliskiren (such as Amturnide, Tekturna, Tekamlo).

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Get up slowly and steady yourself to prevent a fall.

Call your doctor at once if you have swelling in your hands or feet, rapid weight gain, pounding heartbeats or fluttering in your chest, urinating less than usual, jaundice (yellowing of the skin or eyes), chest pain or heavy feeling, pain spreading to the arm or shoulder, nausea, sweating, general ill feeling, or if you feel like you might pass out.

Before taking this medicine

You should not use Azor if you are allergic to amlodipine or olmesartan.

If you have diabetes, do not use Azor together with any medication that contains aliskiren (a blood pressure medicine).

You may also need to avoid taking amlodipine and olmesartan with aliskiren if you have kidney disease.

To make sure Azor is safe for you, tell your doctor if you have:

- · kidney disease;
- · liver disease;
- · coronary artery disease;
- angina (chest pain), congestive heart failure;
- high levels of potassium in your blood;
- if you are on a low-salt diet;
- if you are 75 years or older; or
- if you recently had a heart attack.

Do not use if you are pregnant. If you become pregnant, stop taking this medicine and tell your doctor right away. Olmesartan can cause injury or death to the unborn baby if you take the medicine during your second or third trimester.

You should not breastfeed while using this medicine.

Azor is not approved for use by anyone younger than 18 years old.

Azor pregnancy and breastfeeding warnings (more detail)

How should I take Azor?

Take Azor exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose.

You may take Azor with or without food.

Your blood pressure will need to be checked often. Your kidney function may also need to be checked.

Certain conditions can cause your blood pressure to get too low. This includes vomiting, diarrhea, or heavy sweating. Call your doctor if you are sick with vomiting or diarrhea.

If you need surgery, tell the surgeon ahead of time that you are using Azor.

Keep using Azor as directed, even if you feel well. High blood pressure often has no symptoms. You may need to use blood pressure medication for the rest of your life.

Store at room temperature away from moisture and heat.

Dosing information

Usual Adult Dose for Hypertension:

Initial dose: Amlodipine 5 mg-Olmesartan 20 mg orally once a day; may increase dose after 1 to 2 weeks of therapy.

Maximum dose: Amlodipine 10 mg-Olmesartan 40 mg orally once a day

Uses:

- -Substitution therapy for patients maintained on the individual components separately or for patients not adequately controlled with one of the components or a component within the same antihypertensive class.
- -Initial therapy in patients who are likely to need more than 1 antihypertensive agent to control blood pressure.



What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What to avoid

If you also take colesevelam, avoid taking it for at least 4 hours after you have taken Azor.

Avoid driving or hazardous activity until you know how this medicine will affect you. Your reactions could be impaired. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

Do not use potassium supplements or salt substitutes, unless your doctor has told you to.

Azor side effects

Get emergency medical help if you have **signs of an allergic reaction to Azor:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

When you first start taking this medicine, you may have new or worsening chest pain (angina), or you could have a heart attack. Seek emergency medical attention or call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating.

Call your doctor at once if you have:

- new or worsening chest pain;
- · swelling in your hands or feet, rapid weight gain;
- severe or ongoing diarrhea with weight loss;
- · pounding heartbeats or fluttering in your chest;
- a light-headed feeling, like you might pass out; or
- high potassium level nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement.

Common Azor side effects include:

- · swelling;
- dizziness;
- flushing (warmth, redness, or tingly feeling); or
- pounding heartbeats.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Azor side effects (more detail)

Related/similar drugs

Vasotec

Vasotec (enalapril) is used to treat high blood pressure, heart failure and kidney problems caused ...

Reviews & ratings

4 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Lisinopril

Lisinopril is an ACE inhibitor. It is used to treat high blood pressure, congestive heart failure ...

Reviews & ratings

5.0 / 10

626 Reviews

View more

Atenolol

Atenolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

Reviews & ratings

7.2 / 10

276 Reviews

View more

Valsartan

Valsartan is an angiotensin II receptor blocker (ARB) that may be used to treat high blood pressure ...

Reviews & ratings

5.7 / 10

152 Reviews

View more

Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

Reviews & ratings

5.2 / 10

185 Reviews

View more

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

6.4 / 10

56 Reviews

View more

Furosemide

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

Reviews & ratings

6.8 / 10

121 Reviews

View more

Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

Reviews & ratings

4.5 / 10

578 Reviews

View more

What other drugs will affect Azor?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

If you also take colesevelam, take it 4 hours after you take Azor.

Tell your doctor about all your other medicines, especially:

- · cyclosporine;
- lithium;
- · simvastatin;
- tacrolimus; or
- NSAIDs (nonsteroidal anti-inflammatory drugs) aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others.

This list is not complete. Other drugs may interact with amlodipine and olmesartan, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Azor drug interactions (more detail)

Does Azor interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Azor
+
Enter a drug name

Aug Aug India

References

1. Azor Product Label

More about Azor (amlodipine / olmesartan)

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (59)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- FDA approval history
- Drug class: angiotensin II inhibitors with calcium channel blockers
- En español

Professional resources

· Azor prescribing information

Related treatment guides

• High Blood Pressure

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Azor only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

8.5 / 10

59 Reviews

Images

Azor 5 mg / 20 mg (C73)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













<u>Subscribe to our newsletter</u> for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

 \times