

[Home](#)

2. Vandazole

Vandazole

Generic name: [metronidazole vaginal](#) [*MET-roe-NYE-da-zole*]**Brand names:** [Nuversa](#), Vandazole**Drug classes:** [Topical anti-rosacea agents](#), [Vaginal anti-infectives](#)[Medically reviewed](#) by Drugs.com on Feb 28, 2025. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is Vandazole?

Vandazole (for use in the vagina) is an [antibiotic](#) that is used to treat vaginal infections caused by bacteria.

Do not use [metronidazole](#) to treat any condition that has not been checked by your doctor.

Vandazole may also be used for purposes not listed in this medication guide.

Vandazole side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, itching, warmth or tingling; fever, [joint pain](#); [dry mouth](#), dry vagina; stuffy nose, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, [sore throat](#), burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Rare but serious side effects may include:

- seizure (convulsions); or
- numbness, burning, pain, or tingly feeling in your hands or feet.

Call your doctor at once if you have new or worsening vaginal symptoms.

Common side effects of Vandazole may include:

- vaginal itching or discharge;
- menstrual pain, heavy bleeding;
- breast pain;
- headache;
- rash, itching;
- stomach pain, nausea, diarrhea; or

- [cold symptoms](#) such as stuffy nose, sneezing, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Vandazole side effects](#) (more detail)

Related/similar drugs

Flagyl

Flagyl is used to treat bacterial infections of the vagina, stomach, skin and joints. Learn about ...

Reviews & ratings

6.1 / 10

[393 Reviews](#)

[View more](#)

Cleocin

Cleocin (clindamycin) is used to treat serious bacterial infections. Includes Cleocin side effects ...

Reviews & ratings

5.7 / 10

[17 Reviews](#)

[View more](#)

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

[685 Reviews](#)

[View more](#)

Clindamycin

Clindamycin (Cleocin) is used to treat serious infections caused by bacteria. Includes clindamycin ...

Reviews & ratings

5.7 / 10

[806 Reviews](#)

[View more](#)

Metronidazole

Metronidazole is an antibiotic used to fight bacteria in your body. Learn about side effects ...

Reviews & ratings

6.0 / 10

[2,431 Reviews](#)

[View more](#)

Warnings

You should not use metronidazole if you have taken [disulfiram](#) ([Antabuse](#)) within the past 2 weeks.

Do not drink alcohol or consume foods or medicines that contain propylene glycol while you are using metronidazole and for at least 3 days after you stop using it.

Before taking this medicine

You should not use Vandazole if you are allergic to metronidazole, [secnidazole](#), or [tinidazole](#), or if:

- you are allergic to parabens or polyethylene glycol;
- you recently drank alcohol; or
- you took disulfiram (Antabuse) within the past 14 days.

Tell your doctor if you have ever had:

- nerve problems;
- a seizure; or
- [peripheral vascular disease](#) such as [Raynaud's syndrome](#).

It is not known whether this medicine will harm an unborn baby. Tell your doctor if you are pregnant.

You should not breastfeed within 48 hours after using metronidazole. If you use a breast pump during this time, throw out any milk you collect. Do not feed it to your baby.

Vandazole is not approved for use in children younger than 12 years old, or in girls who have not begun having menstrual periods.

How should I use Vandazole?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Use this medicine at bedtime.

Vandazole is often given as a single dose. However, you might need to use the medicine for 5 nights in a row. **Follow your doctor's dosing instructions very carefully.**

Do not take Vandazole by mouth. It is for use only in the vagina.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions.

Gently wash your vaginal area before applying the medicine. Wash your hands before and after handling the medicine tube or vaginal applicator.

Insert the medicine directly into your vagina using the applicator provided. **Each disposable applicator is for one use only.**

You may need to use a sanitary napkin during treatment, but **do not use a tampon**. Do not use a douche or other vaginal product while using Vandazole.

Use this medicine for the full prescribed length of time, even if your symptoms quickly improve. Skipping doses can increase your risk of [infection](#) that is resistant to medication.

Call your doctor if your symptoms do not improve, or if they get worse.

Do not share this medicine with another person, even if they have the same symptoms you have.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Vandazole.

Vandazole will not treat a vaginal yeast infection. You may even develop a new [vaginal yeast infection](#), which may need to be treated with antifungal medication. Tell your doctor if you have symptoms such as itching or discharge during or after treatment with this medicine.

Store at room temperature away from moisture or heat. Do not refrigerate or freeze.

 [Detailed Vandazole dosage information](#)

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

An overdose of Vandazole is not expected to be dangerous. Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

What should I avoid while using metronidazole vaginal?

While you are using Vandazole and for 3 days after your last dose: Avoid drinking alcohol or consuming foods or other products that contain alcohol or propylene glycol. Drinking alcohol may cause unpleasant effects such as headaches, nausea, vomiting, stomach cramps, and warmth or tingling under your skin.

Avoid wearing tight-fitting, synthetic clothing such as nylon underwear or panty hose that does not allow air circulation. Wear loose-fitting clothing made of cotton and other natural fibers until your infection is healed.

Getting this medicine in your eyes can cause burning or irritation. If contact does occur, rinse with water and call your doctor.

Avoid using other vaginal medications unless your doctor tells you to. Avoid having sex until your vaginal infection has been completely treated.

What other drugs will affect Vandazole?

Tell your doctor about all your other medicines, especially:

- lithium; or
- a blood thinner ([warfarin](#), [Coumadin](#), [Jantoven](#)).

This list is not complete. Other drugs may affect Vandazole, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

 [Vandazole drug interactions](#) (more detail)

Does Vandazole interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Vandazole

+

Enter a drug name

Add

More about Vandazole (metronidazole topical)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(3\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)

- [Drug class: topical anti-rosacea agents](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

- [Vandazole advanced reading](#)

Other brands

[MetroGel](#), [MetroGel-Vaginal](#), [MetroCream](#), [Rozex](#), ... +5 more

Professional resources

- [Vandazole prescribing information](#)
- [MetroNIDAZOLE \(Topical\) \(AHFS Monograph\)](#)

Other brands

[MetroGel](#), [MetroGel-Vaginal](#), [MetroCream](#), [Noritate](#), ... +3 more

Related treatment guides

- [Bacterial Vaginosis](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

Antibiotics 101

[List of Common Antibiotics & Types](#)

[Antibiotics and Alcohol](#)

[Probiotics and Antibiotics](#)

[Antibiotics for Strep Throat](#)

[Antibiotics for a Tooth Infection](#)


[Antibiotics for UTI's](#)

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation


 Risk data available

CSA Schedule*

N/A Not a controlled drug



Approval History

 Drug history at FDA



User Reviews & Ratings

[3 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)


Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



ADVERTISEMENT

