

[Home](#)2. [Motrin Infant Drops](#)

Motrin Infant Drops

Generic name: [ibuprofen](#) [*EYE-bue-PROE-fen*]**Brand names:** [Advil](#), [Advil Migraine](#), Children's Advil, Children's Ibuprofen Berry, [Children's Motrin](#), ... [show all 17 brands](#)**Drug class:** [Nonsteroidal anti-inflammatory drugs](#)[Medically reviewed](#) by Drugs.com on Sep 20, 2023. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

What is Motrin Infant Drops?

Motrin Infant Drops is a nonsteroidal anti-inflammatory drug (NSAID).

Motrin Infant Drops is used to reduce fever and treat pain or inflammation caused by many conditions such as headache, [toothache](#), [back pain](#), [arthritis](#), menstrual cramps, or minor injury. Motrin Infant Drops is used in adults and children who are at least 6 months old.

Motrin Infant Drops may also be used for purposes not listed in this medication guide.

Motrin Infant Drops side effects

Get emergency medical help if you have **signs of an allergic reaction** (hives, difficult breathing, swelling in your face or throat) **or a severe skin reaction** (fever, [sore throat](#), burning eyes, skin pain, red or purple skin rash with blistering and peeling).

Get emergency medical help if you have **signs of a heart attack or stroke**: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath.

Motrin Infant Drops may cause serious side effects. Stop using Motrin Infant Drops and call your doctor at once if you have:

- changes in your vision;
- [shortness of breath](#) (even with mild exertion);
- swelling or rapid weight gain;
- a skin rash, no matter how mild;
- **signs of stomach bleeding**--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds;
- **liver problems**--nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);

- **low red blood cells (anemia)**--pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating; or
- **kidney problems**--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath.

Common side effects of Motrin Infant Drops may include:

- nausea, vomiting, gas;
- bleeding; or
- dizziness, headache.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Motrin Infant Drops side effects](#) (more detail)

Related/similar drugs

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

5.8 / 10

[30 Reviews](#)

[View more](#)

Celebrex

Celebrex is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation. Learn about ...

Reviews & ratings

7.0 / 10

[226 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Advil

Advil (ibuprofen) reduces fever, pain and inflammation caused by headaches, toothaches, arthritis ...

Reviews & ratings

6.9 / 10

[41 Reviews](#)

[View more](#)

Acetaminophen

Acetaminophen is a widely used pain reliever and fever reducer for conditions like headaches ...

Reviews & ratings

6.1 / 10

[104 Reviews](#)

[View more](#)

Naproxen

Naproxen is a nonsteroidal anti-inflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

[683 Reviews](#)

[View more](#)

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-the-counter painkiller and fever ...

Reviews & ratings

6.1 / 10

[13 Reviews](#)

[View more](#)

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

[1,404 Reviews](#)

[View more](#)

Warnings

Motrin Infant Drops can increase your risk of fatal heart attack or stroke. Do not use Motrin Infant Drops just before or after heart bypass surgery (coronary artery bypass graft, or CABG). **This medicine may also cause stomach or intestinal bleeding, which can be fatal.**

Before taking this medicine

Motrin Infant Drops can increase your risk of fatal heart attack or stroke, even if you don't have any risk factors. Do not use this medicine just before or after heart bypass surgery (coronary artery bypass graft, or CABG).

Motrin Infant Drops may also cause stomach or intestinal bleeding, which can be fatal. These conditions can occur without warning while you are using this medicine, especially in older adults.

You should not use Motrin Infant Drops if you are allergic to it, or if you have ever had an asthma attack or severe allergic reaction after taking aspirin or an NSAID.

Ask a doctor or pharmacist if this medicine is safe to use if you have ever had:

- [heart disease](#), [high blood pressure](#), [high cholesterol](#), diabetes, or if you smoke;
- a [heart attack](#), stroke, or blood clot;
- stomach ulcers or bleeding;
- liver or kidney disease;
- asthma; or
- if you take aspirin to prevent heart attack or stroke.

Ask a doctor before using this medicine if you are pregnant or breastfeeding.

If you are pregnant, you should not take Motrin Infant Drops unless your doctor tells you to. Taking an NSAID during the last 20 weeks of pregnancy can cause serious heart or kidney problems in the unborn baby and possible complications with your pregnancy.

Do not give Motrin Infant Drops to a child younger than 6 months old without the advice of a doctor.

How should I take Motrin Infant Drops?

Use exactly as directed on the label, or as prescribed by your doctor. **Use the lowest dose that is effective in treating your condition.**

An Motrin Infant Drops overdose can damage your stomach or intestines. The maximum amount of [ibuprofen](#) for adults is 800 milligrams per dose or 3200 mg per day (4 maximum doses).

A child's dose of ibuprofen is based on the age and weight of the child. **Carefully follow the dosing instructions provided with children's Motrin Infant Drops for the age and weight of your child.** Ask a doctor or pharmacist if you have questions.

Take Motrin Infant Drops with food or milk to lessen [stomach upset](#).

Shake the **oral suspension** (liquid) before you measure a dose. Use the dosing syringe provided, or use a medicine dose-measuring device (not a kitchen spoon).

You must chew the **chewable tablet** before you swallow it.

Store at room temperature away from moisture and heat. Do not allow the liquid medicine to freeze.

 [Detailed Motrin Infant Drops dosage information](#)

What happens if I miss a dose?

Since Motrin Infant Drops is used when needed, you may not be on a dosing schedule. Skip any missed dose if it's almost time for your next dose. **Do not** use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose symptoms may include nausea, vomiting, stomach pain, [drowsiness](#), black or bloody stools, coughing up blood, shallow breathing, fainting, or coma.

What should I avoid while taking Motrin Infant Drops?

Ask a doctor or pharmacist before using other medicines for pain, fever, swelling, or cold/flu symptoms. They may contain ingredients similar to Motrin Infant Drops (such as aspirin, this medicine, [ketoprofen](#), or naproxen).

Avoid taking aspirin unless your doctor tells you to.

If you also take aspirin to prevent stroke or heart attack, taking Motrin Infant Drops can make aspirin less effective in protecting your heart and blood vessels. If you take both medicines, take this medicine at least 8 hours before or 30 minutes after you take aspirin (non-enteric coated form).

Avoid drinking alcohol. It may increase your risk of stomach bleeding.

What other drugs will affect Motrin Infant Drops?

Ask your doctor before using Motrin Infant Drops if you take an antidepressant. Taking certain [antidepressants](#)

with an NSAID may cause you to bruise or bleed easily.

Ask a doctor or pharmacist before using Motrin Infant Drops with any other medications, especially:

- [cyclosporine](#);
- lithium;
- [methotrexate](#);
- a blood thinner (warfarin, Coumadin, Jantoven);
- heart or blood pressure medication, including a diuretic or "water pill"; or
- steroid medicine (such as [prednisone](#)).

This list is not complete. Other drugs may affect Motrin Infant Drops, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). Not all possible [drug interactions](#) are listed here.

 [Motrin Infant Drops drug interactions](#) (more detail)

Does Motrin Infant Drops interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Motrin Infant Drops

+

Enter a drug name

Add

Popular FAQ

- Can I take ibuprofen with Ajovy?

▼
- Naproxen vs ibuprofen: What's the difference?

▼
- Can you take ibuprofen on an empty stomach?

▼
- Can you take expired ibuprofen?

▼
- What's the difference between aspirin and ibuprofen?

▼
- Can you overdose on ibuprofen?

▼
- Can you take tramadol with acetaminophen, ibuprofen, or aspirin?

▼

More FAQ

- [Can you take Ibuprofen if you have COVID-19 \(coronavirus\)?](#)

- [Meloxicam vs Ibuprofen: What's the difference?](#)

More about Motrin Infant Drops (ibuprofen)

- [Check interactions](#)
- [Compare alternatives](#)
- [Latest FDA alerts \(16\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: Nonsteroidal anti-inflammatory drugs](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

[Advil](#), [IBU](#), [Proprinal](#), [Advil Migraine](#), ... +15 more

Professional resources

- [Ibuprofen, Ibuprofen Lysine monograph](#)
- [Ibuprofen Tablets \(FDA\)](#)

Other brands

[Advil](#), [IBU](#), [Advil Liqui-Gels](#), [Advil Migraine](#), ... +5 more

Other formulations

- [Motrin](#)

Related treatment guides

- [Fever](#)
- [Juvenile Rheumatoid Arthritis](#)
- [Pain](#)
- [Spondylolisthesis](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)


Copyright 1996-2025 Cerner Multum, Inc. Version: 24.02.


Motrin Pediatric Dosage Guide

Use these dosage charts for infants and children under 12 years old:

[Motrin Dosage Charts \(Infants & Children\)](#)

DRUG STATUS

Availability
 Rx and/or OTC

Pregnancy & Lactation
 Risk data available

CSA Schedule*
N/A Not a controlled drug



User Reviews & Ratings

[Review this drug](#)

Related News

[NSAIDs Linked To Lower Dementia Risk](#)

[Non-Opioids Can Ease Post-Dental Surgery Pain Better Than Opioids, Study Shows](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.