



[Home](#)

2. Maxifed DMX

Maxifed DMX

Generic name: [dextromethorphan, guaifenesin, and pseudoephedrine](#) [*dex-troe-meth-OR-fan, gwy-e-FEN-e-sin, soo-doe-e-FED-rin*]

Brand names: [Capmist DM](#), [Despec Drops](#), Maxifed DMX, [Poly-Vent DM](#), [Trispec PSE](#), ... [show all 11 brands](#)

Drug class: [Upper respiratory combinations](#)



[Medically reviewed](#) by Drugs.com on Aug 16, 2023. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is Maxifed DMX?

[Dextromethorphan](#) is a cough suppressant. It affects the signals in the brain that trigger cough reflex.

[Guaifenesin](#) is an expectorant. It helps loosen congestion in your chest and throat, making it easier to cough out through your mouth.

[Pseudoephedrine](#) is a decongestant that shrinks blood vessels in the nasal passages. Dilated blood vessels can cause [nasal congestion](#) (stuffy nose).

Maxifed DMX is a combination medicine used to treat cough, stuffy nose, sinus congestion and chest congestion caused by allergies, the [common cold](#), or the flu.

Maxifed DMX will not treat a cough that is caused by smoking, asthma, or emphysema.

Maxifed DMX may also be used for purposes not listed in this medication guide.

Maxifed DMX side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

This medication may cause serious side effects. Stop using this medication and call your doctor at once if you have:

- fast or uneven heart rate;
- severe dizziness or anxiety, feeling like you might pass out;
- severe headache;
- mood changes;
- fever; or

- dangerously [high blood pressure](#) (severe headache, blurred vision, buzzing in your ears, anxiety, chest pain, uneven heartbeats, seizure).

Common side effects of Maxifed DMX may include:

- diarrhea;
- dizziness, drowsiness;
- sleep problems (insomnia); or
- feeling nervous, restless, or irritable.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Maxifed DMX side effects](#) (more detail)

Related/similar drugs

Mucinex D

Mucinex D (guaifenesin and pseudoephedrine) is used to treat nasal and sinus congestion, and to ...

Reviews & ratings

6.1 / 10

[224 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Warnings

Do not use Maxifed DMX if you have used an MAO inhibitor in the past 14 days, such as [isocarboxazid](#), linezolid, [methylene blue](#) injection, [phenelzine](#), rasagiline, selegiline, or [tranylcypromine](#).

Before taking this medicine

You should not use this medicine if you are allergic to dextromethorphan, guaifenesin, or pseudoephedrine.

Do not use this medicine if you have taken an MAO inhibitor in the past 14 days. A dangerous [drug interaction](#) could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Ask a doctor or pharmacist if it is safe for you to take Maxifed DMX if you have:

- high blood pressure or [coronary artery disease](#);
- diabetes;
- glaucoma;
- epilepsy or other seizure disorder;
- [enlarged prostate](#) or urination problems;
- [pheochromocytoma](#) (an adrenal gland tumor); or
- cough with mucus, or cough caused by emphysema or [chronic bronchitis](#).

It is not known whether Maxifed DMX will harm an unborn baby. **Ask a doctor before using this medicine if you are pregnant.**

This medicine may pass into breast milk and may harm a nursing baby. [Decongestants](#) may also slow breast milk production. **Ask a doctor before using this medicine if you are breast-feeding a baby.**

Artificially sweetened cold medicine may contain phenylalanine. If you have [phenylketonuria](#) (PKU), check the medication label to see if the product contains phenylalanine.

How should I take Maxifed DMX?

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. Cough and cold medicine is usually taken only for a short time until your symptoms clear up.

Do not give this medication to a child younger than 4 years old. **Always ask a doctor before giving a cough or cold medicine to a child.** Death can occur from the misuse of cough and cold medicines in very young children.

Do not crush, chew, or break an **extended-release tablet**. Swallow it whole.

Measure liquid medicine with the dosing syringe provided, or with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

Drink extra fluids to help loosen the congestion and lubricate your throat while you are taking this medication.

If you need surgery or medical tests, tell the surgeon or doctor ahead of time if you have taken Maxifed DMX within the past few days.

Do not take for longer than 7 days in a row. Talk with your doctor if your symptoms do not improve after 7 days of treatment, or if you have a fever with a headache or skin rash.

Store at room temperature away from moisture and heat. Do not allow liquid medicine to freeze.

 [Detailed Maxifed DMX dosage information](#)

What happens if I miss a dose?

Since this medicine is taken when needed, you may not be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Maxifed DMX?

Maxifed DMX may cause blurred vision or impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert and able to see clearly.

Drinking alcohol can increase certain side effects of this medicine.

Ask a doctor or pharmacist before using any other cold, cough, or allergy medicine. Maxifed DMX are contained in many combination medicines. Taking certain products together can cause you to get too much of a certain drug. Check the label to see if a medicine contains a decongestant, expectorant, or cough suppressant.

What other drugs will affect Maxifed DMX?

Ask a doctor or pharmacist before using this medicine if you are also using any other drugs, including prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Some medicines can cause unwanted or dangerous effects when used together.** Not all possible interactions are listed in this medication guide.

Taking this medicine with other drugs that make you sleepy or slow your breathing can worsen these effects. Ask your doctor before taking this medicine with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

 [Maxifed DMX drug interactions](#) (more detail)

Does Maxifed DMX interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Maxifed DMX

+

Enter a drug name

Add

More about Maxifed DMX (dextromethorphan / guaifenesin /

pseudoephedrine)

- Check interactions
- Compare alternatives
- Imprints, shape & color data
- Side effects
- Dosage information
- During pregnancy
- Drug class: upper respiratory combinations
- En español

Patient resources

Other brands

Capmist DM, Tusnel, Trispec PSE, Despec DM, ... +6 more

Other formulations

- Maxifed

Related treatment guides

- Cough and Nasal Congestion

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

	Availability
Rx	Rx and/or OTC
OTC	

Pregnancy & Lactation
 Risk data available

CSA Schedule*	
N/A	Not a controlled drug

WADA Class



User Reviews & Ratings

[Review this drug](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Copyright © 2000-2025 Drugs.com. All rights reserved.

