



[Home](#)

2. [Ortho-Novum 7/7/7](#)

Ortho-Novum 7/7/7

Generic name: [ethinyl estradiol and norethindrone \(birth control\)](#) [*ETH-in-il-ess-tra-DYE-ole-and-nor-ETH-in-drone*]

Drug class: [Contraceptives](#)

Medically reviewed by [Sanjai Sinha, MD](#). Last updated on Apr 7, 2025.

[Uses](#) [Warnings](#) [Side effects](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is Ortho-Novum 7/7/7?

Ortho-Novum 7/7/7 is a [combination birth control pill](#) containing female hormones that prevent ovulation (the release of an egg from an ovary). This medication also causes changes in your cervical mucus and uterine lining, making it harder for sperm to reach the uterus and harder for a fertilized egg to attach to the uterus.

Ortho-Novum 7/7/7 is used as contraception to prevent pregnancy. There are many available brands of ethinyl estradiol and norethindrone.

Ortho-Novum 7/7/7 may also be used for purposes not listed in this medication guide.

Warnings

Do not use Ortho-Novum 7/7/7 if you are pregnant or if you have recently had a baby.

You should not use Ortho-Novum 7/7/7 if you have: uncontrolled high blood pressure, heart disease, coronary artery disease, circulation problems (especially with diabetes), undiagnosed vaginal bleeding, liver disease or liver cancer, severe migraine headaches, if you also take certain hepatitis C medication, if you will be having major surgery, if you smoke and are over 35, or if you have ever had a heart attack, a stroke, a blood clot, jaundice caused by pregnancy or birth control pills, or cancer of the breast, uterus/cervix, or vagina.

Taking Ortho-Novum 7/7/7 can increase your risk of blood clots, stroke, or heart attack.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. You should not take Ortho-Novum 7/7/7 if you smoke and are over 35 years old.

Ortho-Novum 7/7/7 side effects

Get emergency medical help if you have **signs of an allergic reaction to Ortho-Novum 7/7/7:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using this medicine and call your doctor at once if you have:

- **signs of a stroke** - sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance;
- **signs of a blood clot** - sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, pain or warmth in one or both legs;
- **heart attack symptoms** - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- **liver problems** - loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- **increased blood pressure** - severe headache, blurred vision, pounding in your neck or ears;
- swelling in your hands, ankles, or feet;
- a change in the pattern or severity of migraine headaches;
- a breast lump; or
- **symptoms of depression** - sleep problems, weakness, tired feeling, mood changes.

Common Ortho-Novum 7/7/7 side effects may include:

- nausea, vomiting;
- breast tenderness;
- breakthrough bleeding;
- headache; or
- problems with contact lenses.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Ortho-Novum 7/7/7 side effects](#) (more detail)

Related/similar drugs

Mirena

Mirena (levonorgestrel) is a plastic intrauterine device placed in the uterus and is used to ...

Reviews & ratings

6.1 / 10

[2,624 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular

management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Depo Provera

Depo-Provera is used as contraception to prevent pregnancy. It is also used to reduce pain cause by ...

Reviews & ratings

4.2 / 10

[1,967 Reviews](#)

[View more](#)

Nexplanon

Nexplanon is a long-term, reversible birth control implant that lasts for three years. Learn about ...

Reviews & ratings

5.0 / 10

[4,864 Reviews](#)

[View more](#)

Yaz

Yaz (drospirenone and ethinyl estradiol) is used as a contraceptive to prevent pregnancy. Includes ...

Reviews & ratings

5.9 / 10

[997 Reviews](#)

[View more](#)

Slynd

Slynd (drospirenone) is a birth control pill used for the prevention of pregnancy. Includes Slynd ...

Reviews & ratings

[View more](#)

Before taking this medicine

Taking Ortho-Novum 7/7/7 can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have high blood pressure, diabetes, high cholesterol, or if you are overweight. Your risk of stroke or blood clot is highest during your first year of taking Ortho-Novum 7/7/7. Your risk is also high when you restart birth control pills after not taking them for 4 weeks or longer.

Smoking can greatly increase your risk of blood clots, stroke, or heart attack. Your risk increases the older you are and the more you smoke. You should not take Ortho-Novum 7/7/7 if you smoke and are over 35 years old.

Do not use if you are pregnant. Stop using this medicine and tell your doctor right away if you become pregnant, or if you miss 2 menstrual periods in a row. If you have recently had a baby, wait at least 4 weeks before taking Ortho-Novum 7/7/7.

You should not take Ortho-Novum 7/7/7 if you have:

- untreated or uncontrolled high blood pressure;
- heart disease (chest pain, coronary artery disease, history of heart attack, stroke, or blood clot);
- an increased risk of having blood clots due to a heart problem or a hereditary blood disorder;
- circulation problems (especially if caused by diabetes);
- a history of hormone-related cancer, or cancer of the breast, uterus/cervix, or vagina;
- unusual vaginal bleeding that has not been checked by a doctor;
- liver disease or liver cancer;
- severe migraine headaches (with aura, numbness, weakness, or vision changes), especially if you are older than 35;
- a history of jaundice caused by pregnancy or birth control pills;
- if you smoke and are over 35 years old; or
- if you take any hepatitis C medication containing ombitasvir/paritaprevir/ritonavir (Technivie).

To make sure Ortho-Novum 7/7/7 are safe for you, tell your doctor if you have ever had:

- heart problems, high blood pressure, or if you are prone to having blood clots;
- high cholesterol or triglycerides, or if you are overweight;
- depression;
- a seizure or migraine headache;
- diabetes, gallbladder disease, underactive thyroid;
- liver or kidney disease;

- irregular menstrual cycles; or
- fibrocystic breast disease, lumps, nodules, or an abnormal mammogram.

The hormones in Ortho-Novum 7/7/7 can pass into breast milk and may harm a nursing baby. This medicine may also slow breast milk production. Do not use if you are breast feeding a baby.

i [Ortho-Novum 7/7/7 pregnancy and breastfeeding warnings](#) (more detail)

How should I take Ortho-Novum 7/7/7?

Take Ortho-Novum 7/7/7 exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Take your first pill on the first day of your period or on the first Sunday after your period begins. You may need to use back-up birth control, such as condoms with spermicide, when you first start using Ortho-Novum 7/7/7. Follow your doctor's instructions.

Take one pill every day, no more than 24 hours apart. When the pills run out, start a new pack the following day. You may get pregnant if you do not take one pill daily. Get your prescription refilled before you run out of pills completely.

Ortho-Novum 7/7/7 packs contain seven green "reminder" pills to keep you on your regular cycle. Your period will usually begin while you are using these reminder pills.

Use a back-up birth control if you are sick with severe vomiting or diarrhea.

You may have breakthrough bleeding, especially during the first 3 months. Tell your doctor if this bleeding continues or is very heavy.

If you need major surgery or will be on long-term bed rest, you may need to stop using this medicine for a short time. Any doctor or surgeon who treats you should know that you are using Ortho-Novum 7/7/7.

While taking Ortho-Novum 7/7/7, you will need to visit your doctor regularly.

Store Ortho-Novum 7/7/7 at room temperature away from moisture and heat.

i [Detailed Ortho-Novum 7/7/7 dosage information](#)

What happens if I miss a dose?

Follow the patient instructions provided in your Ortho-Novum 7/7/7 packet. Missing a pill increases your risk of becoming pregnant.

If you miss 1 active pill, take 2 pills on the day you remember. Then take 1 pill per day for the rest of the pack.

If you miss 2 active pills in a row in Week 1 or 2, take 2 pills per day for 2 days in a row. Then take 1 pill per day for the rest of the pack. Use back-up birth control for at least 7 days following the missed pills.

If you miss 2 active pills in a row in Week 3, throw out the rest of the pack and start a new pack the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss 3 active pills in a row in Week 1, 2, or 3, throw out the rest of the pack and start a new pack on the same day if you are a Day 1 starter. If you are a Sunday starter, keep taking a pill every day until Sunday. On Sunday, throw out the rest of the pack and start a new pack that day.

If you miss 2 or more active pills, you may not have a period during the month. **If you miss a period for 2 months in a row, call your doctor because you might be pregnant.**

If you miss a reminder pill, throw it away and keep taking 1 reminder pill per day until the pack is empty.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose may cause nausea or vaginal bleeding.

What to avoid

Do not smoke while taking Ortho-Novum 7/7/7, especially if you are older than 35 years of age.

Ortho-Novum 7/7/7 will not protect you from sexually transmitted diseases--including HIV and AIDS. Using a condom is the only way to protect yourself from these diseases.

What other drugs will affect Ortho-Novum 7/7/7?

Other drugs may interact with ethinyl estradiol and norethindrone, including prescription and over-the-counter medicines, vitamins, and herbal products. Some drugs can make Ortho-Novum 7/7/7 less effective, which may result in pregnancy. Tell your doctor about all your current medicines and any medicine you start or stop using.

 [Ortho-Novum 7/7/7 drug interactions](#) (more detail)

Does Ortho-Novum 7/7/7 interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Ortho-Novum 7/7/7

+

Enter a drug name

Add

More about Ortho-Novum 7 / 7 / 7 (ethinyl estradiol / norethindrone)

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(20\)](#)
- [Drug images](#)
- [Side effects](#)

- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: contraceptives](#)

Patient resources

Other brands

[Lo Loestrin Fe](#), [Junel Fe 1/20](#), [Loestrin 21 1/20](#), [Larin Fe 1/20](#), ... +32 more

Professional resources

- [Ortho-Novum 777 prescribing information](#)

Other brands

[Lo Loestrin Fe](#), [Blisovi Fe 1/20](#), [Loestrin 21 1/20](#), [Hailey Fe 1/20](#), ... +81 more

Related treatment guides

- [Birth Control](#)

Further information


Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Ortho-Novum 7/7/7 only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 15.02.

DRUG STATUS

Availability	
Rx	Prescription only
Pregnancy & Lactation	
	Risk data available
CSA Schedule*	
N/A	Not a controlled drug
Approval History	
	Drug history at FDA

User Reviews & Ratings

6.8 / 10

[20 Reviews](#)

Related News

[Certain Birth Control Pills, Devices Linked to Higher Heart Risks](#)

Images

[Ortho-novum 7 7 7 ethinyl estradiol 0.035 mg / norethindrone 0.75 mg \(Ortho 75 Ortho 75\)](#)



[View larger images](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)
[Advertising policy](#)
[Attribution & citations](#)

Terms & privacy

[Terms of use](#)
[Editorial policy](#)
[Privacy policy](#)

Support

[Help center](#)
[Sitemap](#)
[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.