

[Home](#)

## 2. Paliperidone

# Paliperidone

**Generic name:** paliperidone (oral) [ *PAL-ee-PER-i-DONE* ]

**Brand name:** [Invega](#)

**Dosage form:** oral tablet, extended release (1.5 mg; 3 mg; 6 mg; 9 mg)

**Drug class:** [Atypical antipsychotics](#)



[Medically reviewed](#) by Drugs.com on Jan 27, 2025. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

## What is paliperidone?

Paliperidone is an antipsychotic medicine that is used to treat [schizophrenia](#) in adults and teenagers who are at least 12 years old.

Paliperidone may also be used for purposes not listed in this medication guide.

## Paliperidone side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

**Stop taking paliperidone and call your doctor at once if you have any of these signs of a serious movement disorder:**

- tremors or shaking in your arms or legs;
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement); or
- any new or unusual muscle movements you cannot control.

**Paliperidone may cause serious side effects. Call your doctor at once if you have:**

- fast or pounding heartbeats, fluttering in your chest, [shortness of breath](#), and sudden dizziness (like you might pass out);
- breast swelling (in women or men), nipple discharge;
- changes in menstrual periods;
- [impotence](#), penis erection that is painful or lasts 4 hours or longer;
- weight gain;

- **low white blood cell counts**--fever, chills, mouth sores, skin sores, [sore throat](#), cough, trouble breathing;
- **high blood sugar**--increased thirst, increased urination, hunger, fruity breath odor; or
- **severe nervous system reaction**--very stiff (rigid) muscles, high fever, fast or pounding heartbeats, [fainting](#).

**Common side effects of paliperidone may include:**

- [drowsiness](#);
- anxiety;
- muscle stiffness, tremors or shaking;
- uncontrolled muscle movements, trouble with walking, balance, or speech;
- weight gain;
- [upset stomach](#), [constipation](#);
- [fast heart rate](#); or
- stuffy nose, sore throat.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Paliperidone side effects](#) (more detail)

## Related/similar drugs

### Fanapt

Fanapt (iloperidone) is an antipsychotic medication used to treat schizophrenia. Includes Fanapt ...

#### Reviews & ratings

6.4 / 10

[30 Reviews](#)

[View more](#)

### Abilify

Abilify is an antipsychotic medicine used to treat the symptoms of schizophrenia and bipolar ...

#### Reviews & ratings

6.1 / 10

[1,187 Reviews](#)

[View more](#)

## Lybalvi

Lybalvi is used to treat adults with manic or mixed episodes associated with bipolar I disorder ...

### Reviews & ratings

8.1 / 10

[16 Reviews](#)

[View more](#)

## Seroquel

Seroquel is used to treat the symptoms of schizophrenia, bipolar disorder and major depressive ...

### Reviews & ratings

7.0 / 10

[694 Reviews](#)

[View more](#)

## Risperidone

Risperidone is used to treat schizophrenia and symptoms of bipolar disorder. Learn about side ...

### Reviews & ratings

5.6 / 10

[731 Reviews](#)

[View more](#)

## Olanzapine

Olanzapine (brand name Zyprexa) is an atypical antipsychotic that may be used to treat adults and ...

### Reviews & ratings

5.9 / 10

[1,091 Reviews](#)

[View more](#)

## Aripiprazole

Easy-to-read patient leaflet for

aripiprazole. Includes indications, proper use, special ...

### Reviews & ratings

6.1 / 10

[1,540 Reviews](#)

[View more](#)

## Quetiapine

Quetiapine is an atypical antipsychotic used to treat schizophrenia, bipolar disorder, and major ...

### Reviews & ratings

7.2 / 10

[2,249 Reviews](#)

[View more](#)

## Fluoxetine

Fluoxetine is used to treat depression or obsessive-compulsive disorder in adults. Learn about side ...

### Reviews & ratings

7.3 / 10

[2,232 Reviews](#)

[View more](#)

## Prozac

Prozac (fluoxetine) is an SSRI antidepressant used to treat depression, OCD, panic disorder ...

### Reviews & ratings

7.2 / 10

[1,239 Reviews](#)

[View more](#)

## Warnings

Paliperidone is not approved for use in older adults with dementia-related [psychosis](#).

## Before taking this medicine

You should not use paliperidone if you are allergic to paliperidone or [risperidone \(Risperdal\)](#).

Paliperidone may increase the risk of death in older adults with dementia-related psychosis and is not approved for this use.

**High doses or long-term use of paliperidone can cause a serious movement disorder that may not be reversible.**

The longer you use paliperidone, the more likely you are to develop this disorder, especially if you are a woman or an older adult.

Tell your doctor if you have ever had:

- heart problems, or a [heart attack](#);
- [long QT syndrome](#) (in you or a family member);
- high or [low blood pressure](#), or fainting spells;
- low white blood cell (WBC) counts;
- a serious neurologic disorder caused by taking an antipsychotic medicine;
- uncontrolled muscle movements in your face;
- a stomach or intestinal disorder;
- liver or kidney disease;
- seizures or epilepsy;
- an electrolyte imbalance (such as [low potassium](#) or magnesium levels in your blood);
- diabetes (paliperidone may raise your blood sugar); or
- [breast cancer](#).

Tell your doctor if you are pregnant or plan to become pregnant while using this medicine.

**Taking antipsychotic medicine in the last 3 months of pregnancy may cause breathing problems, feeding problems, or withdrawal symptoms in the newborn.** However, you may have withdrawal symptoms or other problems if you stop using your medicine during pregnancy. If you get pregnant, tell your doctor right away. **Do not stop using paliperidone without your doctor's advice.**

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of paliperidone on the baby.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk.

## How should I take paliperidone?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

You may take paliperidone with or without food.

Swallow the tablet whole and do not crush, chew, or break it.

Your doctor will need to check your progress on a regular basis.

Store at room temperature away from moisture, heat, and light.

 [Detailed Paliperidone dosage information](#)

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

Get your prescription refilled before you run out of medicine completely.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include severe drowsiness, fast heartbeats, and fainting.

## What should I avoid while taking paliperidone?

While you are taking paliperidone, you may be more sensitive to heat. Avoid getting too hot, or becoming overheated or dehydrated. Drink plenty of fluids, especially in hot weather and during exercise. It is easier to become dangerously overheated and dehydrated while you are taking paliperidone.

**Avoid drinking alcohol.** Dangerous side effects could occur.

Avoid driving or hazardous activity until you know how paliperidone will affect you. Dizziness or drowsiness can cause falls, accidents, or severe injuries.

Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

## What other drugs will affect paliperidone?

Using paliperidone with other drugs that make you dizzy or lower your blood pressure can worsen these effects. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

**Sometimes it is not safe to use certain medications at the same time.** Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

**Many drugs can affect paliperidone.** This includes prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Not all possible interactions are listed here.** Tell your doctor about all your current medicines and any medicine you start or stop using.

 [Paliperidone drug interactions](#) (more detail)

## Does paliperidone interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

paliperidone

+

Enter a drug name

Add

## Popular FAQ

Can you drink alcohol while taking Invega Sustenna?



Invega Sustenna vs Invega Trinza vs Invega Hafyera. What's the difference?



How to transition to Invega Trinza from Invega Sustenna ?



How do you give an Invega Sustenna injection?



How to transition to Invega Hayfera from Invega Trinza or Invega Sustenna?



How is Invega Trinza administered?



## More FAQ

- [What drugs cause tardive dyskinesia?](#)

[View more FAQ...](#)

## More about paliperidone

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(642\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: atypical antipsychotics](#)
- [Breastfeeding](#)

## Patient resources

- [Paliperidone injection drug information](#)

- [Paliperidone \(Intramuscular\) \(Advanced Reading\)](#)
- [Paliperidone \(Oral\) \(Advanced Reading\)](#)
- [Paliperidone Extended-Release Tablets](#)

## Other brands

[Invega](#), [Invega Sustenna](#), [Invega Trinza](#), [Invega Hafyera](#), [Erzofri](#)

## Professional resources

- [Paliperidone monograph](#)
- [Paliperidone \(FDA\)](#)
- [Paliperidone ER Injection \(FDA\)](#)

## Other brands

[Invega](#), [Invega Sustenna](#), [Invega Trinza](#), [Invega Hafyera](#), [Erzofri](#)

## Related treatment guides

- [Bipolar Disorder](#)
- [Schizoaffective Disorder](#)
- [Schizophrenia](#)
- [Borderline Personality Disorder](#)
- [Autism](#)
- [Depression](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

### [Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.02.

### DRUG STATUS

#### Availability

**Rx** Prescription only

#### [Pregnancy & Lactation](#)

 Risk data available




### CSA Schedule\*

N/A Not a controlled drug



### Approval History

 Drug history at FDA



## User Reviews & Ratings

4.4 / 10

[642 Reviews](#)

## Images

[Paliperidone Extended-Release 6 mg \(AN 82\)](#)



[View larger images](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



### About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

### Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

### Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Copyright © 2000-2025 Drugs.com. All rights reserved.