

Home

2. Lyrica

Lyrica

Generic name: pregabalin [pre-GAB-a-lin]

Brand names: Lyrica, Lyrica CR

Drug class: Gamma-aminobutyric acid analogs

Medically reviewed by Sophia Entringer, PharmD. Last updated on Nov 18, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is Lyrica?

Lyrica was originally FDA approved as an anti-epileptic drug, also called an anticonvulsant. It works by slowing down impulses in the brain that cause seizures. Pregabalin also affects chemicals in the brain that send pain signals across the nervous system.

Lyrica is used to treat pain caused by fibromyalgia, or nerve pain in people with diabetes (diabetic neuropathy), herpes zoster (post-herpetic neuralgia), or spinal cord injury.

Lyrica may also be used for purposes not listed in this medication guide.

Warnings

Lyrica can cause a severe allergic reaction. Stop taking this medicine and seek emergency medical help if you have hives or blisters on your skin, trouble breathing, or swelling in your face, mouth, or throat.

Some people have thoughts about suicide while taking Lyrica. Stay alert to changes in your mood or symptoms. Report any new or worsening symptoms to your doctor.

If you have diabetes or heart problems, call your doctor if you have weight gain or swelling in your hands or feet while taking Lyrica.

Do not stop using this medicine suddenly, even if you feel fine. Stopping suddenly may cause withdrawal symptoms.

Do not change your dose without your doctor's advice. Tell your doctor if the medication does not seem to work as well in treating your condition.

Before taking this medicine

You should not use Lyrica if you are allergic to pregabalin.

To make sure Lyrica is safe for you, tell your doctor if you have ever had:

- lung disease, such as chronic obstructive pulmonary disease (COPD);
- a mood disorder, depression, or suicidal thoughts;
- heart problems (especially congestive heart failure);
- a bleeding disorder, or low levels of platelets in your blood;
- kidney disease (or if you are on dialysis);
- diabetes (unless you are taking pregabalin to treat diabetic neuropathy);
- · drug or alcohol addiction; or
- a severe allergic reaction (angioedema).

Do not give this medicine to a child without medical advice.

- Pregabalin is not approved for use by anyone younger than 18 years old to treat nerve pain caused by fibromyalgia, diabetes, herpes zoster, or spinal cord injury.
- Pregabalin is not approved for seizures in anyone younger than 1 month old.

Some people have thoughts about suicide while taking Lyrica. Your doctor will need to check your progress at regular visits. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Follow your doctor's instructions about taking seizure medication if you are pregnant. Seizure control is very important during pregnancy, and having a seizure could harm both mother and baby. **Do not start or stop taking this medicine** without your doctor's advice, and tell your doctor right away if you become pregnant.

If you are pregnant, your name may be listed on a pregnancy registry to track the effects of pregabalin on the baby.

Pregabalin can temporarily decrease sperm count and may affect fertility in men (your ability to have children). In animal studies, pregabalin also caused birth defects in the offspring of males treated with this medicine. However, it is not known whether these effects would occur in humans. Ask your doctor about your risk.

You should not breastfeed while using pregabalin.

1 Lyrica pregnancy and breastfeeding warnings (more detail)

How should I take Lyrica?

Take Lyrica exactly as prescribed by your doctor. Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Take the medicine at the same time each day, with or without food.

Do not crush, chew, or break an extended-release tablet. Swallow the tablet whole.

Measure liquid medicine with the dosing syringe provided, or with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

Call your doctor if your symptoms do not improve, or if they get worse.

Do not stop using Lyrica suddenly, even if you feel fine. Stopping suddenly may cause increased seizures or unpleasant withdrawal symptoms. Follow your doctor's instructions about tapering your dose for at least 1 week before

stopping completely.

In case of emergency, wear or carry medical identification to let others know you take seizure medication.

Store at room temperature away from moisture, heat, and light.

Detailed Lyrica dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What to avoid

Avoid drinking alcohol. It may increase certain side effects of Lyrica.

Avoid driving or hazardous activity until you know how this medicine will affect you. Your reactions could be impaired.

Lyrica side effects

Lyrica can cause a severe allergic reaction. Stop taking this medicine and get emergency medical help if you have: hives or blisters on your skin; difficult breathing; swelling of your face, lips, tongue, or throat.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself.

Call your doctor at once if you have:

- weak or shallow breathing;
- blue-colored skin, lips, fingers, and toes;
- confusion, extreme drowsiness or weakness;
- vision problems;
- skin sores (if you have diabetes);
- · easy bruising, unusual bleeding;
- swelling in your hands or feet, rapid weight gain (especially if you have diabetes or heart problems); or
- unexplained muscle pain, tenderness, or weakness (especially if you also have fever or don't feel well).

Pregabalin can cause life-threatening breathing problems. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Breathing problems may be more likely in older adults or in people with COPD.

If you have diabetes, tell your doctor right away if you have any new sores or other skin problems.

Common Lyrica side effects may include:

- dizziness, drowsiness;
- · swelling in your hands and feet;
- trouble concentrating;
- · increased appetite;
- · weight gain;
- dry mouth; or
- blurred vision.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Lyrica side effects (more detail)

Related/similar drugs

Lamictal

Lamictal is an anti-epileptic medication used treat seizures in adults and children over 2 years ...

Reviews & ratings

8.0 / 10

792 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Diazepam

Diazepam is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.4 / 10

884 Reviews

View more

Cymbalta

Cymbalta (duloxetine) is used to treat major depressive disorder, general anxiety disorder and ...

Reviews & ratings

6.3 / 10

1,899 Reviews

View more

Keppra

Keppra is an anti-epileptic drug used to treat partial onset seizures in people with epilepsy ...

Reviews & ratings

5.7 / 10

307 Reviews

View more

Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

Reviews & ratings

7.1 / 10

484 Reviews

View more

Clonazepam

Clonazepam is used to treat seizures and panic disorder. Learn about side effects, interactions and ...

Reviews & ratings

View more

What other drugs will affect Lyrica?

Using Lyrica with other drugs that slow your breathing can cause dangerous side effects or death. Ask your doctor before using opioid medication, a sleeping pill, cold or allergy medicine, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- oral diabetes medicine pioglitazone, rosiglitazone; or
- an ACE inhibitor benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril, or trandolapril.
- · any other seizure medications

This list is not complete. Other drugs may interact with pregabalin, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Lyrica drug interactions (more detail)

Does Lyrica interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Lyrica	
+	
Enter a drug name	Add

Popular FAQ

How long does it take for Lyrica to work?	~
Does Lyrica cause weight gain?	~
How long does Lyrica stay in your system?	~
Can you stop Lyrica cold turkey?	~
Is Lyrica a controlled substance / narcotic?	~

Does Lyrica make you sleepy?	~
Is there a generic for Lyrica?	~
What is Lyrica used to treat?	~

More FAQ

• Lyrica vs Gabapentin: What's the difference?

View more FAQ...

More about Lyrica (pregabalin)

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (925)
- Drug images
- Side effects
- Dosage information
- Patient tips
- During pregnancy
- · Generic availability
- Support group
- FDA approval history
- Drug class: gamma-aminobutyric acid analogs
- Breastfeeding
- En español

Patient resources

- Lyrica (Pregabalin Capsules) patient information
- Lyrica (Pregabalin Oral Solution)

Other brands

Lyrica CR

Professional resources

• Lyrica prescribing information

• Pregabalin (AHFS Monograph)

Other brands

Lyrica CR

Related treatment guides

- Neuropathic Pain
- Fibromyalgia
- Neuralgia
- Diabetic Peripheral Neuropathy
- Postherpetic Neuralgia
- Epilepsy

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Lyrica only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 9.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ം Risk data available

CSA Schedule*

5 Some potential for abuse

Approval History

10+ years FDA approved 2004

User Reviews & Ratings

6.4 / 10

925 Reviews

Images





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







