

[Home](#)

2. Travoprost

Travoprost ophthalmic

Generic name: travoprost ophthalmic [*TRA-voe-prost-off-THAL-mik*]**Brand names:** [Travatan Z](#), Travatan, [iDose TR](#)**Dosage form:** ophthalmic solution (0.004%)**Drug class:** [Ophthalmic glaucoma agents](#)[Medically reviewed](#) by Drugs.com on Aug 14, 2023. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is travoprost ophthalmic?

Travoprost lowers pressure inside the eye by increasing the amount of fluid that drains from the eye.

Travoprost ophthalmic (for the eyes) is used to treat certain types of [glaucoma](#) and other causes of high pressure inside the eye.

Travoprost ophthalmic may also be used for purposes not listed in this medication guide.

Travoprost ophthalmic side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Travoprost may cause serious side effects. Stop using travoprost and call your doctor at once if you have:

- eye swelling, redness, severe discomfort, crusting or drainage (may be signs of [infection](#));
- red, swollen, or itchy eyelids;
- increased sensitivity to light;
- vision changes; or
- severe burning, stinging, or irritation after using travoprost ophthalmic.

Common side effects of travoprost ophthalmic may include:

- pain, [itching](#), or redness of the eyes;
- puffy eyelids; or
- blurred vision.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088.

 [Travoprost ophthalmic side effects](#) (more detail)

Related/similar drugs

Lumigan

Lumigan is used to treat open angle glaucoma. Learn about side effects, interactions and indications.

Reviews & ratings

5.2 / 10

[25 Reviews](#)

[View more](#)

Xalatan

Xalatan (latanoprost) is used to treat certain types of glaucoma. Includes Xalatan side effects ...

Reviews & ratings

7.3 / 10

[13 Reviews](#)

[View more](#)

FEATURED

Omvox

Omvox is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Cosopt

Cosopt reduces the amount of fluid in the eye and is used to treat glaucoma. Learn about side ...

Reviews & ratings

4.4 / 10

[8 Reviews](#)

[View more](#)

Combigan

Combigan (brimonidine and timolol) is used to treat glaucoma or ocular hypertension. Includes ...

Reviews & ratings

4.9 / 10

[25 Reviews](#)

[View more](#)

Alphagan P

Alphagan P is used to treat glaucoma or ocular hypertension (high pressure inside the eye). Learn ...

Reviews & ratings

5.0 / 10

[8 Reviews](#)

[View more](#)

Warnings

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Before taking this medicine

You should not use travoprost ophthalmic if you are allergic to it.

To make sure travoprost ophthalmic is safe for you, tell your doctor if you have ever had:

- swelling or an infection in your eye;
- retinal detachment; or
- eye [surgery](#) or injury affecting the lens of your eye.

Travoprost ophthalmic may cause a gradual change in the color of your eyes or eyelids and lashes, usually an increase in brown pigment. You may also notice increased growth or thickness of your eyelashes. These changes occur slowly and you may not notice them for months or years. Color changes may be permanent even after your treatment ends, and may occur only in the eye being treated. This could result in a cosmetic difference in eye or eyelash color from one eye to the other.

It is not known whether travoprost ophthalmic will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

It is not known whether travoprost ophthalmic passes into breast milk or if it could affect the nursing baby. Tell your doctor if you are breast-feeding.

Travoprost ophthalmic is not approved for use by anyone younger than 16 years old.

How should I use travoprost ophthalmic?

Follow all directions on your prescription label. Do not use this medicine in larger or smaller amounts or for longer than recommended.

The usual dose of this medicine is 1 drop into the affected eye every evening. Follow your doctor's dosing instructions very carefully.

Do not use this medicine while wearing contact lenses. Travoprost ophthalmic may contain a preservative that can discolor soft contact lenses. Wait at least 15 minutes after using this medicine before putting in your contact lenses.

Wash your hands before using the eye drops.

To apply the **eye drops**:

- Tilt your head back slightly and pull down your lower eyelid to create a small pocket. Hold the dropper above the eye with the tip down. Look up and away from the dropper and squeeze out a drop.
- Close your eyes for 2 or 3 minutes with your head tipped down, without blinking or squinting. Gently press your finger to the inside corner of the eye for about 1 minute, to keep the liquid from draining into your tear duct.
- Wait at least 5 minutes before using any other eye drops your doctor has prescribed.

Do not touch the tip of the eye dropper or place it directly on your eye. A contaminated dropper can infect your eye, which could lead to serious vision problems.

Do not use the eye drops if the liquid has changed colors or has particles in it. Call your pharmacist for new medicine.

Tell your doctor right away if you have any eye injury or eye infection.

If you need surgery (including eye surgery), tell the surgeon ahead of time that you are using travoprost ophthalmic. You may need to stop using the medicine for a short time.

Store at cool room temperature, away from moisture and heat. Keep the bottle tightly closed when not in use.

 [Detailed Travoprost ophthalmic dosage information](#)

What happens if I miss a dose?

Skip the missed dose and use the medicine at the next regularly scheduled time. **Do not** use extra medicine to make up the missed dose.

What happens if I overdose?

An overdose of travoprost ophthalmic is not expected to be dangerous. Seek emergency medical attention or call the Poison Help line at 1-800-222-1222 if anyone has accidentally swallowed the medication.

What should I avoid while using travoprost ophthalmic?

Do not use other eye medications unless your doctor tells you to.

What other drugs will affect travoprost ophthalmic?

It is not likely that other drugs you take orally or inject will have an effect on travoprost used in the eyes. But many drugs can interact with each other. Tell each of your healthcare providers about all medicines you use, including prescription and over-the-counter medicines, [vitamins](#), and [herbal products](#).

 [Travoprost ophthalmic drug interactions](#) (more detail)

Does travoprost ophthalmic interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

travoprost ophthalmic

+

Enter a drug name

Add

More about travoprost ophthalmic

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(15\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: ophthalmic glaucoma agents](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

- [Travoprost advanced reading](#)

- [Travoprost](#)

Other brands

[Travatan Z](#), [iDose TR](#), [Izba](#)

Professional resources

- [Travoprost monograph](#)
- [Travoprost Ophthalmic Solution \(FDA\)](#)

Other brands

[Travatan Z](#)

Related treatment guides

- [Glaucoma, Open Angle](#)
- [Intraocular Hypertension](#)

Further information



Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

| | |
|---|-----------------------|
| Availability | |
| Rx | Prescription only |
| Pregnancy & Lactation | |
|  | Risk data available |
| CSA Schedule* | |
| N/A | Not a controlled drug |
| Approval History | |
|  | Drug history at FDA |



User Reviews & Ratings

5.8 / 10

[15 Reviews](#)

Images

[IDose TR \(travoprost\) 75 mcg intracameral implant](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.