

[Home](#)

2. Nardil

# Nardil

**Generic name:** [phenelzine](#) [ *FEN-el-zeen* ]**Drug class:** [Monoamine oxidase inhibitors](#)[Medically reviewed](#) by Drugs.com on Feb 21, 2025. Written by [Cerner Multum](#).[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#) [FAQ](#)

## What is Nardil?

Nardil is a monoamine oxidase inhibitor (MAOI) that is used to treat symptoms of atypical depression in adults when other medicines have not been effective.

Nardil is not for treating severe depression or [bipolar disorder](#) ([manic depression](#)).

Nardil may also be used for purposes not listed in this medication guide.

## Nardil side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

**Report any new or worsening symptoms to your doctor**, such as: mood or behavior changes, anxiety, [panic attacks](#), trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

**Nardil may cause serious side effects. Call your doctor at once if you have:**

- sudden and severe headache, neck pain or stiffness;
- pounding heartbeats or fluttering in your chest;
- fast or slow heartbeats;
- chest pain;
- a light-headed feeling, like you might pass out;
- sweating (sometimes with fever and sometimes with cold, clammy skin);
- nausea, vomiting; or
- dilated pupils (your eyes may be more sensitive to light).

**Common side effects of Nardil may include:**

- dizziness, feeling light-headed;
- drowsiness, sleep problems;
- headache;
- feeling weak or tired;
- tremors, [muscle twitching](#);
- dry mouth, stomach discomfort, constipation;
- swelling, weight gain; or
- sexual problems.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Nardil side effects](#) (more detail)

## Related/similar drugs

### Lexapro

Lexapro is used to treat anxiety and major depressive disorder. Learn about side effects ...

#### Reviews & ratings

7.6 / 10

[2,376 Reviews](#)

[View more](#)

### Zoloft

Zoloft is an antidepressant used to treat major depression, obsessive-compulsive disorder, panic ...

#### Reviews & ratings

7.3 / 10

[1,972 Reviews](#)

[View more](#)

#### FEATURED

### Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

#### Reviews & ratings

#### Reviews & ratings

5.3 / 10

[685 Reviews](#)

[View more](#)

### Seroquel

Seroquel is used to treat the symptoms of schizophrenia, bipolar disorder and major depressive ...

#### Reviews & ratings

7.0 / 10

[694 Reviews](#)

[View more](#)

### Prozac

Prozac (fluoxetine) is an SSRI antidepressant used to treat depression, OCD, panic disorder ...

#### Reviews & ratings

7.2 / 10

[1,239 Reviews](#)

[View more](#)

### Cymbalta

Cymbalta (duloxetine) is used to treat major depressive disorder, general anxiety disorder and ...

#### Reviews & ratings

6.3 / 10

[1,899 Reviews](#)

[View more](#)

### Wellbutrin

Wellbutrin (bupropion) is used to treat major depressive disorder and seasonal affective disorder ...

#### Reviews & ratings

6.8 / 10

[756 Reviews](#)

[View more](#)

## Celexa

Celexa (citalopram) is an antidepressant in a group of drugs called selective serotonin reuptake ...

### Reviews & ratings

7.3 / 10

[866 Reviews](#)

[View more](#)

## Concerta

Concerta is used to treat attention deficit disorder (ADD) and attention deficit hyperactivity ...

### Reviews & ratings

6.9 / 10

[424 Reviews](#)

[View more](#)

## Citalopram

Citalopram is an antidepressant (selective serotonin reuptake inhibitor) and is used to treat ...

### Reviews & ratings

7.4 / 10

[2,231 Reviews](#)

[View more](#)

## Warnings

Some young people have thoughts about suicide when first taking an antidepressant. Stay alert to changes in your mood or symptoms. **Report any new or worsening symptoms to your doctor.**

**There are many other drugs, foods, and beverages that can cause dangerously high blood pressure if you take them together with Nardil.** Tell your doctor about all your current medicines and any you start or stop using. Avoid drinking alcohol, and learn about the foods you should avoid.

**Symptoms of dangerously high blood pressure include:** a sudden and severe headache, nausea, vomiting, fever,

cold sweat, dilated pupils, light sensitivity, fast or pounding heartbeats, neck stiffness, weakness, or problems with vision or speech.

## Before taking this medicine

You should not use Nardil if you are allergic to it, or if you have:

- [pheochromocytoma](#) (tumor of the adrenal gland).
- congestive [heart failure](#);
- severe kidney disease; or
- a history of liver problems or abnormal liver function tests.

**A dangerous drug interaction can occur between Nardil and certain other medicines you use within 14 days before or after taking this medicine.** Your doctor may change your treatment plan if you need to use any of these medicines, including:

- epinephrine or [norepinephrine](#);
- bupropion, buspirone, [carbamazepine](#), guanethidine, levodopa, meperidine, methyldopa, tryptophan;
- ADHD medication;
- diet pills, cough and cold or allergy medicines;
- opioid medicine;
- other [antidepressants](#);
- **another MAO inhibitor**--furazolidone, [isocarboxazid](#), linezolid, [methylene blue](#) injection, [procarbazine](#), rasagiline, selegiline, [tranylcypromine](#), and others; or
- **drugs that affect serotonin levels in your body**--stimulant medicine, opioid medicine, [herbal products](#), or medicine for depression, mental illness, [Parkinson's disease](#), migraine headaches, serious infections, or prevention of nausea and vomiting.

**This is not a complete list and there may be other medicines you should not take while you are taking Nardil.**

Tell your doctor if you have ever had:

- [high blood pressure](#), [heart disease](#);
- diabetes;
- [schizophrenia](#);
- epilepsy or other seizure disorder; or
- if you have taken another antidepressant within the past 5 weeks.

Some young people have thoughts about suicide when first taking an antidepressant. Your doctor should check your progress at regular visits. Your family or other caregivers should also be alert to changes in your mood or symptoms.

Tell your doctor if you are pregnant or breastfeeding.

Nardil is not approved for use by anyone younger than 18 years old.

## How should I take Nardil?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Your blood pressure will need to be checked often.

You may need to stop using Nardil for a short time before any type of surgery or medical procedure. **Tell any doctor who treats you that you take this medicine.**

It may take a few weeks before you receive the full benefit of taking Nardil. Your doctor will determine how long to treat you with Nardil.

**Do not stop using Nardil suddenly**, or you could have unpleasant withdrawal symptoms. Ask your doctor how to safely stop using this medicine.

Store at room temperature away from moisture and heat.

 [Detailed Nardil dosage information](#)

## What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

## What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. Overdose symptoms may include feeling drowsy or dizzy, severe headache, neck pain or stiffness, hallucinations, shallow breathing, fast and uneven heart rate, cold sweats, feeling like you might pass out, or seizure (convulsions).

## What should I avoid while taking Nardil?

**Do not drink alcohol.** Dangerous side effects could occur.

You must not eat certain foods, including:

- air dried, aged, smoked, or fermented meats, including sausage, pepperoni, Lebanon bologna, or salami;
- beer, wine, reduced-alcohol or alcohol-free beer or wine;
- pickled herring;
- aged cheeses, including blue, boursault, brie, camembert, cheddar, gruyere, mozzarella, parmesan, Romano, Roquefort, and Swiss;
- moldy or improperly stored meat, fish, poultry, or liver;
- soy sauce, miso soup, bean curd, fava beans;
- sauerkraut;
- yogurt;

- yeast extracts, Marmite; or
- excessive amounts of chocolate or caffeine.

**Eating these foods while you are taking Nardil can raise your blood pressure to dangerous levels**, causing life-threatening side effects. Also avoid these foods for 2 weeks after you stop taking this medicine.

Nardil may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert.

## What other drugs will affect Nardil?

**When you start or stop taking Nardil**, your doctor may need to adjust the doses of any other medicines you take on a regular basis.

**There are many other drugs that can cause serious medical problems if you take them together with Nardil.** This includes prescription and over-the-counter medicines, vitamins, and herbal products. **Not all possible interactions are listed in this medication guide.** Tell your doctor about all your current medicines and any medicine you start or stop using.

 [Nardil drug interactions](#) (more detail)

## Does Nardil interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Nardil

+

Enter a drug name

Add

## Frequently asked questions

- [What are some common side effects of antidepressants?](#)

## More about Nardil (phenelzine)

- [Check interactions](#)
- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(124\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)

- [Generic availability](#)
- [Drug class: monoamine oxidase inhibitors](#)
- [Breastfeeding](#)
- [En español](#)

## Patient resources

- [Nardil advanced reading](#)

## Professional resources

- [Nardil prescribing information](#)
- [Phenelzine \(AHFS Monograph\)](#)

## Related treatment guides

- [Depression](#)

## Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

### [Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 3.01.

### DRUG STATUS

#### Availability

**Rx** Prescription only

#### [Pregnancy & Lactation](#)

 Risk data available

#### CSA Schedule\*

**N/A** Not a controlled drug

#### Approval History

**10+ years** FDA approved 1961



## User Reviews & Ratings

8.0 / 10

[124 Reviews](#)



## Images

[Nardil 15 mg \(PD 270\)](#)

## Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



### About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

### Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

### Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices  
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.