

Home

2. Prednisone

Prednisone do

Pronunciation: *PRED-ni-sone* **Generic name:** prednisone

Brand names: Rayos, Sterapred, Deltasone

Drug class: Glucocorticoids

Medically reviewed by Melisa Puckey, BPharm. Last updated on May 13, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is prednisone?

Prednisone is a corticosteroid medicine used to decrease inflammation and keep your immune system in check, if it is overactive. Prednisone is used to treat allergic disorders, skin conditions, ulcerative colitis, Crohn's disease, arthritis, lupus, psoriasis, asthma, chronic obstructive pulmonary disease (COPD) and many more conditions.

Warnings

Prednisone treats many different conditions such as allergic disorders, skin conditions, ulcerative colitis, arthritis, lupus, psoriasis, or breathing disorders.

You should avoid taking prednisone if you have a fungal infection that requires oral antifungals. Topical antifungals may not be an issue, but always let your doctor know what medicines you're taking before starting this medicine.

Steroid medication can weaken your immune system, making it easier for you to get an infection. Avoid being near people who are sick or have infections. Do not receive a "live" vaccine while using prednisone.

Call your doctor at once if you have shortness of breath, severe pain in your upper stomach, bloody or tarry stools, severe depression, changes in personality or behavior, vision problems, or eye pain.

You should not stop using prednisone suddenly. Follow your doctor's instructions about tapering your dose.

Before taking this medicine

You should not use this medication if you are allergic to prednisone, or if you have a fungal infection that requires oral antifungal treatment.

Steroid medication can weaken your immune system, making it easier for you to get an infection or worsening an infection you already have or have recently had. Tell your doctor about any illness or infection you have had within the past several weeks.

To make sure prednisone is safe for you, tell your doctor if you have:

- any illness that causes diarrhea;
- liver disease (such as cirrhosis);
- · kidney disease;
- · heart disease, high blood pressure, low levels of potassium in your blood;
- a thyroid disorder;
- · diabetes;
- a history of malaria;
- · tuberculosis;
- · osteoporosis;
- glaucoma, cataracts, or herpes infection of the eyes;
- stomach ulcers, ulcerative colitis, or a history of stomach bleeding;
- a muscle disorder such as myasthenia gravis; or
- · depression or mental illness.

Long-term use of steroids may lead to bone loss (osteoporosis), especially if you smoke, if you do not exercise, if you do not get enough vitamin D or calcium in your diet, or if you have a family history of osteoporosis. Talk with your doctor about your risk of osteoporosis.

Prednisone can cause low birth weight or birth defects if you take the medicine during your first trimester. Tell your doctor if you are pregnant or plan to become pregnant while using this medication. Use effective birth control.

Prednisone can pass into breast milk and may harm a nursing baby. Tell your doctor if you are breast-feeding a baby.

Steroids can affect growth in children. Talk with your doctor if you think your child is not growing at a normal rate while using this medicine.

Prednisone pregnancy and breastfeeding warnings (more detail)

How should I take prednisone?

Take prednisone exactly as prescribed by your doctor. Follow all directions on your prescription label. Your doctor may change your dose to make sure you get the best results. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Take prednisone with food.

Your dosage needs may change if you have any unusual stress such as a serious illness, fever or infection, or if you have surgery or a medical emergency. Do not change your medication dose or schedule without your doctor's advice.

Measure liquid medicine with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

Do not crush, chew, or break a delayed-release tablet. Swallow it whole.

While using this medicine, you may need frequent blood tests at your doctor's office. Your blood pressure may also need to be checked.

This medication can cause unusual results with certain medical tests. Tell any doctor who treats you that you are using this medicine.

You should not stop using prednisone suddenly. Follow your doctor's instructions about tapering your dose.

Wear a medical alert tag or carry an ID card stating that you take prednisone. Any medical care provider who treats you should know that you are using a steroid.

Store at room temperature away from moisture and heat.

i Detailed Prednisone dosage information

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

An overdose of prednisone is not expected to produce life threatening symptoms. However, long term use of high steroid doses can lead to symptoms such as thinning skin, easy bruising, changes in the shape or location of body fat (especially in your face, neck, back, and waist), increased acne or facial hair, menstrual problems, impotence, or loss of interest in sex.

What should I avoid?

Avoid being near people who are sick or have infections. Call your doctor for preventive treatment if you are exposed to chicken pox or measles. These conditions can be serious or even fatal in people who are using a steroid.

Do not receive a "live" vaccine while using this medicine. Prednisone may increase your risk of harmful effects from a live vaccine. Live vaccines include measles, mumps, rubella (MMR), rotavirus, yellow fever, varicella (chickenpox), one type of the typhoid vaccine and nasal flu (influenza) vaccine.

Avoid drinking alcohol while you are taking prednisone.

Prednisone side effects

Get emergency medical help if you have any of these signs of an allergic reaction to prednisone: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- blurred vision, eye pain, or seeing halos around lights;
- swelling, rapid weight gain, feeling short of breath;

- severe depression, feelings of extreme happiness or sadness, changes in personality or behavior, seizure (convulsions);
- bloody or tarry stools, coughing up blood;
- pancreatitis (severe pain in your upper stomach spreading to your back, nausea and vomiting, fast heart rate);
- low potassium (confusion, uneven heart rate, extreme thirst, increased urination, leg discomfort, muscle weakness or limp feeling); or
- dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure).

Other common prednisone side effects may include:

- sleep problems (insomnia), mood changes;
- increased appetite, gradual weight gain;
- · acne, increased sweating, dry skin, thinning skin, bruising or discoloration;
- slow wound healing;
- headache, dizziness, spinning sensation;
- nausea, stomach pain, bloating; or
- changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Prednisone side effects (more detail)

Related/similar drugs

Nplate

Nplate is used to prevent bleeding episodes in people with chronic immune thrombocytopenic purpura ...

Reviews & ratings

7.5 / 10

11 Reviews

View more

Otezla

Otezla (apremilast) is used to treat plaque psoriasis, psoriatic arthritis, and oral ulcers ...

Reviews & ratings

View more

FEATURED

Benlysta

Benlysta infusion is used to treat active systemic lupus erythematosus (SLE) and active lupus ...

Reviews & ratings

7.9 / 10

63 Reviews

View more

Saphnelo

Saphnelo is used for moderate to severe SLE (lupus) for adults who are also receiving other lupus ...

Reviews & ratings

8.4 / 10

10 Reviews

View more

Skyrizi

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

Reviews & ratings

7.9 / 10

56 Reviews

View more

Dupixent

Dupixent may be used to treat eczema, eosinophilic or oral-corticosteroid-dependent asthma, chronic ...

Reviews & ratings

7.2 / 10

499 Reviews

View more

Breztri Aerosphere

Breztri

(budesonide/glycopyrrolate/formoterol fumarate) is a combination inhaler that may be used ...

Reviews & ratings

6.9 / 10

119 Reviews

View more

Kesimpta

Kesimpta (ofatumumab) is used to treat relapsing forms of multiple sclerosis. Includes Kesimpta ...

Reviews & ratings

6.8 / 10

74 Reviews

View more

Xeljanz

Xeljanz (tofacitinib) is an oral Janus kinase (JAK) inhibitor that may be used alone or with other ...

Reviews & ratings

5.7 / 10

145 Reviews

View more

Stelara

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

Reviews & ratings

7.6 / 10

252 Reviews

View more

What other drugs will affect prednisone?

Many drugs can interact with prednisone. Not all possible interactions are listed here. Tell your doctor about all your medications and any you start or stop using during treatment with prednisone, especially:

- amphotericin B;
- cyclosporine;
- · digoxin, digitalis;
- St. John's wort;
- an antibiotic such as clarithromycin or telithromycin;
- antifungal medication such as itraconazole, ketoconazole, posaconazole, voriconazole;
- · birth control pills and other hormones;
- a blood thinner such as warfarin, Coumadin;
- a diuretic or "water pill";
- the hepatitis C medications boceprevir or telaprevir;
- HIV or AIDS medicine such as atazanavir, delavirdine, efavirenz, fosamprenavir, indinavir, nelfinavir, nevirapine, ritonavir, saquinavir;
- insulin or diabetes medications you take by mouth;
- a non-steroidal anti-inflammatory drug (NSAID) such as aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others;
- seizure medications such as carbamazepine, fosphenytoin, oxcarbazepine, phenobarbital, phenytoin, primidone; or
- the tuberculosis medications isoniazid, rifabutin, rifapentine, or rifampin.

This list is not complete and many other drugs can interact with prednisone. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Give a list of all your medicines to any healthcare provider who treats you.

Prednisone drug interactions (more detail)

Does prednisone interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

prednisone
+
Enter a drug name
Add

What is considered a high dose of prednisone?	~
Can you take prednisone with Keytruda?	~
What's a good prednisone taper schedule?	~
Is it OK to drink alcohol with Prednisone?	~
How long can you take prednisone safely?	~
Prednisone: What are 12 Things You Should Know?	~
How long does it take for prednisone to work?	~
Do I take all 6 prednisone pills at once on the first day? How long does prednisone stay in your system?	~
Tiom long does produited to stay in your system:	•

View more FAQ...

More about prednisone

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (867)
- Drug images
- Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: glucocorticoids
- Breastfeeding
- En español

Patient resources

• Prednisone Delayed-Release Tablets patient information

Other brands

Deltasone, Rayos, Sterapred, Prednicot, Sterapred DS

Professional resources

- Prednisone monograph
- Prednisone (FDA)
- Prednisone Tablets (FDA)

Other brands

Deltasone, Rayos

Related treatment guides

- Allergic Reactions
- Adrenocortical Insufficiency
- · Congenital Adrenal Hyperplasia
- Acute Lymphocytic Leukemia

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 9.01.

DRUG STATUS



User Reviews & Ratings

7.5 / 10

867 Reviews

Images

Prednisone 20 mg (060)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





