

Home

Benicar HCT

Benicar HCT

 $\textbf{Generic name:} \ \text{hydrochlorothiazide and olmesartan} \ [\ \textit{HYE-droe-KLOR-oh-THYE-a-zide-and-ol-me-SAR-tan}\]$

Drug class: Angiotensin II inhibitors with thiazides



Medically reviewed by Drugs.com on Feb 5, 2025. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Benicar HCT?

Hydrochlorothiazide is a diuretic (water pill). Olmesartan is an angiotensin II receptor blocker (sometimes called an ARB blocker).

Benicar HCT is a combination medicine used to treat high blood pressure (hypertension). Lowering blood pressure may lower your risk of a stroke or heart attack.

Benicar HCT is usually given after other blood pressure medicines did not work or have stopped working.

Benicar HCT may also be used for purposes not listed in this medication guide.

Benicar HCT side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

In rare cases, Benicar HCT can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine.

Also call your doctor at once if you have:

- a light-headed feeling, like you might pass out;
- · eye pain, vision problems;
- · an unusual skin rash;
- · easy bruising, unusual bleeding;
- jaundice (yellowing of the skin or eyes);
- **kidney problems**--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath;

- low levels of sodium in the body--headache, confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady;
- high potassium--slow or unusual heart rate, weakness, loss of movement; or
- low potassium--leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling.

Common side effects include:

- nausea;
- · dizziness:
- cold symptoms such as stuffy nose, sneezing, sore throat; or
- high levels of uric acid in your blood.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Benicar HCT side effects (more detail)

Related/similar drugs

Vasotec

Vasotec (enalapril) is used to treat high blood pressure, heart failure and kidney problems caused ...

Reviews & ratings

4 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Lisinopril

Lisinopril is an ACE inhibitor. It is used to treat high blood pressure, congestive heart failure ...

Reviews & ratings

- - / 4 -

View more

Atenolol

Atenolol is used to treat angina (chest pain) and hypertension (high blood pressure). Learn about ...

Reviews & ratings

7.2 / 10

276 Reviews

View more

Valsartan

Valsartan is an angiotensin II receptor blocker (ARB) that may be used to treat high blood pressure ...

Reviews & ratings

5.7 / 10

152 Reviews

View more

Hydrochlorothiazide

HCTZ (hydrochlorothiazide) used to treat high blood pressure (hypertension) and edema. Includes ...

Reviews & ratings

5.2 / 10

185 Reviews

View more

Lasix

Lasix is a loop diuretic used to treat fluid retention from heart, liver, or kidney conditions, and ...

Reviews & ratings

6.4 / 10

56 Reviews

View more

Furosemide

Furosemide is a loop diuretic used to treat fluid retention and high blood pressure by increasing ...

Reviews & ratings

6.8 / 10

121 Reviews

View more

Losartan

Losartan is used to treat high blood pressure (hypertension). It is also used to lower the risk of ...

Reviews & ratings

4.5 / 10

578 Reviews

View more

Warnings

You should not use this medicine if you are unable to urinate.

Do not use if you are pregnant. Use effective birth control, and tell your doctor if you become pregnant during treatment.

If you have diabetes, do not use Benicar HCT together with any medication that contains aliskiren (a blood pressure medicine).

Before taking this medicine

You should not use Benicar HCT if you are allergic to hydrochlorothiazide or olmesartan, or if you are unable to urinate.

If you have diabetes, do not use Benicar HCT together with any medication that contains aliskiren (a blood pressure medicine).

You may also need to avoid taking Benicar HCT with aliskiren if you have kidney disease.

Tell your doctor if you have ever had:

· kidney disease;

- · liver disease;
- · congestive heart failure;
- · glaucoma;
- low or high levels of potassium in your blood;
- asthma or allergies;
- high cholesterol or triglyceride levels;
- · gout;
- lupus;
- · diabetes; or
- an allergy to penicillin or sulfa drugs.

Do not use if you are pregnant. Stop using the medicine and tell your doctor right away if you become pregnant. Olmesartan can cause injury or death to the unborn baby if you take the medicine during your second or third trimester.

You should not breastfeed while using this medicine.

How should I take Benicar HCT?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Your blood pressure will need to be checked often.

Call your doctor if you have ongoing vomiting or diarrhea, or if you are sweating more than usual. You can easily become dehydrated while taking this medication, which can lead to severely low blood pressure or a serious electrolyte imbalance.

If you need surgery or medical tests, tell the surgeon ahead of time that you are using Benicar HCT.

It may take up to 4 weeks for this medication to control your blood pressure. **Keep using this medicine as directed, even if you feel well.** High blood pressure often has no symptoms. You may need to use blood pressure medication for the rest of your life.

Store at room temperature away from moisture and heat.

Detailed Benicar HCT dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

Overdose symptoms may include fast or slow heart rate, feeling light-headed, or fainting.

What should I avoid while taking Benicar HCT?

Hydrochlorothiazide may increase your risk of skin cancer. Avoid sunlight or tanning beds. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors. Your doctor may want you to have skin examinations on a regular basis.

Do not use potassium supplements or salt substitutes, unless your doctor has told you to.

Drinking alcohol can further lower your blood pressure and may cause side effects.

Avoid driving or hazardous activity until you know how Benicar HCT will affect you. Your reactions could be impaired. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy.

What other drugs will affect Benicar HCT?

If you also take cholestyramine or colestipol, take your Benicar HCT dose 4 hours before or 4 hours after you take the other medicine.

Tell your doctor about all your other medicines, especially:

- · steroids:
- any other blood pressure medications;
- · insulin or oral diabetes medicine;
- · lithium; or
- **NSAIDs (nonsteroidal anti-inflammatory drugs)**--aspirin, ibuprofen (Advil, Motrin), naproxen (Aleve), celecoxib, diclofenac, indomethacin, meloxicam, and others.

This list is not complete. Other drugs may affect Benicar HCT, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Benicar HCT drug interactions (more detail)

Does Benicar HCT interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



More about Benicar HCT (hydrochlorothiazide / olmesartan)

· Check interactions

- · Compare alternatives
- Pricing & coupons
- Reviews (29)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Drug class: angiotensin II inhibitors with thiazides
- En español

Professional resources

Benicar HCT prescribing information

Other formulations

Benicar

Related treatment guides

• High Blood Pressure

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation


& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

~



User Reviews & Ratings

6.9 / 10

29 Reviews

Images

Benicar HCT 12.5 mg / 20 mg (SANKYO C22)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com Advertising policy Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and









Copyright © 2000-2025 Drugs.com. All rights reserved.

 \times