

Home

2. Dexilant

Dexilant do

Generic name: dexlansoprazole [DEX-lan-SOE-pra-zol]

Drug class: Proton pump inhibitors



Medically reviewed by Drugs.com on Dec 26, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is Dexilant?

Dexilant is used to treat heartburn caused by gastroesophageal reflux disease (GERD), and to heal erosive esophagitis (damage to the esophagus from stomach acid).

Dexilant may also be used for purposes not listed in this medication guide.

Dexilant side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Dexilant may cause serious side effects. Call your doctor at once if you have:

- severe stomach pain, diarrhea that is watery or bloody;
- a seizure (convulsions);
- sudden pain or trouble moving your hip, wrist, or back;
- **kidney problems**-- fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain;
- low magnesium--dizziness, fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling; or
- new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight.

Taking Dexilant long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk.

If you use Dexilant for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it.

Common side effects of Dexilant may include:

- nausea, vomiting, stomach pain, gas;
- diarrhea;
- · headache;
- mouth pain, sore throat; or
- stuffy nose, sinus pain, or other cold symptoms.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Dexilant side effects (more detail)

Related/similar drugs

Protonix

Protonix (pantoprazole) is used to treat erosive esophagitis and other conditions involving excess ...

Reviews & ratings

7.2 / 10

83 Reviews

View more

Nexium

Nexium is used to treat GERD and other conditions involving excessive stomach acid. Learn about ...

Reviews & ratings

6.2 / 10

211 Reviews

View more

FEATURED

Repatha

Repatha (evolocumab) is a PCSK9 inhibitor used to lower high cholesterol alongside dietary changes ...

Reviews & ratings

5.3 / 10

685 Reviews

View more

Pepcid

Pepcid is used to treat ulcers in the stomach and for conditions where the stomach produces too ...

Reviews & ratings

6.0 / 10

10 Reviews

View more

Prilosec

Prilosec (omeprazole) is used to treat symptoms of GERD and other conditions caused by excess ...

Reviews & ratings

6.9 / 10

53 Reviews

View more

Prevacid

Prevacid is used to treat and prevent stomach and intestinal ulcers. Learn about side effects ...

Reviews & ratings

7.9 / 10

59 Reviews

View more

Aciphex

Aciphex is used to treat gastroesophageal reflux disease (GERD) and other conditions involving

...

Reviews & ratings

9.1 / 10

54 Reviews

View more

Ranitidine

Ranitidine is used to treat and prevent ulcers in the stomach and intestines. Includes ranitidine ...

Reviews & ratings

7.2 / 10

163 Reviews

View more

Warnings

Dexilant can cause kidney problems. Tell your doctor if you are urinating less than usual, or if you have blood in your urine.

Diarrhea may be a sign of a new infection. Call your doctor if you have diarrhea that is watery or has blood in it.

Dexilant may cause new or worsening symptoms of lupus. Tell your doctor if you have joint pain and a skin rash on your cheeks or arms that worsens in sunlight.

You may be more likely to have a broken bone while taking Dexilant long term or more than once per day.

Before taking this medicine

Heartburn can mimic early symptoms of a heart attack. **Get emergency medical help if you have** chest pain that spreads to your jaw or shoulder and you feel anxious or light-headed.

You should not take this medicine if you are allergic to Dexilant, or if:

- you had breathing problems, kidney problems, or a severe allergic reaction after taking this medicine in the past; or
- you also take a medicine that contains rilpivirine (Complera, Edurant, Odefsey).

Tell your doctor if you have ever had:

- severe liver disease;
- osteoporosis;
- low bone mineral density (osteopenia); or
- low levels of magnesium in your blood.

You may be more likely to have a broken bone in your hip, wrist, or spine while taking a proton pump inhibitor long-term or more than once per day. Talk with your doctor about ways to keep your bones healthy.

Dexilant may harm an unborn baby. Use effective birth control to prevent pregnancy, and tell your doctor if you become pregnant.

It may not be safe to breastfeed while using this medicine. Ask your doctor about any risk.

Dexilant is not approved for use by anyone younger than 12 years old.

How should I take Dexilant?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Take Dexilant with a full glass of water.

Dexilant may be taken with or without food.

Swallow the capsule whole and do not crush, chew, break, or open it.

If you cannot swallow a capsule whole, open it and sprinkle the medicine into a spoonful of applesauce. Swallow the mixture right away without chewing. Do not save it for later use.

To heal erosive esophagitis and relieve heartburn, Dexilant is usually given for up to 6 months in adults, and for 4 to 16 weeks in children ages 12 through 17. **Follow your doctor's dosing instructions very carefully.**

Use this medicine for the full prescribed length of time, even if your symptoms quickly improve.

Call your doctor if your symptoms do not improve, or if they get worse while using Dexilant.

This medicine can affect the results of certain medical tests. Tell any doctor who treats you that you are using Dexilant.

Dexilant may also affect a drug-screening urine test and you may have false results. Tell the laboratory staff that you use this medicine.

Store at room temperature away from moisture and heat.

1 Detailed Dexilant dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking Dexilant?

Dexilant can cause diarrhea, which may be a sign of a new infection. **If you have diarrhea that is watery or bloody, call your doctor.** Do not use anti-diarrhea medicine unless your doctor tells you to.

What other drugs will affect Dexilant?

Tell your doctor about all your current medicines. Many drugs can affect Dexilant, especially:

- methotrexate:
- rifampin;
- St. John's wort; or
- warfarin (Coumadin, Jantoven).

This list is not complete and many other drugs may affect Dexilant. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Dexilant drug interactions (more detail)

Does Dexilant interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



More about Dexilant (dexlansoprazole)

- · Check interactions
- · Compare alternatives
- Pricing & coupons
- Reviews (207)
- Drug images
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- Support group
- Drug class: proton pump inhibitors
- Breastfeeding
- En español

Patient resources

Other brands

Dexilant SoluTab, Kapidex

Professional resources

- · Dexilant prescribing information
- Dexlansoprazole (AHFS Monograph)

Related treatment guides

- · Barrett's Esophagus
- Erosive Esophagitis
- GERD

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 15.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

സ്യ Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

6.6 / 10

207 Reviews

Images

Dexilant 60 mg (TAP 60)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.