

Home

Kenalog

Kenalog ⊲

Generic name: triamcinolone topical [trye-am-SIN-oh-lone]

Brand names: Cinolar, Kenalog, Oralone, Pediaderm TA, Triamcinolone Acetonide in Absorbase, Trianex, Triderm

Drug class: Topical steroids

Medically reviewed by Sanjai Sinha, MD. Last updated on Jul 10, 2024.

Uses Warnings Before taking Dosage Side effects Interactions FAQ

What is Kenalog topical?

Kenalog contains triamcinolone, a potent corticosteroid. Triamcinolone reduces the actions of chemicals in the body that cause inflammation.

Kenalog topical is used to treat the inflammation caused by a number of conditions such as allergic reactions, eczema, and psoriasis. The dental paste form of triamcinolone (Kenalog Orabase) is used to treat mouth ulcers.

Kenalog topical may also be used for purposes not listed in this medication guide.

Warnings

Use Kenalog exactly as directed on the label, or as it has been prescribed by your doctor. Do not use the medication in larger amounts or for longer than recommended.

Do not cover treated skin areas with a bandage or other covering unless your doctor has told you to. If you are treating the diaper area of a baby, do not use plastic pants or tight-fitting diapers. Covering the skin that is treated with Kenalog can increase the amount of the drug your skin absorbs, which may lead to unwanted side effects. Follow your doctor's instructions.

Avoid using Kenalog on your face, near your eyes, or on body areas where you have skin folds or thin skin.

Do not use Kenalog on a child without a doctor's advice. Children are more sensitive to the effects of Kenalog.

Kenalog topical will not treat a bacterial, fungal, or viral skin infection.

Contact your doctor if your condition does not improve or if it gets worse after using this medication for several days.

Before using Kenalog

Do not use Kenalog if you are allergic to triamcinolone.

To make sure you can safely use Kenalog, tell your doctor if you have any of these other conditions:

- any skin infection, especially tuberculosis infection of the skin;
- chicken pox or herpes infection (including cold sores);
- · diabetes; or
- · a stomach ulcer.

It is not known whether Kenalog will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medication. It is not known whether triamcinolone passes into breast milk or if it could harm a nursing baby. Do not use Kenalog without telling your doctor if you are breastfeeding a baby. Do not use this medication on a child without a doctor's advice. Children are more sensitive to the effects of Kenalog.

Kenalog pregnancy and breastfeeding warnings (more detail)

How should I use Kenalog?

Use Kenalog exactly as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label.

Kenalog topical will not treat a bacterial, fungal, or viral skin infection.

Wash your hands before and after each application, unless you are using Kenalog to treat a hand condition.

Apply a small amount to the affected area and rub it gently into the skin.

Avoid using Kenalog on your face, near your eyes or mouth, or on body areas where you have skin folds or thin skin.

If you are using the dental paste, apply the medication in a thin layer, just enough to cover the mouth ulcer. The paste may stick better if you dry the mouth ulcer before applying the medication.

Do not cover treated skin areas with a bandage or other covering unless your doctor has told you to. If you are treating the diaper area of a baby, do not use plastic pants or tight-fitting diapers. Covering the skin that is treated with Kenalog can increase the amount of the drug your skin absorbs, which may lead to unwanted side effects. Follow your doctor's instructions. Contact your doctor if your condition does not improve or if it gets worse after using this medication for several days. It is important to use Kenalog regularly to get the most benefit.

Store at room temperature away from moisture and heat.

Detailed Kenalog dosage information

What happens if I miss a dose?

Use the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222, especially if anyone has accidentally swallowed this medicine. An overdose of Kenalog applied to the skin is not expected to produce life-threatening

symptoms.

What should I avoid?

Avoid getting Kenalog in your eyes, mouth, and nose, or on your lips. If it does get into any of these areas, wash with water. Do not use Kenalog on sunburned, windburned, irritated, or broken skin. Also avoid using Kenalog in open wounds.

Avoid using skin products that can cause irritation, such as harsh soaps or skin cleansers, or skin products with alcohol, spices, astringents, or lime. Avoid using other medications on the areas you treat with Kenalog unless your doctor tells you to.

Kenalog side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Kenalog: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using this medication and call your doctor at once if you have any of these signs that you may be absorbing triamcinolone through your skin or gums:

- blurred vision, or seeing halos around lights;
- · uneven heartbeats;
- · mood changes;
- sleep problems (insomnia);
- · weight gain, puffiness in your face; or
- · feeling tired.

Less serious Kenalog side effects may include:

- skin redness, burning, itching, or peeling;
- · thinning of your skin; or
- · blistering skin; or
- · stretch marks.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Kenalog side effects (more detail)

Related/similar drugs

Dupixent

Dupixent may be used to treat eczema, eosinophilic or oralcorticosteroid-dependent asthma, chronic ...

Reviews & ratings

7.2 / 10

499 Reviews

View more

Stelara

Stelara (ustekinumab) is used to treat Crohn's disease, ulcerative colitis, plaque psoriasis, and ...

Reviews & ratings

7.6 / 10

252 Reviews

View more

FEATURED

Skyrizi

Skyrizi (risankizumab) is used to treat plaque psoriasis, psoriatic arthritis, ulcerative colitis ...

Reviews & ratings

7.9 / 10

56 Reviews

View more

Rinvoq

Rinvoq (upadacitinib) is used to treat rheumatoid arthritis, psoriatic arthritis, atopic ...

Reviews & ratings

7.0 / 10

223 Reviews

View more

Humira

Humira is a tumor necrosis factor blocker that is used to treat many inflammatory conditions such ...

Reviews & ratings

View more

Carafate

Carafate is an anti-ulcer medication used to treat active duodenal ulcers. Learn about side ...

Reviews & ratings

6.2 / 10

72 Reviews

View more

Remicade

Remicade is used to treat rheumatoid arthritis, psoriatic arthritis, ulcerative colitis and Crohn's ...

Reviews & ratings

7.0 / 10

213 Reviews

View more

Adalimumab

Adalimumab is used to treat plaque psoriasis, Crohn's disease, ulcerative colitis, rheumatoid ...

Reviews & ratings

6.4 / 10

737 Reviews

View more

Clobetasol topical

Clobetasol is available as a cream, gel, ointment, lotion, solution, foam or spray and is used to ...

Reviews & ratings

8.2 / 10

164 Reviews

What other drugs will affect Kenalog?

It is not likely that other drugs you take orally or inject will have an effect on topically applied Kenalog. But many drugs can interact with each other. Tell your doctor about all medications you use. This includes prescription, over-the-counter, vitamin, and herbal products. Do not start a new medication without telling your doctor.

Kenalog drug interactions (more detail)

Does Kenalog interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



Popular FAQ

Where should you not use triamcinolone acetonide cream?

More about Kenalog (triamcinolone topical)

- · Check interactions
- · Compare alternatives
- Reviews (31)
- Side effects
- Dosage information
- During pregnancy
- Drug class: topical steroids
- Breastfeeding
- En español

Patient resources

- · Kenalog (Triamcinolone Cream and Ointment) patient information
- Kenalog (Triamcinolone Lotion)
- Kenalog (Triamcinolone Topical Aerosol Solution)

Other brands

Triderm, Trianex, Aristocort A, Triacet, ... +3 more

Professional resources

- Triamcinolone Acetonide (Topical) monograph
- Triamcinolone in Absorbase (FDA)

Other brands

Triderm, Trianex, DermacinRx Trizapak, Dermasorb TA, ... +7 more

Other formulations

- Kenalog-40
- Kenalog-10

Related treatment guides

- Atopic Dermatitis
- Lichen Planus
- Eczema
- Psoriasis
- Dermatitis
- Stomatitis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Kenalog only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 8.02.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ჟ^ა Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

WADA Class

Anti-Doping Classification

User Reviews & Ratings

8.3 / 10

31 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







