

Home

2. Imipramine

Imipramine ⊲

Generic name: imipramine [im-IP-ra-meen]

Brand names: Tofranil, Tofranil-PM

Dosage forms: oral capsule (pamoate 100 mg; pamoate 125 mg; pamoate 150 mg; pamoate 75 mg), oral tablet (10 mg; 25 mg; 50 mg)

Drug class: Tricyclic antidepressants



Medically reviewed by Drugs.com on Aug 3, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is imipramine?

Imipramine is a tricyclic antidepressant used to treat symptoms of depression. Imipramine is sometimes used to treat bed-wetting in children ages 6 and older.

Imipramine may also be used for purposes not listed in this medication guide.

Imipramine side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself.

Imipramine may cause serious side effects. Call your doctor at once if you have:

- easy bruising, unusual bleeding, purple or red spots under your skin;
- tunnel vision, eye pain or swelling, or seeing halos around lights;
- a light-headed feeling, like you might pass out;
- new or worsening chest pain, pounding heartbeats or fluttering in your chest;
- sudden numbness or weakness, problems with vision, speech, or balance;
- fever, sore throat;
- · confusion, hallucinations, unusual thoughts or behavior;
- painful or difficult urination;
- seizure (convulsions); or

• jaundice (yellowing of the skin or eyes).

Common side effects of imipramine may include:

- increased blood pressure;
- tingly feeling, weakness, lack of coordination;
- dry mouth, nausea, vomiting; or
- breast swelling (in men or women).

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

i <u>Imipramine side effects</u> (more detail)

Related/similar drugs

Tylenol

Tylenol is a pain reliever and a fever reducer used to treat many conditions such as headaches ...

Reviews & ratings

5.8 / 10

30 Reviews

View more

Oxycodone

Oxycodone is an opioid analgesic used to treat moderate to severe pain; it has a high potential for ...

Reviews & ratings

6.9 / 10

1,142 Reviews

View more

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Lexapro

Lexapro is used to treat anxiety and major depressive disorder. Learn about side effects ...

Reviews & ratings

7.6 / 10

2,376 Reviews

View more

Naproxen

Naproxen is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.8 / 10

683 Reviews

View more

Paracetamol

Paracetamol (Panadol, Calpol, Alvedon) is a widely used over-thecounter painkiller and fever ...

Reviews & ratings

6.1 / 10

13 Reviews

View more

Hydroxyzine

Hydroxyzine is an antihistamine used to treat itching, hives, and anxiety. It also acts as a ...

Reviews & ratings

5.9 / 10

1,404 Reviews

View more

ZUIUIL

Zoloft is an antidepressant used to treat major depression, obsessivecompulsive disorder, panic ...

Reviews & ratings

7.3 / 10

1,972 Reviews

View more

Meloxicam

Meloxicam is a nonsteroidal antiinflammatory drug used to treat pain or inflammation caused by ...

Reviews & ratings

6.3 / 10

600 Reviews

View more

Cyclobenzaprine

Cyclobenzaprine is a muscle relaxant and works by blocking pain sensations. Includes ...

Reviews & ratings

6.0 / 10

618 Reviews

View more

Warnings

Some young people have thoughts about suicide when first taking an antidepressant. Stay alert to changes in your mood or symptoms. **Report any new or worsening symptoms to your doctor**.

Before taking this medicine

You should not use imipramine if you are allergic to it, or if:

- you recently had a heart attack; or
- you are allergic to other antidepressants (amitriptyline, amoxapine, clomipramine, desipramine, doxepin, nortriptyline, protriptyline, or trimipramine).

Do not use imipramine if you have used an MAO inhibitor in the past 14 days. A dangerous drug interaction could

occur. MAO inhibitors include isocarboxazid, linezolid, methylene blue injection, phenelzine, tranylcypromine, and others.

Tell your doctor if you have used an "SSRI" antidepressant in the past 5 weeks, such as citalopram, escitalopram, fluoxetine (Prozac), fluoxamine, paroxetine, sertraline (Zoloft), trazodone, or vilazodone.

Tell your doctor if you have ever had:

- heart disease, stroke, or seizures;
- bipolar disorder (manic-depression), schizophrenia or other mental illness;
- kidney or liver disease;
- · overactive thyroid;
- narrow-angle glaucoma; or
- problems with urination.

Some young people have thoughts about suicide when first taking an antidepressant. Stay alert to changes in your mood or symptoms. Report any new or worsening symptoms to your doctor.

It is not known whether imipramine will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant.

You should not breastfeed while using imipramine.

Do not give this medicine to a child without medical advice. Imipramine is not approved to treat depression in anyone younger than 18 years old. Imipramine should not be used to treat bed-wetting in a child younger than 6 years old.

How should I take imipramine?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Tell your doctor if you have a planned surgery.

Call your doctor if your symptoms do not improve, or if they get worse.

You should not stop using imipramine suddenly. Follow your doctor's instructions about tapering your dose.

Store at room temperature away from moisture, heat, and light. Keep the bottle tightly closed when not in use.

Detailed Imipramine dosage information

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of imipramine can be fatal.**

Overdose symptoms may include irregular heartbeats, feeling like you might pass out, seizures, or coma.

What should I avoid while taking imipramine?

Do not drink alcohol. Dangerous side effects or death can occur when alcohol is combined with imipramine.

Avoid driving or hazardous activity until you know how imipramine will affect you. Your reactions could be impaired.

Avoid exposure to sunlight or tanning beds. Imipramine can make you sunburn more easily. Wear protective clothing and use sunscreen (SPF 30 or higher) when you are outdoors.

What other drugs will affect imipramine?

Taking this medicine with other drugs that make you sleepy can worsen this effect. Ask your doctor before taking imipramine with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

Many drugs can affect imipramine. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here. Tell your doctor about all other medicines you use.

1 Imipramine drug interactions (more detail)

Does imipramine interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

imipramine	
+	
Enter a drug name	Add

More about imipramine

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (79)
- Drug images
- · Side effects
- Dosage information
- During pregnancy

- Drug class: tricyclic antidepressants
- Breastfeeding
- En español

Patient resources

- · Imipramine advanced reading
- Imipramine Tablets
- Imipramine Capsules

Other brands

Tofranil, Tofranil-PM

Professional resources

- Imipramine monograph
- Imipramine (FDA)
- Imipramine Pamoate (FDA)

Other brands

Tofranil

Related treatment guides

- Interstitial Cystitis
- Primary Nocturnal Enuresis
- Enuresis
- Pain

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 14.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

7.3 / 10

79 Reviews

Images

Imipramine Hydrochloride 10 mg (par 54)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.







Copyright © 2000-2025 Drugs.com. All rights reserved.

 \times