

Home

2. GlipiZIDE XL

GlipiZIDE XL 🖘

Generic name: glipizide [*GLIP-i-zide*] **Brand names:** GlipiZIDE XL, Glucotrol XL

Drug class: Sulfonylureas



Medically reviewed by Drugs.com on May 6, 2024. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is GlipiZIDE XL?

GlipiZIDE XL can be used alone or with other medications, diet, and exercise to improve blood sugar control in adults with type 2 diabetes mellitus (not for type 1 diabetes).

GlipiZIDE XL may also be used for purposes not listed in this medication guide.

GlipiZIDE XL side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives, difficult breathing, swelling of your face, lips, tongue, or throat.

Call your doctor at once or seek emergency medical attention if you have:

- **liver problems**--loss of appetite, nausea, vomiting, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes);
- signs of low or severely low blood sugar--headache, hunger, sweating, confusion, irritability, dizziness, fast heart rate, feeling jittery, extreme weakness, nausea, tremors, trouble speaking, or seizure; or
- signs of hemolytic anemia--pale or yellowed skin, dark colored urine, fever, confusion or weakness.

Common side effects of GlipiZIDE XL may include:

- diarrhea, nausea, gas;
- dizziness, drowsiness;
- · headache, tremors, feeling nervous; or
- · skin rash, redness, or itching.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Related/similar drugs

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Rybelsus

Rybelsus tablets are used to improve blood sugar control in adults with type 2 diabetes, and may ...

Reviews & ratings

6.4 / 10

282 Reviews

View more

FEATURED

Mounjaro

Mounjaro (tirzepatide) is FDAapproved to manage type 2 diabetes in adults. It lowers blood sugar ...

Reviews & ratings

8.5 / 10

1,278 Reviews

View more

Trulicity

Trulicity is an injectable diabetes medicine that is used together with diet and exercise to ...

Reviews & ratings

5.3 / 10

970 Reviews

View more

Lantus

Lantus is a long acting form of insulin used to treat type 1 or type 2 diabetes. Learn about side ...

Reviews & ratings

6.5 / 10

41 Reviews

View more

Tresiba

Tresiba (insulin degludec) is used to treat diabetes mellitus. Includes Tresiba side effects ...

Reviews & ratings

4.3 / 10

152 Reviews

View more

Victoza

Victoza helps control blood sugar levels and reduce the risk of serious heart problems in people ...

Reviews & ratings

7.6 / 10

766 Reviews

View more

Basaglar

Basaglar (insulin glargine) is a longacting insulin that is used to improve blood sugar control in ...

Reviews & ratings

2.7 / 10

148 Reviews

View more

Levemir

Levemir (insulin detemir) is a long acting insulin used to treat diabetes in adults and children ...

Reviews & ratings

4.6 / 10

70 Reviews

View more

Novolog

NovoLog is a fast-acting insulin used to treat diabetes to control blood sugar levels in adults and ...

Reviews & ratings

5.5 / 10

32 Reviews

View more

Warnings

Use only as directed. Tell your doctor if you use other medicines or have other medical conditions or allergies.

Before taking this medicine

You should not use GlipiZIDE XL XL if you are allergic to GlipiZIDE XL or any of its ingredients or if you have:

- type 1 diabetes;
- a condition called diabetic ketoacidosis; or
- an allergy to sulfa drugs.

Tell your doctor if you have or have ever had:

- · diabetic ketoacidosis;
- a blockage in your stomach or intestines caused by disease or surgery;
- · severe or ongoing diarrhea;
- an enzyme deficiency called glucose-6-phosphate dehydrogenase deficiency (G6PD); or
- · liver or kidney disease.

GlipiZIDE XL may increase your risk of heart problems. Ask your doctor about this risk.

It is not known if GlipiZIDE XL will harm an unborn baby. Tell your doctor if you are pregnant or plan to become

pregnant. Follow your doctor's instructions about using this medicine if you are pregnant.

Having high blood sugar during pregnancy may cause complications or medical problems in both mother and baby. Controlling diabetes is very important during pregnancy, and you may need to stop taking GlipiZIDE XL for a short time just before your due date.

Ask a doctor if it is safe to breastfeed while using this medicine.

How should I take GlipiZIDE XL?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Your dose needs may change if you switch to a different brand, strength, or form of this medicine. **Avoid medication** errors by using exactly as directed on the label, or as prescribed by your doctor.

Take GlipiZIDE XL 30 minutes before a meal.

Take the GlipiZIDE XL extended-release tablet once a day with breakfast or your first meal of the day.

Swallow the extended-release tablet whole and do not crush, break, chew or dissolve it.

Your blood sugar will need to be checked on a regular basis, and you may need other medical tests.

Part of the extended-release tablet shell may appear in your stool but this will not make the medicine less effective.

Blood sugar can be affected by stress, illness, surgery, exercise, alcohol use, or skipping meals.

Low blood sugar (hypoglycemia) can make you feel very hungry, dizzy, irritable, or shaky. To quickly treat hypoglycemia, eat or drink hard candy, crackers, raisins, fruit juice, or non-diet soda. Your doctor may prescribe glucagon injection in case of severe hypoglycemia.

Tell your doctor if you have frequent symptoms of high blood sugar (hyperglycemia) such as increased thirst or urination. Ask your doctor before changing your medication dosage.

Your treatment may also include diet, exercise, weight control, and special medical care.

Store tightly closed at room temperature in the original container away from moisture and heat.

Detailed GlipiZIDE XL dosage information

What happens if I miss a dose?

Take your dose as soon as you can, **but only if you are getting ready to eat a meal.** If you skip a meal, skip the missed dose and wait until your next meal. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. An overdose can be fatal.

What should I avoid while taking GlipiZIDE XL?

Do not drink alcohol. Dangerous side effects could occur.

Avoid driving or hazardous activity until you know how GlipiZIDE XL XL will affect you. Your reactions could be impaired.

What other drugs will affect GlipiZIDE XL?

When you start or stop taking GlipiZIDE XL, your doctor may need to adjust the doses of other medicines you use on a regular basis.

Take your GlipiZIDE XL dose 4 hours before taking colesevelam.

Tell your doctor about all your other medicines, especially miconazole or fluconazole.

Many other drugs can affect your blood sugar and may also affect GlipiZIDE XL. This includes prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible interactions are listed here.

GlipiZIDE XL drug interactions (more detail)

Does GlipiZIDE XL interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Enter a drug name Add

More about GlipiZIDE XL (glipizide)

- · Check interactions
- Compare alternatives
- Reviews (10)
- Drug images
- Side effects
- Dosage information
- During pregnancy
- Drug class: sulfonylureas
- Breastfeeding
- En español

Patient resources

Other brands

Glucotrol, Glucotrol XL

Professional resources

- Glipizide XL prescribing information
- Glipizide (AHFS Monograph)

Other brands

Glucotrol, Glucotrol XL

Related treatment guides

Type 2 Diabetes

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 15.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

প্% Risk data available

CSA Schedule*

N/A Not a controlled drug

User Reviews & Ratings

5.8 / 10

10 Reviews

Images

Glipizide XL 10 mg (GLIPIZIDE XL 10)



Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com
Advertising policy
Attribution & citations

Terms & privacy

Terms of use Editorial policy Privacy policy

Support

Help center Sitemap Contact us











שא

Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.