

Home

Nasacort Allergy 24HR

Nasacort Allergy 24HR Nasal Spray

Generic name: triamcinolone (nasal) [TRYE-am-SIN-oh-lone]

Brand name: Nasacort Allergy 24HR

Drug class: Nasal steroids

Medically reviewed by Philip Thornton, DipPharm. Last updated on Aug 23, 2023.

Uses Warnings Side effects Before taking Dosage Interactions

What is Nasacort Allergy 24HR?

Nasacort Allergy 24HR Nasal Spray contains triamcinolone, a corticosteroid medicine that prevents the release of substances in the body that cause inflammation.

Nasacort Allergy 24HR is used to treat sneezing, itching, and runny or stuffy nose caused by seasonal allergies or hay fever.

Nasacort Allergy 24HR may also be used for purposes not listed in this medication guide.

Warnings

Nasacort Allergy 24HR should not be given to a child younger than 2 years old.

Follow all directions on your medicine label and package. Tell each of your healthcare providers about all your medical conditions, allergies, and all medicines you use.

Nasacort side effects

Get emergency medical help if you have **signs of an allergic reaction to Nasacort:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- fever, chills, body aches, flu symptoms;
- · nosebleeds; or
- blurred vision, tunnel vision, eye pain, or seeing halos around lights.

Triamcinolone can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine.

Less serious Nasacort side effects may be more likely, and you may have none at all.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Nasacort Allergy 24HR side effects (more detail)

Related/similar drugs

Promethazine

Promethazine is a phenothiazine and antihistamine used to treat allergies, motion sickness, nausea ...

Reviews & ratings

7.2 / 10

388 Reviews

View more

Loratadine

Loratadine is a non-drowsy antihistamine used to relieve allergy symptoms like sneezing, runny ...

Reviews & ratings

4.7 / 10

139 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Methylprednisolone

Methylprednisolone is used to treat conditions such as allergies, arthritis, lupus and ulcerative ...

Reviews & ratings

7.1 / 10

484 Reviews

View more

Cetirizine

Cetirizine treats cold or allergy symptoms such as sneezing, itching, watery eyes, or runny nose ...

Reviews & ratings

5.8 / 10

294 Reviews

View more

Montelukast

Montelukast is a daily oral medication used to prevent asthma attacks, exercise-induced ...

Reviews & ratings

5.6 / 10

571 Reviews

View more

Zyrtec

Zyrtec (cetirizine) is used to treat allergy symptoms such as sneezing, itching, watery eyes, or ...

Reviews & ratings

5.8 / 10

197 Reviews

....

View more

Xolair

Xolair injection (omalizumab) is used to reduce the risk of severe food allergy reactions ...

Reviews & ratings

7.5 / 10

229 Reviews

View more

Before taking this medicine

You should not use Nasacort Allergy 24HR if you are allergic to triamcinolone.

Ask a doctor or pharmacist if this medicine is safe to use if you have ever had:

- an eye infection;
- glaucoma or cataracts;
- ulcers in your nose; or
- surgery or injury to your nose.

Ask a doctor before using this medicine if you are pregnant or breastfeeding.

Nasacort Allergy 24HR should not be given to a child younger than 2 years old.

Nasacort Allergy 24HR pregnancy and breastfeeding warnings (more detail)

How should I use Nasacort Allergy 24HR?

Use Nasacort exactly as directed on the label, or as prescribed by your doctor.

Do not take by mouth. Nasacort Allergy 24HR is for use only in your nose.

Read and carefully follow any Instructions for Use provided with your medicine. Ask your doctor or pharmacist if you do not understand these instructions.

Before your first use, prime the nasal spray pump by spraying the medicine into the air until a fine mist appears. If the nasal spray has not been used for longer than 2 weeks, prime it by spraying the medicine into the air until a fine mist appears.

Shake the Nasacort nasal spray bottle well just before each use.

To use the nasal spray:

- Blow your nose gently. Keep your head upright and insert the tip of the bottle into one nostril. Press your other nostril
 closed with your finger. Breathe in quickly and gently spray the medicine into your nose. Then use the spray in your
 other nostril.
- The usual dose is 1 to 2 sprays per nostril once daily.
- Do not blow your nose for at least a few minutes after using the nasal spray.
- Do not use the nasal spray more than one time in 24 hours.
- Keep the nozzle of the nasal spray bottle clean, following the cleaning instructions provided with your medicine.
- If the nasal spray has not been used for longer than 2 weeks, prime it by spraying 1 or 2 times into a tissue.

Stop using Nasacort Allergy 24HR and call your doctor if your symptoms do not improve after 7 days of treatment, or if you have a fever with a headache, cough, or skin rash.

Do not share the nasal spray with another person, or you could spread germs or disease.

Store at room temperature away from moisture and heat. Keep the bottle tightly closed when not in use.

Dosing information

Usual Adult Dose of Nasacort Nasal Spray for Allergic Rhinitis:

Nasal aerosol:

2 actuations in each nostril once a day; may increase after 4 to 7 days up to 4 actuations in each nostril once a day, or 2 actuations in each nostril twice daily, or 1 actuation in each nostril 4 times a day.

Nasal spray:

2 sprays in each nostril once a day; titrate to lowest effective dose after symptoms are controlled. Usual maintenance dose is 1 spray in each nostril once a day.

Usual Pediatric Dose of Nasacort Nasal Spray for Allergic Rhinitis:

Nasal spray:

2 years to 5 years: 1 spray in each nostril once a day;

6 years to 11 years: 1 spray in each nostril once a day; maximum dose is 2 sprays in each nostril once a day, titrate to lowest effective dose after symptoms are controlled.

12 years or older: 2 sprays in each nostril once a day; titrate to lowest effective dose after symptoms are controlled. Usual maintenance dose is 1 spray in each nostril once a day.

Detailed Nasacort Allergy 24HR dosage information

What happens if I miss a dose?

Use the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. Do not use two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

An overdose of Nasacort Allergy 24HR is not expected to produce life threatening symptoms. However, long term use of high steroid doses can lead to symptoms such as thinning skin, easy bruising, changes in the shape or location of body fat (especially in your face, neck, back, and waist), increased acne or facial hair, menstrual problems, impotence, or loss of interest in sex.

What should I avoid while using triamcinolone nasal?

Avoid getting Nasacort Allergy 24HR in your eyes.

Avoid being near people who are sick or have infections. Call your doctor for preventive treatment if you are exposed to chickenpox or measles. These conditions can be serious or even fatal in people who are using triamcinolone.

What other drugs will affect Nasacort Allergy 24HR?

Ask a doctor or pharmacist before using Nasacort Allergy 24HR with any other medications, especially:

• steroid medicine for asthma, allergies or skin rash.

This list is not complete. Other drugs may interact with triamcinolone, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

Nasacort Allergy 24HR drug interactions (more detail)

Does Nasacort Allergy 24HR interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Nasacort Allergy 24HR	
+	
Enter a drug name	Add

References

1. Nasacort Product Label

More about Nasacort Allergy 24HR (triamcinolone nasal)

- · Check interactions
- Compare alternatives
- · Pricing & coupons
- Reviews (14)
- · Side effects
- Dosage information

- During pregnancy
- Drug class: nasal steroids
- Breastfeeding
- En español

Patient resources

Nasacort Allergy 24HR patient information

Other brands

Nasacort AQ, Allernaze

Professional resources

- Nasacort Allergy 24HR prescribing information
- Triamcinolone Acetonide (EENT) (AHFS Monograph)

Other brands

Nasacort, Nasacort AQ

Related treatment guides

Allergic Rhinitis

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Nasacort nasal spray only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 10.03.

DRUG STATUS

Availability

OTC Over the counter

Pregnancy & Lactation

্যু Risk data available

CSA Schedule*

N/A Not a controlled drug

WADA Class

്റ്റ് Anti-Doping Classification

User Reviews & Ratings

7.1 / 10

14 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.





