



[Home](#)

2. [Macrobid](#)

Macrobid

Generic name: [nitrofurantoin](#) [*NYE-troe-fue-RAN-toin*]

Drug class: [Urinary anti-infectives](#)

Medically reviewed by [Sanjai Sinha, MD](#). Last updated on Aug 26, 2024.

[Uses](#) [Warnings](#) [Before taking](#) [Side effects](#) [Dosage](#) [Interactions](#) [FAQ](#)

What is Macrobid?

Macrobid is an antibiotic that fights bacteria in the body.

Macrobid is used to treat [urinary tract infections](#).

Macrobid may also be used for purposes not listed in this medication guide.

Warnings

You should **not** take Macrobid if you have severe kidney disease, **complicated** urinary tract infection, or a history of jaundice or liver problems caused by nitrofurantoin.

Do not take Macrobid if you are in the last 2 to 4 weeks of pregnancy.

Before taking this medicine

You should not take Macrobid if you are allergic to nitrofurantoin, or if you have:

- severe kidney disease;
- a history of jaundice or liver problems caused by taking nitrofurantoin;
- if you are urinating less than usual or not at all; or
- if you are in the last 2 to 4 weeks of pregnancy.

Do not take Macrobid if you are in the last 2 to 4 weeks of pregnancy.

To make sure Macrobid is safe for you, tell your doctor if you have:

- kidney disease;
- anemia;

- diabetes;
- an electrolyte imbalance or vitamin B deficiency;
- glucose-6-phosphate dehydrogenase (G6PD) deficiency; or
- any type of debilitating disease.

FDA pregnancy category B. Macrobid is not expected to be harmful to an unborn baby during early pregnancy. Tell your doctor if you are pregnant or plan to become pregnant during treatment.

Nitrofurantoin can pass into breast milk and may harm a nursing baby. You should not breast-feed while you are taking Macrobid.

Macrobid should not be given to a child younger than 1 month old.

 [Macrobid pregnancy and breastfeeding warnings](#) (more detail)

How should I take Macrobid?

Take Macrobid exactly as prescribed by your doctor. Follow all directions on your prescription label. Do not take this medicine in larger or smaller amounts or for longer than recommended.

Take Macrobid with food.

Shake the oral suspension (liquid) well just before you measure a dose. Measure liquid medicine with the dosing syringe provided, or with a special dose-measuring spoon or medicine cup. If you do not have a dose-measuring device, ask your pharmacist for one.

You may mix your liquid dose with water, milk, or fruit juice to make it easier to swallow. Drink the entire mixture right away.

Use this medicine for the full prescribed length of time. Your symptoms may improve before the infection is completely cleared. Skipping doses may also increase your risk of further infection that is resistant to antibiotics. Macrobid will not treat a viral infection such as the common cold or flu.

Macrobid is usually given for 5 days in females and 7 days in males for uncomplicated UTI.

If you use this medicine long-term, for prevention of UTI, you may need frequent medical tests at your doctor's office.

Macrobid can cause unusual results with certain lab tests for glucose (sugar) in the urine. Tell any doctor who treats you that you are using Macrobid.

Store at room temperature away from moisture, heat, and light.

What happens if I miss a dose?

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What to avoid

Antibiotic medicines can cause diarrhea, which may be a sign of a new infection. If you have diarrhea that is watery or has blood in it, call your doctor. Do not use any medicine to stop the diarrhea unless your doctor has told you to.

Avoid using antacids without your doctor's advice. Use only the type of antacid your doctor recommends. Some antacids can make it harder for your body to absorb nitrofurantoin.

Macrobid side effects

Get emergency medical help if you have any of these **signs of an allergic reaction to Macrobid**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Call your doctor at once if you have:

- diarrhea that is watery or bloody;
- sudden chest pain or discomfort, wheezing, dry cough or hack;
- new or worsening cough, trouble breathing;
- fever, chills, body aches, tiredness, unexplained weight loss;
- numbness, tingling, or pain in your hands or feet;
- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes); or
- lupus-like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color.

Serious side effects may be more likely in older adults and those who are ill or debilitated.

Common Macrobid side effects may include:

- headache, dizziness;
- gas, upset stomach;
- mild diarrhea; or
- vaginal itching or discharge.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Macrobid side effects](#) (more detail)

Related/similar drugs

[Amoxicillin/clavulanate](#)

Amoxicillin and clavulanate potassium is a combination antibiotic used to treat bacterial ...

Reviews & ratings

5.3 / 10

[910 Reviews](#)

[View more](#)

Levofloxacin

Levofloxacin is a fluoroquinolone antibiotic used to treat serious bacterial infections and prevent ...

Reviews & ratings

5.1 / 10

[741 Reviews](#)

[View more](#)

FEATURED

Omvox

Omvox is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

[1 Reviews](#)

[View more](#)

Augmentin

Augmentin is a prescription antibiotic combining amoxicillin and clavulanate to treat bacterial ...

Reviews & ratings

5.4 / 10

[447 Reviews](#)

[View more](#)

Ciprofloxacin

Ciprofloxacin is an antibiotic belong to a group of drugs called fluoroquinolones. Learn about side ...

Reviews & ratings

5.0 / 10

[935 Reviews](#)

[View more](#)

Cephalexin

Cephalexin is a cephalosporin antibiotic used to treat bacterial infections like respiratory, skin ...

Reviews & ratings

6.4 / 10

[509 Reviews](#)

[View more](#)

Bactrim

Bactrim (sulfamethoxazole and trimethoprim) is an antibiotic used to treat ear infections, urinary ...

Reviews & ratings

5.4 / 10

[616 Reviews](#)

[View more](#)

Keflex

Keflex (cephalexin) is used to treat infections caused by bacteria, including respiratory ...

Reviews & ratings

7.3 / 10

[91 Reviews](#)

[View more](#)

Dosing information

Usual Adult Dose for Cystitis:

Regular release: 50 to 100 mg orally 4 times a day for 1 week or for at least 3 days after urine sterility is obtained

Dual release: 100 mg orally twice a day for 7 days

Usual Adult Dose for Cystitis Prophylaxis:

Regular release: 50 to 100 mg orally once a day at bedtime

Usual Pediatric Dose for Cystitis:

Regular release:

1 month or older: 5 to 7 mg/kg/day (up to 400 mg/day) orally in 4 divided doses

Dual release:

Greater than 12 years: 100 mg orally twice a day for 7 days

Usual Pediatric Dose for Cystitis Prophylaxis:

Regular release:

1 month or older: 1 to 2 mg/kg/day (up to 100 mg/day) orally in 1 to 2 divided doses

 [Detailed Macrobid dosage information](#)

What other drugs will affect Macrobid?

Other drugs may interact with nitrofurantoin, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

 [Macrobid drug interactions](#) (more detail)

Does Macrobid interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

Macrobid

+

Enter a drug name

Add

Frequently asked questions

- [What is "dual release" nitrofurantoin?](#)
- [Is meloxicam a sulfa drug?](#)
- [Does Macrobid make birth control pills less effective?](#)
- [What antibiotics are used to treat UTI?](#)

[View more FAQ...](#)

More about Macrobid (nitrofurantoin)

- [Check interactions](#)

- [Compare alternatives](#)
- [Pricing & coupons](#)
- [Reviews \(827\)](#)
- [Drug images](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Generic availability](#)
- [Support group](#)
- [Drug class: urinary anti-infectives](#)
- [Breastfeeding](#)
- [En español](#)

Patient resources

Other brands

[Furadantin](#)

Professional resources

- [Macrobid prescribing information](#)
- [Nitrofurantoin \(AHFS Monograph\)](#)

Other brands

[Macrochantin](#), [Furadantin](#)

Related treatment guides

- [Urinary Tract Infection](#)
- [Prevention of Bladder infection](#)
- [Bladder Infection](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use Macrobid only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Antibiotics 101

[List of Common Antibiotics & Types](#)

[Antibiotics and Alcohol](#)

[Probiotics and Antibiotics](#)

[Antibiotics for Strep Throat](#)

[Antibiotics for a Tooth Infection](#)


[Antibiotics for UTI's](#)

DRUG STATUS

Availability

Rx Prescription only


Pregnancy & Lactation

 Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

 Drug history at FDA



User Reviews & Ratings

4.5 / 10

[827 Reviews](#)

Images

[Macrobid 100 mg \(Macrobid Norwich Eaton\)](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)


[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Ad Choices
ADVERTISEMENT



Copyright © 2000-2025 Drugs.com. All rights reserved.