

Home

Veltin

Veltin

Generic name: clindamycin and tretinoin (topical) [klin-da-MYE-sin-and-TRET-i-noyn-TOP-i-kal]

Brand names: Veltin, Ziana

Drug class: Topical acne agents

Medically reviewed by Drugs.com. Last updated on Jul 1, 2024.

Uses Warnings Before taking Dosage Side effects Interactions

What is Veltin?

Veltin is a gel containing a combination of clindamycin and tretinoin. Clindamycin is an antibiotic. It fights bacteria in the body.

Tretinoin is a form of vitamin A. It reduces the amount of oil released by oil glands in your skin, and helps your skin renew itself more quickly.

Veltin gel is a precription medicine used on the skin to treat acne vulgaris in adults and children 12 years and older.

Veltin may also be used for purposes other than those listed in this medication guide.

Warnings

Avoid getting Veltin in your eyes, mouth, or nose (or in the creases of your nose), or on your lips. If it does get into any of these areas, wash with water. Do not apply Veltin to sunburned, windburned, dry, chapped, irritated, or broken skin.

You should not use this medication if you have ulcerative colitis, Crohn's disease, inflammation of your intestines (also called enteritis), or if you have ever had severe diarrhea caused by antibiotic medicine.

You should not use Veltin if you are also taking another antibiotic called erythromycin, or using other skin medications that contain erythromycin.

Although Veltin gel is applied to the skin, your body may absorb enough clindamycin to cause serious side effects.

Avoid exposure to sunlight or artificial UV rays (sunlamps or tanning beds). Veltin gel can make your skin more sensitive to sunlight and sunburn may result. Use a sunscreen (minimum SPF 15) and wear protective clothing if you must be out in the sun.

Your skin may be more sensitive to weather extremes such as cold and wind. Protect your skin with clothing and use a moisturizing cream or lotion as needed.

Before using Veltin

Although Veltin gel is applied to the skin, your body may absorb enough clindamycin to affect certain medical conditions you may have.

You should not use this medication if you are allergic to clindamycin or tretinoin, or if you have:

- inflammation of your intestines (also called enteritis);
- · Crohn's disease, ulcerative colitis; or
- if you have ever had severe diarrhea caused by antibiotic medicine.

You should not use Veltin if you are also taking another antibiotic called erythromycin, or using other skin medications that contain erythromycin.

To make sure Veltin is safe for you, tell your doctor if you have:

- · any allergies; or
- a history of stomach or intestinal disorders.

FDA pregnancy category C. It is not known whether Veltin will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medication.

It is not known whether clindamycin and tretinoin topical passes into breast milk or if it could harm a nursing baby. You should not breast-feed while using this medicine.

This medicine should not be used on a child younger than 12 years old.

1 Veltin pregnancy and breastfeeding warnings (more detail)

How should I use Veltin?

Use Veltin gel exactly as your doctor has prescribed it for you. Follow all directions on your prescription label. Do not use this medicine in larger or smaller amounts or for longer than recommended. Using more medicine or applying it more often than prescribed will not make it work any faster, and may increase side effects.

Wash your hands before and after applying Veltin gel.

Wash your face with a mild soap or cleanser and pat the skin dry with a clean towel.

Use only a pea-sized amount of Veltin gel to cover the face. Do not apply more often than once daily (at bedtime). More frequent application will NOT make for faster results and may increase irritation.

Do not wash your face more than 3 times daily while using this medication. Washing more often or scrubbing your face when you wash it can make your acne worse.

Avoid getting Veltin gel in your eyes, mouth, or nose (or in the creases of your nose), or on your lips. If it does get into any of these areas, wash with water. Do not apply this medicine to sunburned, windburned, dry, chapped, irritated, or broken skin.

Use this medication for the full prescribed length of time, even if you think it is not working. Call your doctor if skin irritation becomes severe or if your acne does not improve after several weeks of using this medicine.

If you need surgery, tell the surgeon ahead of time that you are using Veltin.

Store Veltin gel at room temperature away from moisture, heat, and light. Do not freeze.

Detailed Veltin dosage information

What happens if I miss a dose?

Use the medication as soon as you remember the missed dose. If it is almost time for your next dose, skip the missed dose and use the medicine at your next regularly scheduled time. Do not use extra medicine to make up the missed dose.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid?

Avoid washing treated skin or applying other skin products for at least 1 hour after applying this medicine.

Avoid using other medications on the areas you treat with Veltin gel unless your doctor tells you to. Avoid using skin products that can cause irritation, such as harsh soaps, shampoos, or skin cleansers, hair coloring or permanent chemicals, hair removers or waxes, or skin products with alcohol, spices, astringents, or lime.

Antibiotic medicines can cause diarrhea, which may be a sign of a new infection. If you have diarrhea that is watery or bloody, stop using this medicine and call your doctor. Do not use anti-diarrhea medicine unless your doctor tells you to.

Veltin can make you sunburn more easily. Apply sunscreen (SPF 30 or higher) every morning and throughout the day as needed. Avoid exposure to sunlight or tanning beds.

Your skin may be more sensitive to weather extremes such as cold and wind. Protect your skin with clothing and use a moisturizing cream or lotion as needed.

Veltin side effects

Get emergency medical help if you have any of these signs of an allergic reaction to Veltin: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Stop using this medicine and call your doctor at once if you have:

- severe redness, burning, stinging, or scaling of treated skin areas; or
- diarrhea that is watery or bloody.

Common Veltin side effects may include:

- mild skin irritation, dryness or peeling;
- mild skin redness; or
- · itching of treated skin.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

1 Veltin side effects (more detail)

Related/similar drugs

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Soolantra

Soolantra (ivermectin) is used to treat rosacea. Includes Soolantra side effects, interactions and ...

Reviews & ratings

7.7 / 10

299 Reviews

View more

Tazorac

Tazorac cream and gel is used to treat acne and plaque psoriasis. Includes Tazorac side effects ...

Reviews & ratings

7.9 / 10

38 Reviews

View more

What other drugs will affect Veltin?

It is not likely that other drugs you take orally or inject will have an effect on topically applied clindamycin and tretinoin. But many drugs can interact with each other. Tell each of your health care providers about all medicines you use, including prescription and over-the-counter medicines, vitamins, and herbal products.



Does Veltin interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Veltin	
+	
Enter a drug name	Add

More about Veltin (clindamycin / tretinoin topical)

- · Check interactions
- Compare alternatives
- Reviews (42)
- · Side effects
- Dosage information
- During pregnancy
- · Generic availability
- FDA approval history
- · Drug class: topical acne agents
- En español

Patient resources

Other brands

Ziana

Professional resources

· Veltin Gel prescribing information

Other brands

Ziana

Related treatment guides

Acne

Further information

• Your pharmacist has information about Veltin written for health professionals that you may read.

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and

use Veltin gel only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 2.01.

Antibiotics 101

List of Common Antibiotics & Types

Antibiotics and Alcohol

Probiotics and Antibiotics

Antibiotics for Strep Throat

Antibiotics for a Tooth Infection

Antibiotics for UTI's

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

7.4 / 10

42 Reviews

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





Advertising policy
Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

×