

Home

2. Haloperidol

Haloperidol ⊲[∞]

Generic name: haloperidol (oral) [HAL-oh-PER-i-dol]

Brand name: Haldol

Dosage forms: oral concentrate (2 mg/mL), oral tablet (0.5 mg; 1 mg; 10 mg; 2 mg; 20 mg; 5 mg)

Drug class: Miscellaneous antipsychotic agents



Medically reviewed by Drugs.com on Aug 4, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions FAQ

What is haloperidol?

Haloperidol is an antipsychotic medicine that is used to treat schizophrenia.

Haloperidol is also used to control motor and speech tics in people with Tourette's syndrome.

Haloperidol may also be used for purposes not listed in this medication guide.

Haloperidol side effects

Get emergency medical help if you have **signs of an allergic reaction**: hives; difficult breathing; swelling of your face, lips, tongue, or throat.

High doses or long-term use of haloperidol can cause a serious movement disorder that may not be reversible.

The longer you use haloperidol, the more likely you are to develop this disorder, especially if you are a woman or an older adult.

Haloperidol may cause serious side effects. Call your doctor at once if you have:

- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement);
- muscle spasms in your neck, tightness in your throat, trouble swallowing;
- rapid changes in mood or behavior;
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out);
- cough with mucus, chest pain, feeling short of breath;
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat, cough, trouble breathing; or
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven

heartbeats, tremors, feeling like you might pass out.

Common side effects of haloperidol may include:

- · drowsiness;
- · headache;
- · dizziness, spinning sensation;
- uncontrolled muscle movements;
- · feeling restless or anxious;
- sleep problems (insomnia); or
- breast enlargement, irregular menstrual periods.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

• Haloperidol side effects (more detail)

Related/similar drugs

Abilify

Abilify is an antipsychotic medicine used to treat the symptoms of schizophrenia and bipolar ...

Reviews & ratings

6.1 / 10

1,187 Reviews

View more

Diazepam

Diazepam is used to treat anxiety disorders, alcohol withdrawal symptoms, or muscle spasms. Learn

Reviews & ratings

8.4 / 10

884 Reviews

View more

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment, weight

management, cardiovascular ...

Reviews & ratings

6.6 / 10

1,504 Reviews

View more

Olanzapine

Olanzapine (brand name Zyprexa) is an atypical antipsychotic that may be used to treat adults and ...

Reviews & ratings

5.9 / 10

1,091 Reviews

View more

Dexamethasone

Dexamethasone is used to treat inflammatory conditions such as allergies, skin conditions ...

Reviews & ratings

5.9 / 10

162 Reviews

View more

Lorazepam

Lorazepam is in a group of drugs called benzodiazepines and is used to treat anxiety disorders ...

Reviews & ratings

7.9 / 10

1,054 Reviews

View more

Aripiprazole

Easy-to-read patient leaflet for aripiprazole. Includes indications, proper use, special ...

Reviews & ratings

1,540 Reviews

View more

Risperidone

Risperidone is used to treat schizophrenia and symptoms of bipolar disorder. Learn about side ...

Reviews & ratings

5.6 / 10

731 Reviews

View more

Promethazine

Promethazine is a phenothiazine and antihistamine used to treat allergies, motion sickness, nausea ...

Reviews & ratings

7.2 / 10

388 Reviews

View more

Zofran

Zofran (ondansetron) is used to prevent nausea and vomiting caused by surgery or cancer medicines ...

Reviews & ratings

8.0 / 10

270 Reviews

View more

Warnings

Haloperidol is not approved for use in older adults with dementia-related psychosis.

Before taking this medicine

You should not use haloperidol if you are allergic to it, or if you have:

- · Parkinson's disease; or
- certain conditions that affect your central nervous system (such as severe drowsiness, or slowed thinking caused by taking other medicines or drinking alcohol).

Haloperidol may increase the risk of death in older adults with dementia-related psychosis and is not approved for this use.

Tell your doctor if you have ever had:

- heart problems, angina (chest pain);
- long QT syndrome (in you or a family member);
- low blood pressure;
- · a seizure;
- a thyroid disorder;
- breast cancer; or
- an electrolyte imbalance (such as low blood levels of potassium or magnesium).

Using antipsychotic medicine in the last 3 months of pregnancy may cause serious problems in the newborn. If you get pregnant, tell your doctor right away. Do not stop the medicine without your doctor's advice.

Ask a doctor if it is safe to breastfeed while using haloperidol.

How should I take haloperidol?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Your doctor may occasionally change your dose. Use the medicine exactly as directed.

Taking too much haloperidol can cause a serious heart rhythm disorder or sudden death. **Never take more than your prescribed dose.**

Measure liquid medicine with the supplied syringe or a dose-measuring device (not a kitchen spoon).

Your symptoms may not improve for several weeks.

You may have withdrawal symptoms if you stop using haloperidol suddenly. Ask your doctor before stopping the medicine.

Store at room temperature away from moisture, heat, and light. Do not allow liquid medicine to freeze.

Detailed Haloperidol dosage information.

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222. **An overdose of haloperidol can be fatal.**

What should I avoid while taking haloperidol?

Drinking alcohol with haloperidol can cause side effects.

Avoid driving or hazardous activity until you know how this medicine will affect you. Your reactions could be impaired. Avoid getting up too fast from a sitting or lying position, or you may feel dizzy. Dizziness or severe drowsiness can cause falls, fractures, or other injuries.

What other drugs will affect haloperidol?

Haloperidol can cause a serious heart problem. Your risk may be higher if you also use certain other medicines for infections, asthma, heart problems, high blood pressure, depression, mental illness, cancer, malaria, or HIV.

Using haloperidol with other drugs that make you drowsy can worsen this effect. Ask your doctor before using opioid medication, a sleeping pill, a muscle relaxer, or medicine for anxiety or seizures.

Tell your doctor about all your other medicines, especially:

- lithium;
- rifampin;
- medicine to treat Parkinson's disease;
- seizure medicine; or
- a blood thinner--warfarin, Coumadin, Jantoven.

This list is not complete. Other drugs may affect haloperidol, including prescription and over-the-counter medicines, vitamins, and herbal products. Not all possible drug interactions are listed here.

<u>Haloperidol drug interactions</u> (more detail)

Does haloperidol interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

haloperidol	
+	
Enter a drug name	Add

Frequently asked questions

What drugs cause tardive dyskinesia?

More about haloperidol

- · Check interactions
- Compare alternatives
- Pricing & coupons
- Reviews (203)
- Drug images
- · Side effects
- Dosage information
- Patient tips
- During pregnancy
- Support group
- Drug class: miscellaneous antipsychotic agents
- Breastfeeding

Patient resources

- Haloperidol injection drug information
- Haloperidol Decanoate Injection
- Haloperidol Oral Solution

Other brands

Haldol, Haldol Decanoate

Professional resources

- Haloperidol monograph
- Haloperidol (FDA)
- Haloperidol Decanoate Injection (FDA)
- Haloperidol Lactate Injection (FDA)
- Haloperidol Oral Solution (FDA)

Other brands

Haldol, Haldol Decanoate

Related treatment guides

- Aggressive Behavior
- Dementia
- ICU Agitation

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 17.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

ন্ত Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

Drug history at FDA

User Reviews & Ratings

4.4 / 10

203 Reviews

Images

Haloperidol 5 mg (ZC 07)





View larger images

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

About Drugs.com

Advertising policy

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us











Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

×