

#### Home

Alka-Seltzer Plus Cold

### Alka-Seltzer Plus Cold

Generic name: acetaminophen, chlorpheniramine, and phenylephrine [ a-SEET-a-MIN-oh-fen, KLOR-fen-EER-a-meen, FEN-ill-EFF-rin ]

Brand names: Comtrex Flu Therapy, Comtrex Severe Cold and Sinus, Flonase Headache and Allergy Relief, Norel AD, Relief-PE, ... show all 32 brands

Drug class: Upper respiratory combinations



Medically reviewed by Drugs.com on Jul 26, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions



The **Alka-Seltzer Plus Cold** product containing acetaminophen, chlorpheniramine, and phenylephrine has been discontinued in the U.S.

A similar, reformulated Alka-Seltzer Plus Cold (aspirin, chlorpheniramine, and phenylephrine) is currently available.

### What is Alka-Seltzer Plus Cold?

Acetaminophen is a pain reliever and fever reducer.

Chlorpheniramine is an antihistamine that reduces the effects of natural chemical histamine in the body. Histamine can produce symptoms of sneezing, itching, watery eyes, and runny nose.

Phenylephrine is a decongestant that shrinks blood vessels in the nasal passages. Dilated blood vessels can cause nasal congestion (stuffy nose).

Alka-Seltzer Plus Cold is a combination medicine used to treat headache, fever, body aches, runny or stuffy nose, sneezing, itching, watery eyes, and sinus congestion caused by allergies, the common cold, or the flu.

Alka-Seltzer Plus Cold may also be used for purposes not listed in this medication guide.

### Alka-Seltzer Plus Cold side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Alka-Seltzer Plus Cold and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen.

### This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have:

- chest pain, rapid pulse, fast or uneven heart rate;
- confusion, hallucinations, severe nervousness;
- tremor, seizure (convulsions);
- easy bruising or bleeding, unusual weakness;
- · little or no urinating;
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes); or
- dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, chest pain, shortness of breath, uneven heartbeats, seizure).

Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

### Common side effects of Alka-Seltzer Plus Cold may include:

- dizziness, drowsiness;
- mild headache, blurred vision;
- dry mouth, nose, or throat;
- · constipation;
- · feeling nervous; or
- sleep problems (insomnia);

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Alka-Seltzer Plus Cold side effects (more detail)

### Related/similar drugs

### **Benadryl**

Benadryl is an antihistamine and is used to treat sneezing, runny nose, itching and other ...

### **Reviews & ratings**

7.0 / 10

79 Reviews

View more

#### FEATURED

#### **Omvoh**

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

#### **Reviews & ratings**

1 Reviews

View more

### **Nasonex 24HR Allergy**

Nasonex 24HR Allergy (mometasone) is used to treat nasal allergies such as congestion, sneezing ...

#### **Reviews & ratings**

Add a review

View more

### Warnings

Do not use this medicine if you have used an MAO inhibitor in the past 14 days, such as isocarboxazid, linezolid, methylene blue injection, phenelzine, rasagiline, selegiline, or tranylcypromine.

Do not take more of this medication than is recommended. **An overdose of acetaminophen can damage your liver or cause death.** Call your doctor at once if you have nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, or jaundice (yellowing of your skin or eyes).

In rare cases, acetaminophen may cause a severe skin reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling.

### Before taking this medicine

You should not use this medication if you are allergic to acetaminophen, chlorpheniramine, or phenylephrine.

Do not use this medicine if you have taken an MAO inhibitor in the past 14 days. A dangerous drug interaction could occur. MAO inhibitors include isocarboxazid, linezolid, phenelzine, rasagiline, selegiline, and tranylcypromine.

Ask a doctor or pharmacist if it is safe for you to take this medicine if you have other medical conditions, especially:

- asthma or COPD, cough with mucus, or cough caused by smoking, emphysema, or chronic bronchitis;
- a blockage in your stomach or intestines;
- liver disease, alcoholism, or if you drink more than 3 alcoholic beverages per day;
- · kidney disease;
- high blood pressure, heart disease, coronary artery disease, or recent heart attack;
- enlarged prostate or urination problems;
- glaucoma;
- · diabetes:
- · overactive thyroid;
- pheochromocytoma (an adrenal gland tumor); or
- if you take potassium (Cytra, Epiklor, K-Lyte, K-Phos, Kaon, Klor-Con, Polycitra, Urocit-K).

It is not known whether Alka-Seltzer Plus Cold will harm an unborn baby. **Do not use this medicine without your doctor's advice if you are pregnant.** 

This medication may pass into breast milk and may harm a nursing baby. Antihistamines and decongestants may also slow breast milk production. Do not use this medicine without your doctor's advice if you are breast-feeding a baby.

### How should I take Alka-Seltzer Plus Cold?

Use exactly as directed on the label, or as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. This medicine is usually taken only for a short time until your symptoms clear up.

Do not take more of this medication than is recommended. **An overdose of acetaminophen can damage your liver or cause death.** 

Measure liquid medicine with a special dose-measuring spoon or medicine cup, not with a regular table spoon. If you do not have a dose-measuring device, ask your pharmacist for one.

Drop the effervescent tablets into a glass of water (at least 4 ounces, or one-half cup). Stir this mixture and drink all of it right away.

Do not give this medication to a child younger than 4 years old. Always ask a doctor before giving a cough or cold medicine to a child. Death can occur from the misuse of cough and cold medicines in very young children.

**Do not take for longer than 7 days in a row.** Stop taking the medicine and call your doctor if you still have a fever after 3 days of use, you still have pain after 7 days (or 5 days if treating a child), if your symptoms get worse, or if you have a skin rash, ongoing headache, or any redness or swelling.

If you need surgery or medical tests, tell the surgeon or doctor ahead of time if you have taken this medicine within the

past few days.

Store at room temperature away from moisture and heat. Do not allow liquid medicine to freeze.

Detailed Alka-Seltzer Plus Cold dosage information

### What happens if I miss a dose?

Since Alka-Seltzer Plus Cold is taken when needed, you may not be on a dosing schedule. If you are taking the medication regularly, take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. **Do not** take extra medicine to make up the missed dose.

### What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1 800 222 1222. **An overdose of acetaminophen can be fatal.** 

The first signs of an acetaminophen overdose include loss of appetite, nausea, vomiting, stomach pain, sweating, and confusion or weakness. Later symptoms may include pain in your upper stomach, dark urine, and yellowing of your skin or the whites of your eyes.

### What should I avoid while taking Alka-Seltzer Plus Cold?

Ask a doctor or pharmacist before using any other cold, allergy, pain, or sleep medication. Acetaminophen (sometimes abbreviated as APAP) is contained in many combination medicines. **Taking certain products together can cause you to get too much acetaminophen which can lead to a fatal overdose.** Check the label to see if a medicine contains acetaminophen or APAP.

Avoid drinking alcohol. It may increase your risk of liver damage while taking acetaminophen, and can increase certain side effects of chlorpheniramine.

This medicine may cause blurred vision or impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert and able to see clearly.

### What other drugs will affect Alka-Seltzer Plus Cold?

Ask a doctor or pharmacist before using Alka-Seltzer Plus Cold if you are also using any other drugs, including prescription and over-the-counter medicines, vitamins, and herbal products. Some medicines can cause unwanted or dangerous effects when used together. Not all possible interactions are listed in this medication guide.

Taking this medicine with other drugs that make you sleepy or slow your breathing can worsen these effects. Ask your doctor before taking Alka-Seltzer Plus Cold with a sleeping pill, narcotic pain medicine, muscle relaxer, or medicine for anxiety, depression, or seizures.

Alka-Seltzer Plus Cold drug interactions (more detail)

Enter medications to view a detailed interaction report using our Drug Interaction Checker.

Alka-Seltzer Plus Cold	
+	
Enter a drug name	Add

# More about Alka-Seltzer Plus Cold (acetaminophen / chlorpheniramine / phenylephrine)

- · Check interactions
- · Compare alternatives
- Reviews (9)
- Drug images
- Side effects
- Dosage information
- Drug class: upper respiratory combinations

### **Patient resources**

### Other brands

Norel AD, Comtrex Flu Therapy, Medicidin-D, Relief-PE, ... +9 more

### Related treatment guides

- Cold Symptoms
- Sinus Symptoms

### **Further information**

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

#### Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 11.01.

#### **DRUG STATUS**

**Availability** 



## CSA Schedule\* N/A Not a controlled drug

### **User Reviews & Ratings**

8.3 / 10

9 Reviews

### **Images**

Alka-Seltzer Plus Cold aspirin 325 mg / chlorpheniramine maleate 2 mg / phenylephrine bitartrate 7.8 mg (ALKA SELTZER +)

### **Drugs.com Mobile App**

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





### **About**

About Drugs.com

Advertising policy

Attribution & citations

#### Terms & privacy

Terms of use

**Editorial policy** 

Privacy policy

### **Support**

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

**Drugs.com** provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









