

[Home](#)

2. Bazedoxifene/conjugated estrogens

Bazedoxifene and conjugated estrogens

Generic name: bazedoxifene and conjugated estrogens [*BA-ze-DOX-i-feen-and-KON-joo-GAY-ted-ES-troe-jens*]

Brand name: [Duavee](#)

Dosage form: oral tablet (20 mg-0.45 mg)

Drug class: [Sex hormone combinations](#)



[Medically reviewed](#) by Drugs.com on Feb 6, 2025. Written by [Cerner Multum](#).

[Uses](#) [Side effects](#) [Warnings](#) [Before taking](#) [Dosage](#) [Interactions](#)

What is bazedoxifene and conjugated estrogens?

Bazedoxifene blocks estrogen from reaching certain cells in the uterine lining, to protect against overgrowth of uterine tissue.

Estrogen is a female sex hormone produced by the ovaries. [Conjugated estrogens](#) are a mixture of estrogen hormones.

Bazedoxifene and conjugated estrogens is a combination medicine used short-term in women after menopause to treat moderate to severe [hot flashes](#), and to prevent [osteoporosis](#). **bazedoxifene and conjugated estrogens is used in women who have NOT had a hysterectomy (uterus removed).**

Bazedoxifene and conjugated estrogens may also be used for purposes not listed in this medication guide.

Bazedoxifene and conjugated estrogens side effects

Get emergency medical help if you have **signs of an allergic reaction:** hives; difficult breathing; swelling of your face, lips, tongue, or throat.

Bazedoxifene and conjugated estrogens may cause serious side effects. Call your doctor at once if you have:

- **heart attack symptoms**--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating;
- **signs of a stroke**--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance;
- **signs of a blood clot**--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, pain or warmth in one or both legs;
- sudden vision loss;
- swelling, rapid weight gain;
- jaundice (yellowing of the skin or eyes);

- memory problems, confusion, unusual behavior;
- unusual vaginal bleeding, pelvic pain;
- a lump in your breast; or
- **low calcium level**--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes).

Common side effects of bazedoxifene and conjugated estrogens may include:

- dizziness;
- nausea, stomach pain or discomfort, diarrhea;
- [muscle spasm](#); or
- throat pain, neck pain.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

 [Bazedoxifene / conjugated estrogens side effects](#) (more detail)

Related/similar drugs

Paxil

Paxil is used to treat depression, obsessive-compulsive disorder and anxiety. Learn about side ...

Reviews & ratings

7.1 / 10

[960 Reviews](#)

[View more](#)

Prolia

Prolia (denosumab) is an injection that is administered subcutaneously (under the skin) once every ...

Reviews & ratings

2.9 / 10

[374 Reviews](#)

[View more](#)

FEATURED

Ozempic

Learn about Ozempic (semaglutide) for type 2 diabetes treatment. weight

for type 2 diabetes treatment, weight management, cardiovascular ...

Reviews & ratings

6.6 / 10

[1,504 Reviews](#)

[View more](#)

Fosamax

Fosamax (alendronate) is used to treat or prevent postmenopausal osteoporosis and steroid-induced ...

Reviews & ratings

2.8 / 10

[76 Reviews](#)

[View more](#)

Premarin

Premarin is used to treat symptoms of menopause such as hot flashes, and vaginal dryness. Learn ...

Reviews & ratings

7.7 / 10

[84 Reviews](#)

[View more](#)

Reclast

Reclast (zoledronic acid) is used to treat osteoporosis in women after menopause and also to treat ...

Reviews & ratings

5.1 / 10

[275 Reviews](#)

[View more](#)

Estrace

Estrace (estradiol) is used to treat symptoms of menopause such as hot flashes and vaginal dryness ...

Reviews & ratings

[View more](#)

Zometa

Zometa is used to treat high blood levels of calcium caused by cancer and multiple myeloma. Learn ...

Reviews & ratings

4.8 / 10

[43 Reviews](#)[View more](#)

Warnings

You should not use bazedoxifene and conjugated estrogens if you have: undiagnosed vaginal bleeding, [liver disease](#), a [bleeding disorder](#), if you will have major surgery, or if you have ever had a [heart attack](#), a stroke, a blood clot, or cancer of the breast, uterus/cervix, or vagina.

Using this medicine can increase your risk of blood clots, stroke, heart attack, or cancer of the breast, uterus, or ovaries. Estrogens should not be used to prevent [heart disease](#), stroke, or dementia.

Estrogens may increase your risk of developing a condition that may lead to [uterine cancer](#). **Report any unusual vaginal bleeding right away.**

Bazedoxifene and conjugated estrogens is for short-term use at the lowest dose needed to treat your condition.

Do not take any other estrogens while you are taking this medicine.

Before taking this medicine

You should not use this medicine if you are allergic to bazedoxifene or conjugated estrogens, or if you have:

- unusual vaginal bleeding that has not been checked by a doctor;
- liver disease;
- a history of heart attack, stroke, or blood clot;
- an increased risk of having blood clots due to a heart problem or a hereditary blood disorder; or
- a history of hormone-related cancer, or cancer of the breast, uterus/cervix, or vagina.

Do not use other estrogen products while you are taking bazedoxifene and conjugated estrogens.

Using this medicine can increase your risk of blood clots, stroke, or heart attack. You are even more at risk if you have [high blood pressure](#), diabetes, [high cholesterol](#), if you are [overweight](#), or if you smoke.

Bazedoxifene and conjugated estrogens should not be used to prevent heart disease, stroke, or dementia, because this medicine may actually increase your risk of developing these conditions.

Women who are 75 or older should not take bazedoxifene and conjugated estrogens.

Tell your doctor if you have ever had:

- a [hysterectomy](#);
- heart disease;
- liver problems, or jaundice caused by pregnancy or taking hormones;
- kidney disease;
- [gallbladder disease](#);
- asthma;
- epilepsy or other seizure disorder;
- migraines;
- lupus;
- [endometriosis](#) or uterine fibroid tumors;
- [hereditary angioedema](#);
- porphyria (a genetic enzyme disorder that causes symptoms affecting the skin or nervous system);
- a thyroid disorder; or
- high levels of calcium in your blood.

Using bazedoxifene and conjugated estrogens may increase your risk of cancer of the breast, uterus, or ovaries. Talk with your doctor about this risk.

This medicine is for use only in women who can no longer get pregnant. **Bazedoxifene and conjugated estrogens can harm an unborn baby. Do not use if you are pregnant.** Use effective [birth control](#) if you are not past menopause. Tell your doctor if you think you may be pregnant.

You should not breast-feed while using this medicine.

How should I take bazedoxifene and conjugated estrogens?

Follow all directions on your prescription label and read all medication guides or instruction sheets. Use the medicine exactly as directed.

Bazedoxifene and conjugated estrogens is for short-term use at the lowest dose needed to treat your condition.

You may take bazedoxifene and conjugated estrogens with or without food, at the same time each day.

Swallow the tablet whole and do not crush, chew, or break it.

Your doctor should check your progress on a regular basis to determine whether you should continue this treatment. Self-examine your breasts for lumps on a monthly basis, and have regular mammograms.

If you need major surgery or will be on long-term bed rest, you may need to stop using this medicine for a short time. Any doctor or surgeon who treats you should know that you are using bazedoxifene and conjugated estrogens.

Your doctor may have you take extra calcium or vitamin D while you are taking bazedoxifene and conjugated estrogens. Take only the amount that your doctor has prescribed.

Store at room temperature away from moisture and heat. Keep each tablet in its blister pack until you are ready to take it. Do not use a pill box for this medicine.

Write down the date you open a Duavee foil pouch. Use the medicine within 60 days after opening the pouch.

i [Detailed Bazedoxifene / conjugated estrogens dosage information](#)

What happens if I miss a dose?

Take the medicine as soon as you can, but skip the missed dose if it is almost time for your next dose. **Do not** take two doses at one time.

What happens if I overdose?

Seek emergency medical attention or call the Poison Help line at 1-800-222-1222.

What should I avoid while taking bazedoxifene and conjugated estrogens?

Do not take progestins or any other estrogen while you are taking bazedoxifene and conjugated estrogens.

What other drugs will affect bazedoxifene and conjugated estrogens?

Sometimes it is not safe to use certain medications at the same time. Some drugs can affect your blood levels of other drugs you take, which may increase side effects or make the medications less effective.

Many drugs can affect bazedoxifene and conjugated estrogens. This includes prescription and over-the-counter medicines, vitamins, and [herbal products](#). **Not all possible interactions are listed here.** Tell your doctor about all your current medicines and any medicine you start or stop using.

i [Bazedoxifene / conjugated estrogens drug interactions](#) (more detail)

Does bazedoxifene/conjugated estrogens interact with my other drugs?

Enter medications to view a detailed interaction report using our [Drug Interaction Checker](#).

bazedoxifene/conjugated estrogens

+

Enter a drug name

Add

More about bazedoxifene / conjugated estrogens

- [Check interactions](#)
- [Compare alternatives](#)
- [Reviews \(35\)](#)
- [Side effects](#)
- [Dosage information](#)
- [During pregnancy](#)
- [Drug class: sex hormone combinations](#)
- [En español](#)

Patient resources

- [Conjugated estrogens and bazedoxifene advanced reading](#)

Other brands

[Duavee](#)

Professional resources

- [Bazedoxifene monograph](#)

Other brands

[Duavee](#)

Related treatment guides

- [Hot Flashes](#)
- [Osteoporosis](#)
- [Postmenopausal Symptoms](#)
- [Prevention of Osteoporosis](#)

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

[Medical Disclaimer](#)

Copyright 1996-2025 Cerner Multum, Inc. Version: 2.03.

DRUG STATUS

Availability

Rx Prescription only


Pregnancy & Lactation

 Risk data available


CSA Schedule*

N/A Not a controlled drug

Approval History

 Drug history at FDA

WADA Class

 Anti-Doping Classification



User Reviews & Ratings

6.7 / 10

[35 Reviews](#)

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.



About

[About Drugs.com](#)

[Advertising policy](#)

[Attribution & citations](#)

Terms & privacy

[Terms of use](#)

[Editorial policy](#)

[Privacy policy](#)

Support

[Help center](#)

[Sitemap](#)

[Contact us](#)



[Subscribe to our newsletter](#) for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment.

Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.



Copyright © 2000-2025 Drugs.com. All rights reserved.