

Home

Hydroxocobalamin

Hydroxocobalamin (injection)

Generic name: hydroxocobalamin (injection) [hye-DROX-o-koe-BAL-a-min]

Brand names: Cyanokit, Hydroxy-Cobal, Hydro-Cobex, Cobalin-H, Neo-Cytamen

Dosage forms: injectable solution (1000 mcg/mL), intravenous powder for injection (5 g)

Drug class: Vitamins



Medically reviewed by Drugs.com on Aug 14, 2023. Written by Cerner Multum.

Uses Side effects Warnings Before taking Dosage Interactions

What is hydroxocobalamin?

Hydroxocobalamin is a form of vitamin B-12. It is used as an antidote to cyanide poisoning. Hydroxocobalamin works by helping cells in the body convert cyanide to a form that can be removed from the body through urination.

Hydroxocobalamin is used in an emergency to treat cyanide poisoning. This type of poisoning can occur if you are exposed to smoke from a house or industrial fire, if you swallow or breathe in cyanide, or if you get cyanide on your skin.

Hydroxocobalamin may also be used for purposes not listed in this medication guide.

Hydroxocobalamin side effects

Get emergency medical help if you have any of these **signs of an allergic reaction:** hives; chest tightness, difficulty breathing; swelling of your face, lips, tongue, or throat.

You may develop an acne-like skin rash within 1 to 4 weeks after you are treated with hydroxocobalamin. This rash should go away without treatment. Call your doctor if you have a rash that lasts longer than 4 weeks.

Call your doctor at once if you have a serious side effect such as:

- bright red blood in your stools;
- chest pain, fast or uneven heart rate;
- severe shortness of breath, wheezing, gasping for breath, cough with foamy mucus;
- · swelling in your feet or ankles;
- · a light-headed feeling, like you might pass out; or
- dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure).

Common side effects of Hydroxocobalamin may include:

- nausea;
- · headache;
- acne, skin rash or redness;
- red coloring of your urine (may last 2 to 5 weeks); or
- pain, swelling, or irritation of your skin where the injection was given.

This is not a complete list of side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Hydroxocobalamin side effects (more detail)

Related/similar drugs

FEATURED

Omvoh

Omvoh is used to treat moderate to severe ulcerative colitis or Crohn's disease in adults. This ...

Reviews & ratings

1 Reviews

View more

Cyanokit

Cyanokit is used in an emergency to treat cyanide poisoning. Learn about side effects, interactions ...

Reviews & ratings

Add a review

View more

Warnings

In an emergency situation it may not be possible to tell your caregivers about your health conditions. Make sure any doctor caring for you afterward knows you have received Hydroxocobalamin.

Before taking this medicine

If possible before you receive hydroxocobalamin, tell your caregivers if you have:

• high blood pressure;

- heart disease;
- · congestive heart failure;
- liver disease;
- · kidney disease (or if you are on dialysis); or
- if you have ever had a severe allergic reaction to hydroxocobalamin, Vitamin B12, or cyanocobalamin (Nascobal, Cobolin, Cyomin, and others).

FDA pregnancy category C. Tell your doctor if you are pregnant. It is not known whether hydroxocobalamin will harm an unborn baby. However, the benefits of treating cyanide poisoning may outweigh any risks posed by hydroxocobalamin, for both you and your baby.

It is not known whether hydroxocobalamin passes into breast milk or if it could harm a nursing baby. You should not breast-feed after you have been treated with hydroxocobalamin.

In an emergency situation, it may not be possible before you are treated with hydroxocobalamin to tell your caregivers if you are pregnant or breast feeding. Make sure any doctor caring for your pregnancy or your baby knows you have received this medication.

How should I take hydroxocobalamin?

Hydroxocobalamin is injected into a vein through an IV. You will receive this injection in a clinic or hospital setting. Hydroxocobalamin must be given slowly, and the IV infusion can take at least 15 minutes to complete.

Hydroxocobalamin is usually given only once. However, you may receive a second dose if needed.

Your breathing, blood pressure, oxygen levels, heart function, and other vital signs will be watched closely while you are receiving hydroxocobalamin.

This medication can cause unusual results with certain medical tests. Tell any doctor who treats you that you have recently received a hydroxocobalamin injection.

1 Detailed Hydroxocobalamin dosage information

What happens if I miss a dose?

Since hydroxocobalamin is given by a healthcare professional in an emergency setting, you are not likely to miss a dose.

What happens if I overdose?

Since this medication is given by a healthcare professional in a medical setting, an overdose is unlikely to occur.

What should I avoid while taking hydroxocobalamin?

If your skin turns red after receiving this medication, avoid exposure to sunlight or tanning beds. Hydroxocobalamin can make you sunburn more easily while your skin is still red. Wear protective clothing and use sunscreen (SPF 30 or higher) until your skin color returns to normal.

What other drugs will affect hydroxocobalamin?

Other drugs may interact with hydroxocobalamin, including prescription and over-the-counter medicines, vitamins, and herbal products. Tell each of your health care providers about all medicines you use now and any medicine you start or stop using.

Hydroxocobalamin drug interactions (more detail)

Does hydroxocobalamin interact with my other drugs?

Enter medications to view a detailed interaction report using our Drug Interaction Checker.



More about hydroxocobalamin

- · Check interactions
- Compare alternatives
- Reviews (2)
- Latest FDA alerts (1)
- · Side effects
- Dosage information
- During pregnancy
- Drug class: vitamins
- En español

Patient resources

- Hydroxocobalamin advanced reading
- Hydroxocobalamin Injection (Cyanide Poisoning)

Other brands

Cyanokit

Professional resources

• Hydroxocobalamin prescribing information

Other brands

Cyanokit

Related treatment guides

- Vitamin B12 Deficiency
- · Cyanide Poisoning
- Transcobalamin II Deficiency
- Schilling Test

Further information

Remember, keep this and all other medicines out of the reach of children, never share your medicines with others, and use this medication only for the indication prescribed.

Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer

Copyright 1996-2025 Cerner Multum, Inc. Version: 5.01.

DRUG STATUS

Availability

Rx Prescription only

Pregnancy & Lactation

& Risk data available

CSA Schedule*

N/A Not a controlled drug

Approval History

□ Drug history at FDA

User Reviews & Ratings

Review this drug

Drugs.com Mobile App

Access drug & treatment information, identify pills, check interactions and set up personal medication records.





About

Attribution & citations

Terms & privacy

Terms of use

Editorial policy

Privacy policy

Support

Help center

Sitemap

Contact us













Subscribe to our newsletter for the latest medication news, new drug approvals and FDA alerts.

Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include Micromedex (updated 7 Apr 2025), Cerner Multum™ (updated 13 Apr 2025), ASHP (updated 10 Apr 2025) and others.









Copyright © 2000-2025 Drugs.com. All rights reserved.

×